VOL. 29. Ser. A. Title-page and Index.

[Issued August' 1942.]

THE REVIEW OF APPLIED ENTOMOLOGY.

SERIES A: AGRICULTURAL.

ISSUED BY THE IMPERIAL INSTITUTE OF ENTOMOLOGY.

LONDON: THE IMPERIAL INSTITUTE OF ENTOMOLOGY 41. QUEEN'S GATE, S.W. 7.

Price 6s. net.

All Rights Reserved.

IMPERIAL INSTITUTE OF ENTOMOLOGY.

Executive Council.

SHAMALDHARI LALL, C.I.E., Officiating High Commissioner, Chairman, India and Burma.

J. A. CALDER, C.M.G., Vice-Chairman, Colonies, Protectorates and Mandated Territories.

Sir Donald Fergusson, K.C.B., United Kingdom.

F. L. McDougall, C.M.G., Australia.

F. J. DU TOIT, Union of South Africa.

NEVILL L. WRIGHT, F.I.C., D.I.C., New Zealand.

J. M. ADAMS, F.R.C.Sc. (I), Eire.

D. James Davies, C.B.E., Newfoundland.

W. C. ROBERTSON, Southern Rhodesia.

Sir David Chadwick, K.C.M.G., C.S.I., C.I.E., Secretary.

Director and Editor.

Dr. S. A. NEAVE, C.M.G., O.B.E.

Assistant Director.

Dr. W. R. THOMPSON, F.R.S.

Head Office—British Museum (Natural History), Cromwell Road, London, S.W.7.

Publication Office and Library-41, Queen's Gate, London, S.W.7.

INDEX OF AUTHORS.

A reference in heavy type indicates that a paper by the author has been noticed.

Abe, T., 102. Abercrombie, W. F., 224. Adamson, A. M., 543. Afanas'eva, O. V., 581. Afzal Husain, M., 266, 272, 341, 516. Agacino, E. Morales, 462. Ahmad, T., 267, 341. Albrecht, H. R., 404. Alden, C. H., 192. Alekseev, Ya. A., 582. Alexandrov, V. R., 240. Allen, H. W., 130, 294. Allen, N., 404, 596. Allen, T. C., 135, 246, 301. Allman, S. L., 54, 337, 494. Amaral, S. F. do, 511. Anderson, H. W., 224. Anderson, L. D., 46, 121, 152, 249, 503, 504. Andison, H., 417, 508. Andreeva, N. V., 583. Andrewartha, H. G., 540. Annand, P. N., 204. Apple, J. W., 556. Arant, F. S., 404. Araujo, R. L., 292. Ark, P. A., 535. Armitage, R. W., 290, 546. Armstrong, T., 225, 306. Arrow, G. J., 388. Ashworth, J. T., 161. Autuori, M., 40, 344. Avens, A. W., 412. Ayyar, P. N. Krishna, 268, 270, 515, 521. Ayyar, T. V. Ramakrishna, 319,

Back, E. A., **137**, **536**. Baerg, W. J., **2**, **3**. Bailey, S. F., **210**. Bailey, V. A., 397, 491. Bailey, W. K., **261**. Baird, A. B., **307**.

480.

Baker, H., 633. Balch, R. E., 37, 347, 531. Barber, G. W., 217, 367, 535. Bare, C. O., 199. Barnes, H. F., 312, 375, 524. Bartlett, B. R., **561.** Basheer, M., **518.** Baumhofer, L. G., 636. Baur, K. E., 418. Beard, R. L., 120, 162, 163, 442. Beaulieu, A. A., 224. Becker, G., 233. Becnel, I. J., **599**, **602**. Bedford, E. C. G., 320. Bedford, H. W., 386. Beeson, C. F. C., 74, 289. Bei-Bienko, G. Ya., 655. Belanovs'kiĭ, I. D., 618. Bell, A. F., 388. Benlloch, M., 457, 459, 463. Bennett, S. H., 110. Benson, R. B., 544. Bentley, E. W., 527. Berezina, V. M., 571. Berg, V. L., 280. Bertholf, L. M., 550. Betrem, J. G., 88. Bhatia, D. R., 519. Bianchi, F. A., 272, 273. Bibby, F. F., 510. Bienko, G. Ya. Bei-, 655. Bigger, J. H., 187, 245. Bishop, H. J., 435. Bissell, T. L., 159, 405, 598. Black, L. M., 500, 502. Blair, C. R., **356.** Blair, K. G., 343. Blake, C. H., 499. Blanchard, E. E., 169, 344. Blanchard, R. A., 187. Blanton, F. S., 415. Bliss, C. I., 352, 433. Bloxham, H. P., 407. Bobinskaya, S. G., 572. Bodine, J. H., 328. Bogachev, A. V., 235.

Bogdarina, A. A., 585. Bohart, R. M., 408. Bohorquez, R., 462. Bondar, G., 9, 391, 468. Bonnemaison, L., 24. Bosq, J. M., 512. Bottger, G. T., 43. Boughton, D. C., 220. Bovey, P., **66**, **617**. Bovien, P., 369. Boyce, A. M., 295, 349, 561. Boyce, H. R., 51. Brain, C. K., 84. Brair, J. H., 622, 623. Brammanis, L., 85. Breakey, E. P., 418. Bredo, H. J., 492, 610. Bremer, H., 231. Briceño-Iragorry, L., 292. Brierley, P., **532.** Brigham, W. T., **161.** Brimblecombe, A. R., 150. Brindley, T. A., 408. Brittin, G., 58. Bromley, S. W., 118, 606, 607. Bronson, T. E., 353. Brooks, J. W., 135, 301. Brown, A. W. A., 328, 530. Brown, W. J., 152. Brunson, A. M., 250, 367. Brunson, M. H., 129. Bryant, S. A., 178. Bryson, H. R., 592. Buchanan, L. L., 370, 530. Buchanan, W. D., 119, 365. Buckell, E. R., 241. Buckner, R. P., 410. Bugher, J. C., **349.** Buhl, C., 231. Bulger, J. W., 47, 123. Burdick, H. L., 569. Burgess, A. F., 212, 412. Burgess, E. D., 126, 364, 595. Burkholder, C. L., 351. Burks, B. D., 189, 401. Bushnell, R. J., 215, 220. Butac, F. L., 648. Butcher, F. G., 254. Butt, F. H., 328.

Caldwell, N. E. H., **436, 480.** Calinisan, M. R., 317. Callan, E. McC., **146, 468.** Callenbach, J. A., **361.**

Cameron, A. E., 381, 382, 383. Cameron, E., 489. Campbell, W. G., 178. Cañizo, J. Del, 458, 459. Cann, F. F., 146, 447. Cánovas, C., 458. Caprio, A. De, 164. Carlson, F. W., 559. Carlson, J. W., **567.** Carman, P. C., 397. Carpenter, J. R., 11. Carpenter, T. L., 246. Carruth, L. A., 481. Carter, R. H., 260. Carter, W. B., 113. Cartwright, W. B., 215. Caspari, E., 451. Cassil, C. C., 208, 555. Castro, A. Ruiz, 461. Celino, M. S., 316. Chaboussou, F., 26. Chamberlain, R., 63. Chamberlin, W. J., 625. Chan (Konyil), 153. Chandler, S. C., 196. Chapman, P. J., 1, 2, 127, 364. Chapman, R. N., 396. Charles, V. K., 500, 535. Charmoy, D. d'Emmerez de, 424. Charpentier, L. J., **601.** Chatterjee, N. C., **289**, 382. Chen, S. H., 524, 649 (note). Cherian, M. C., 20, 268, 270, 343, 518, 524. Chernova, F. I., 585. Chernuisheva, M. G., 579. Chiao, C. Y., 153. Chilton, St. J. P., 165. Chisholm, R. D., 419. Christensen, C. M., 165. Christensen, J. R., 496. Christenson, L. D., 403. Christian, M. B., 447. Clark, E. P., 397 Clausen, C. P., 431. Cleare, L. D., 538. Clemente, F. Gómez, 457, 458, 464. Coaton, W. G. H., 338. Cockerham, K. L., 601. Cole, J. R., 159. Collins, D. L., 502, 563, 644, 645. Colman, W., 216, 363. Compere, H., 290, 385.

Comstock, J. H., 275.

Cooke, O. B., 161. Coon, B. F., 134. Cooper, J. Omer-, 227, 614. Copenhafer, L. M., 190. Cory, E. N., 125, 202, 504. Costa, A. S., 141. Costas, L. A., 553. Cotterell, G. S., 106, 516. Cotton, R. T., 452, 605. Couch, J. N., 500. Coulter, R. W., 559. Cox, J. A., 195, 299. Craig, F. W., 562. Craig, R., 535. Crawford, J. C., 86. Crombie, A. C., 451. Crosthwait, S. L., 125. Cushman, R. A., 220, 293. Cuthbertson, A., 655. Cutright, C. R., 195.

da Fonseca, J. Pinto, 292, 344, 345. Daguerre, J. B., 497, 528. Dahms, R. G., 202, 258, 299. Daviault, L., 625, 626. Davis, A. C., 137. Dawsey, L. H., 112, 351, 352. Dean, A. G., 632. Dean, R. W., 563. De Caprio, A., 164. de Charmoy, D. d'Emmerez, 424. Decker, G. C., 326. Deen, O. T., 602. Deev, S. S., 571, 572. de Figueiredo jr., E. R., 292. de Fluiter, H. J., 493. de la Peña, F., 529. Del Cañizo, J., 458, 459. DeLong, D. M., 39, 248, 439. Demaree, J. B., 159. d'Emmerez de Charmoy, D., 424. Denier, P. C., 445. Derevyanikin, 581. De Santis, L., 445, 655. de Toledo, A. A., 624. Diakonoff, A., **524.** Dibble, C. B., **204.** Dick, R. D., 334. Dicker, G. H. L., 287. Dickson, R. C., 399, 553, 564. Dieter, C. E., 569. Dietrich, H., 645.

Dillon, G. F., 592.

(1349)

Ditman, L. P., 123, 152, 202. do Amaral, S. F., 511. Dobbins, T. N., **594.** Dodd, A. P., 587. Dohanian, S. M., 405. Dominick, C. B., 200. Doner, M. H., 592. Donohoe, H. C., 123. Dowden, P. B., 422, 424, 552. Downes, W., 417, 508. Dowson, W. J., 288. Drake, C. J., 326. Drews, E. A., 88. Driggers, B. F., 130. Duarte, A., 62. Dudley, H. C., 444. Dudley jr., J. E., 353. Dugas, A. L., 161, 601. Dumbleton, L. J., 150, 335, 541, 545. Dungan, G. H., 245. du Plessis, C., 612. Durocher-Yvon, F., 610. Dutky, S. R., 125, 369.

Eagleson, C., 260. Easter, S. S., 208, 413. Ebeling, W., 410. Eckert, J. E., 407. Eddy, C. O., **599**, **600**, **601**, **602**. Edel'man, N. M., **573**. Edwards, E. E., 526. Edwards, W. H., 293, 445. Ehrenhardt, H., 68. Eichmann, R. D., 376, 413, 419. Eidel'man, Z. M., 652. el Gammal, Mohammed Fouad, 338. Elliott, C., 244. Ellisor, L. O., 356, 601. Emden, F. I. van, **480.** Emerson, A. E., 396. Endo, K., 100. English, L. L., 597. Enser, K., 232. Enzie, J. V., 638. Erichson, W. F., 649 (note). Esaki, T., 101. Ess, M. W. Van, 407. Essig, E. O., 216, 416, 418. Evans, I. B. Pole, 17, 338. Evans, J. W., 152, 291, 467, 547. Evenden, J. C., 422, 647. Everly, R. T., 88.

Evlakhova, A. A., **583.** Ewer, D. W., **527.** Eyer, J. R., **414, 638.**

Fahey, J. E., 206, 412, 558. Fallscheer, H., 411. Farstad, C., 393. Fawcett, G. L., 608. Felt, E. P., 118, 212, 606. Fennah, R. G., 262. Fenton, F. A., 251, 258, 356, 635. Fenwick, E. M., 614. Ferguson, W. C., 247, 248. Ferreira, L., 63. Ferrière, C., 79, 80, 513. Fiedler, H., 449. Figueiredo jr., E. R. de, 292. Filatov, K. N., 570. Finlayson, L. R., 394. Fisher, E. H., 251. Fisher, R. A., 352, 408, 464. Fisher, R. C., 315, 593. 323, Fitzgerald, D. Vesey-, 82, 340. Fitzgerald, J. S., 480. Flanders, S. E., 219, 297, 355, 363, 385, 416. Fleming, W. E., **126**, **364**, **595**. Fletcher, R. K., **198**. Flint, W. P., **205**, **224**, **245**. Flock, R. A., 562. Floyd, E. H., 600, 602. Fluiter, H. J. de, 493. Fonseca, J. Pinto da, 292, 344, 345. Ford, O. W., 351. Fouad el Gammal, Mohammed, 338. Fox, C. C., 557, 559. Fox, H., 592. Fraenkel, G. S., 178. Frear, D. E. H., 371. Freitag, J. H., 500, 501. Friend, R. B., 161. Froggatt, J. L., 377, 379. Frost, S. W., 36. Fukushi, T., 272. Fullaway, D. T., 280. Fulton, B. B., 218, 259. Fulton, R. A., 240, 533. Furniss, R. L., 552. Furuichi, S., 96.

Gadd, C. H., 173. Gaenko, E. P., 575. Gahan, A. B., 569. Gahan, J. B., 556, 557. Gaines, J. C., 257. Gaines, R. C., 257, 358, 359, 600. Gallego, F. L., 8. Gallotti, M., 332. Galloway, A. G., 412. Gambrell, F. L., 120. Gammal, Mohammed Fouad el, 338. Gardner, J. C. M., 480. Garlick, W. G., 51. Garman, P., 87, 161, 162, 533. Garrison, G. L., 358. Garthwaite, P. F., 75. Gaufin, A. R., 400. Gay, F. J., 480. Geigy, R., 375. Geijskes, D. C., 169. Geissler, G. H., 363. Gemignani, E. V., 497. Genest, M. E., 36. Gentner, L. G., 359. Georgobiani, T. A., 653. Gerasimov, A. M., 478. Ghilarov (Gilyarov), M. S., 484. Gibson, A., 152. Gibson, A. L., 647. Giddings, N. J., **501.** Gilyarov (see Ghilarov). Gimingham, C. T., 40. Ginsburg, J. M., 42, 127, 194. Gippius, I. L., 580. Girth, H. B., 369, 370. Glaser, R. W., 369, 370. Glasgow, H., 131. Glass, E. H., 200. Glick, P. A., 50. Glover, P. M., 263, 264, 523. Gnadinger, C. B., 559. Gobeil, A. R., 311. Goddard, W. H., 288. Golding, F. D., 88, 515. Gómez Clemente, F., 457, 458, 464. Gómez-Menor Ortega, J., 498. Gonçalves, C. R., 167, 168. Gonzales, S. S., **649**. Gooden, E. L., 397. Goodey, T., 174. Goodhue, L. D., 127, 397, 555. Goot, P. van der, 182. Gorham, R. P., 302.

Gould, E., 363. Gourlay, E. S., 545. Graham, C., 41, 202. Graham. L. T., 406, 601. Graham, S. A., 396, 507. Granovsky, A. A., 501. Gray, E., 288. Grayson, J. M., 134, 554. Green, T., 394. Greenall, A. F., 89. Greenslade, R. M., 283. Greenwood, W., 592. Griffin, S. W., 360. Griffith, A. L., 382. Grigor'eva, T. G., 573. Griswold, G. H., 644. Groom, J. L., 238. Guérin-Méneville, F. E., 35. Guill jr., J. H., 123. Gunderson, H., 327. Gunn, D., 227. Gunn, D. L., 178, 224, 527. Guseva, A. V., 576. Guyton, F. E., 251. Gwynn, A. M., 175.

Haasis, F. A., 415.

Hadersold, O., 70. Haeussler, G. J., 294, 372. Haley, W. E., 601. Haller, H. L., 126, 241, 397, 414. Halliday, O. E., 280. Hamilton, A., 334. Hamilton, C. C., 132, 203, 438. Hamilton, D. W., 196. Hamlin, J. C., 628. Hammond, G. H., 89. Handford, R. H., 394. Hansberry, T. R., 206, 353, 556, 644. Hanson, H. S., 323. Hardman, N. F., 535. Hardy, C. H., 284. Hargreaves, H., 175, 228, 516. Harman, S. W., 128, 642. Harmston, F. C., 217, 563. Haroon Khan, M., 266. Harrar, J. G., 116, 500. Harrington, C. D., 42. Harris, R. V., 288. Harris, W. V., 177, 338, 590. Harrison, P. K., 362, 596. Hartzell, A., 136, 595. Hartzell, F. Z., 127.

Haruta, D., 102. Haseman, L., 166, 432. Hawkins, J. H., 560. Hawley, I. M., 118, 594. Hayward, K. J., 36, 263, 280, 498. Hazelhoff, E. H., 452. Headlee, T. J., 131. Hedden, O. K., 276. Heidenreich, E., 385. Heit, C. E., 421, 432. Helfer, J. R., 655. Hely, P. C., 336. Henry, H. K., 421, 432. Heriot, A. D., 444. Herrick, G. W., 275. Hess, A. D., 242. Hesse, A. J., 320. Hester, K. H. C., 527. Hetrick, L. A., 211. Hewitt, J. L., 223. Hey, G. L., 285, 286. Hibraoui, H., 157. Hill, S. O., 117. Hille Ris Lambers, D., 27. Hinman, F. G., 408. Hinton, H. E., 340. Hodson, A. C., 165. Hoffman, W. A., 361. Hoffmann, C. H., 40, 166, 364, 550. Hoffmann, W. E., 152. Hogan, T. W., 332. Holdaway, F. G., 34, 273, 650. Holloway, J. K., 294. Holloway, T. E., 354. Holmes, F. O., 316, 437. Hood, C. E., 438. Hood, J. D., 273. Hopkins, G. H. E., 175. Horsfall, W. R., 566. Hoskins, W. M., 407. Hosny, M., 30. Howard, N. F., 114, 240. Howes, F. N., 483. Howland, R. B., **592**. Hoyt, A. S., 358. Hrenoff, A. K., 647. Hrisafi, C. P., 12. Huber, G. A., 418. Huber, L. L., 253, 529. Huffaker, C. B., 560. Hurst, H., 239. Husain, M. Afzal, 266, 272, 341, 516. Hustache, A., 445, 468, 469.

Hutchins, R. E., 280. Hutson, J. C., 594. Hutson, R., 117, 558. Hyam, G. M., 290.

Ibatulina, F. S., 580.
Ijima, K., 172.
Ikeda, Y., 102.
Ileri, Mesude, 288.
Imerlishvili, M. I., 583.
Ingram, J. W., 601.
Iragorry, L. Briceño-, 292.
Isaac, P. V., 184.
Isely, D., 2, 3, 201.
Ishii, G., 101.
Ishii, T., 40, 172, 344 (note).
Ishikura, H., 172.
Ivanova, N. A., 585, 653.
Iyengar, S. Rangaswami, 382.
Iyer, C. S. Venkatsubba, 270.
Iyriboz, Nihat, 288.

Jack, R. W., 184. Jacob, F. H., 27, 590. Jacques, C., 391. Jamieson, C. A., 473. Jancke, O., 231, 232. Janjua, N. A., 267, 518. Jayaratnam, T. J., 619. Jayewickreme, S. H., 88. Jenkins, C. F. H., 169, 380, 540. Jepson, W. F., 609. Jervis, L. A., 293. Jewett, H. H., **363, 482.** Joachim, A. W. R., **592.** Johannsen, O. A., 328. Johnson, A. C., 47, 123. Johnson, C. G., 40. Johnson, D. L., 351. Johnson, G. V., 204, 214. Johnson, J. P., 161. Johnson, P. C., 357. Johnston, C. O., 366. Jones, D. P., **314**, **315**. Jones, E. T., **366**, **505**. Jones, H. A., 30, 126. Jones, S. C., 261. Julliard, R., 376.

Kagy, J. F., **349**, **411**, **560**, 647. Kakizaki, T., **99**. Kalra, A. N., **272**. Kalshoven, L. G. E., 270. Kamenkova, 581. Kamiya, K., 172. Kandiah, S., 592. Kanzawa, T., 622. Kasikhin, A. N., 573. Kaston, B. J., 40. Katsumata, K., 97. Kawada, A., 99, 173. Kawasaki, M., 100. Kayashima, 95, 103. Kayudin, 570. Kearns, H. G. H., 108, 152. Keifer, H. H., 259, 565. Kerr jr., T. W., **258, 643.** Khabirova, M., **454.** Khalilova, R., 455. Khan, M. Haroon, 266. Khoroshkevich, N. S., 654. Kimberley, A. E., 536. King, C. B. R., 174, 381. King, K. M., 396. Kinoshita, S., 172. Kirichenko, A. N., 480. Kishida, K., 172. Kisliuk jr., M., 133. Kiyashko, P. I., **584.** Kiyoku, M., 102. Klein; H. Z., 339. Klots, A. B., 628. Knight, K. L., 209. Knight, P., 189. Knowlton, G. F., 136, 217, 563. Knutson, H., 477. Koidzumi, K., 104. Kôno, H., 40, 96, 171, 173, 480. Kono, T., 97. Konyil Chan, 153. Koyama, T., 99, 173. Kozhanchikov, I. V., 478. Kozlova, E. N., 584. Kraiter (Kraitor), A. D., 651. Krishna Ayyar, P. N., 268, 270, 515, 521. Krishnaswami, C. S., 542. Kruim, Ya. S., **579.** Kubota, K., 104. Kurosawa, T., 101. Kuwana, Z., 101. Kuwayama, S., 93. Kylasam, M. S., 20.

Laan, P. A. van der, 318, 319. LaDue, J. P., 349.

Lal, K. B., 516. Lambers, D. Hille Ris, 27. Lambert, R., **36**, **431**, **626**. Lamerson, P. G., **632**, **633**. Landis, B. J., 114. Lane, F., 292. Langford, G. S., 125. Langronier, C., 616. Lapina, V. F., 582. Larchenko, K. I., 579. Large, E. C., 111. Large, J. R., 159. La Rivero, I., 562. Lateef, A., 269. Laubscher, F. X., 321. Lauderdale, J. L. E., **628.** Lawrence, R. F., **322.** Lavoipierre, M. M. J., 320. Lea, A., 613. Lea, A. M., 547. Leach, J. G., 116, 165, 239. Leake, C. D., 647. Leefmans, S., 380. Leffert, I., 325. Leroy, J. V., 516. Lesnikovskaya, A. Ya., 585. Leukel, R. W., 165, 477. Lever, R. J. A. W., 147, 169, 279, 389, 390, 495, 496. Levin, C., 201. Lewin, C. J., 491. Lezhava, V. V., 528. Liebermann, J., 511. Lilly, J. H., 205. Lincoln, C. G., 198, 252, 357, 643. Lindford, M. B., 273, 274. Lindgren, D. L., 398, 553, 564. Lindsay, E., 55. Linn, M. B., 646. Linsley, E. G., 213. List, G. M., 21, 278. Liu, C. L., 446. Livingstone, E. M., 208, 302, 555. Livshitz, 582. Lizer y Trelles, C. A., 343, 538. Llanos, V. V., 9, 53. Lloyd, D. C., 64. Lord, F. T., 307. Loughnane, J. B., 329. Lounsbury, C. P., 320. Lowman, M. S., 398. Lozina-Lozinskii, L. K., 235. Lubatti, O. F., 84.

Ludwig, D., 224, 592.

Lühmann, M., 68.

Lundie, A. E., 611. Lutherman, C. Z., 592. Lyle, C., 356, 647.

McCall, G. L., 349, 411, 560. McCarthy, T., 54, 143. McCarty, F. G., 295. McCoy, E. E., 369, 370. McCrory, S. A., 45, 160. McCulloch, E. C., 44. McDaniel, E. I., 117, 277, 563, 605. McDonald, J., **544.** McGovran, E. R., **208, 555.** McGregor, E. A., **530.** Mach, G. E., 572, 573. McIndoo, N. E., 573. McKelvey jr., J. J., 500. McKenna, G. F., 136, 595. McKeown, K. C., 467. Mackie, D. B., **113, 221, 565, 629.** MacLagan, D. S., **63.** McLane, S. R., 432. McLean, D. M., 502. MacLeod, G. F., 543, 544. McLeod, J. H., 309. McNally, A. G., 304. Maercks, H., 68. Magee, C. J. P., 57, 149. Magill, W. W., 2. Mahadevan, V., 343. Maheux, G., 626. Maines, W. W., 126, 595. Mains, E. B., 152. Maki, M., 88. Maksimovich, A. E., 586. Malenotti, E., 155. Mamet, R., 424, 514, 610. Mani, M. S., 514, 522. Manis, H. C., 325. Manninger, A. G., 12. Marcovitch, S., **534.** Margabandhu, V., **480, 515.** Markwood, L. N., **351, 352.** Márquez, V. Moreno, 459, 460, 461. Marshall, Sir G. A. K., 79, 265, 341. Marshall, James, 411. Martin, C. H., 415. Martin, H., 108. Martin, J. H., 299. Martin, J. T., 29, 483, 527. Martland, J. G., 116. Martorell, L. F., 51, 217, 473. Mason, P. W., 45.

Massee, A. M., 282, 283, 284. Mathers, W. G., 328, 415. Mathes, R., 354. Mathew, G. E. A., 434. Mathis, W., 474. Mathur, C. B., 341. Maulik, S., 468. Maxwell, C. W. B., 307. Mayer, E. L., 208. Medler, J. T., 414. Medvedeva, V. I., 583. Mendes, L. O. T., 36. Mendizábal Villalba, M., 463. Méneville, F. E. Guérin-, 35. Menon, K., 270. Menor Ortega, J. Gómez-, 498. Menozzi, C., 316. Merino, G., 649. Metalnikov, S., 583. Metzger, F. W., 118. Meuche, A., 235. Meyer, E., 231. Miao, C. P., 544. Michelbacher, A. E., 401, 402, 409. Middlekauff, W. W., 206. Mikul'skiĭ, A. A., 586. Milanez, F. R., 168. Milisavl'ević, D., 72. Miller, D., 333. Miller, E. V., 592. Miller, H. D. O., 505. Miller, J. W., 444. Miller, N. C. E., 90, 512, 548. Mills, H. B., 536. Mitchell, H. L., 400. Mitchener, A. V., 648. Mitono, T., 103, 104. Miwa, Y., **93, 94, 95.** Moericke, V., **234.** Mohammed Fouad el Gammal, 338. Montgomery, H. B. S., 281. Moore, E. S., 491. Moore, J. B., **557**, **559**. Moore, M. H., **281**. Morales Agacino, E., 462. Moran, E. J., 356. Moreno Márquez, V., 459, 460, 461. Morgan, W. L., 335. Moriyama, T., 88, 94, 95. Morley, P. M., 312. Morofsky, W. F., 224.

Morozov, S. F., 240.

Morris, V. H., 207, 275.

Morrison, F. O., 280.

Morrison, H. E., 249.

Moses, C. S., 166, 364.

Mossop, M. C., 184.

Mote, D. C., 249.

Moutia, L. A., 83.

Moznette, G. F., 159.

Muesebeck, C. F. W., 86, 137.

Mukerji, S., 269.

Mundinger, F. G., 121.

Munro, H. K., 524.

Munro, J. W., 186, 285.

Murayama, J., 592.

Murillo, L. M., 8.

Murray, G. H., 377.

Murrill, W. A., 475.

Mustafa, A. M., 267.

Nakayama, S., 92, 103. Narayanaswami, P. S., 268. Narematsu, I., 99. Nash, K. E., 644. Nash, R. W., 637. Nath, R., 81. Naudé, T. J., 156. Neal, P. A., 444. Neary, M. E., 226. Neiswander, C. R., 207, 275, 276. Nel, R. G., 434. Nel, R. I., 435. Nelson, O. A., 477. Nesterenko, P. A., 15. Nettles, W. C., 253. Neu, W., 488. Newcomer, E. J., 362. Newell, W., 358. Newsom, L. D., 602. Nicholls, H. M., 151. Nicholson, A. J., 397, 490, 491. Nickels, C. B., 133, 159. Nicol, J. M., **616.** Nikol'skiĭ, V. L., **577.** Noble, W. B., **215.** Nolte, M. C. A., 612, 613. Norris, K. R., 587. Norton, L. B., 206, 245, 353, 556, 644. Nosovskaya, T. Ya., 651. Notley, F. B., **175, 589.** Novodranov, Yu. K., **584.**

Nye, W. P., 136.

Oboussier, H., 488. Ocfemia, G. O., 317. O'Connor, B. A., 91, 377, 379. Ohta, Y., 100. Oike, K., 98. Oishi, T., 97. Okado, I., **98.** Okamoto, D., **152, 171.** Okazaki, K., **98.** Olds, H. E., **415**. Oman, P. W., 608 (note). Omer-Cooper, J., 227, 614. Oosthuizen, M. J., 58, 321. Orchard, O. B., **144.** Orelli, O. Schneider-, 67. Orr, L. W., 507. Ortega, J. Gómez-Menor, 498. Osburn, M. R., 135. Osorio, J. M., 538. Ostrovskii, N. I., 576. Ostwald, W., **433**. Otanes, F. Q., 648. Ozaki, S., 97.

Packard, C. M., 300. Page, A. B. P., 84. Paillot, A., 224. Painter, R. H., 202, 366, 367. Palm, C. E., 199, 252, 357, 643. Park, T., 592. Parker, J. H., 366. Parker, K. G., 502, 563, 645. Parker, R. L., 190, 632, 633, 635. Parkin, E. A., 340. Parks, T. H., 254. Parman, D. C., **355.** Parr, W. J., **525.** Parris, G. K., 140 Parson, H. E., 159. Parsons, F. S., 80. Patel, J. S., 446. Paul, L. C., 393. Paulian, R., 547. Pearce, G. W., 412. Pechuman, L. L., 40, 627. Peirson, H. B., 637. Pen, D. S., 153. Peña, F. de la, 529. Pepper, B. B., 122, 217. Pepper, J. H., 161. Pequeño, L., 464. Pérez, M. Quilis, 155, 156. Perrottet, G. S., 35. Persing, C. O., 295. Perzelan, J., **339.**

Pescott, R. T. M., 30, 143, 149, 216, 436, 546. Petch, T., **424, 587.** Peterson, A. G., 501. Petlyuk, P. T., 13. Petrov, A. D., 454. Petrov, P. M., **576.** Petrukhin, K. V., **653.** Petty, B. K., 278. Phillips, A. M., 47, 556, 557. Phillips, J. S., **324.** Phillips, W. J., **367.** Pickett, A. D., **226.** Pickles, A., 52, 373, 648. Pielou, D. P., **224.** Pierce, W. C., **133, 159.** Pilson, J. E., **550.** Pinckney, J. S., 47, 400. Pinto da Fonseca, J., 292, 344, 345. Planes, S., 464. Plank, H. K., 261, 346. Plessis, C. du, **612.** Plumb, G. H., **164.** Poisson, R., 23. Pole Evans, I. B., 17, 338. Polezhaev, V., 487. Pollacci, G., 332. Poos, F. W., 244, 354. Popov, K. I., 181. Popova, A. I., **652.** Porter, C. E., **512.** Potter, C., **29, 591.** Potts, S. F., **256.** Prebble, J. S., 347. Pruthi, H. S., 183, 265, 522. Putman, W. L., 304, 306, 548. Pyenson, L., 404, 624.

Quilis Pérez, M., 155, 156.

Radzievskaya, S. B., 486.
Rahman, K. A., 81, 175, 271, 272, 520, 523.
Ram(a)chandra Rao, Y., 269, 519.
Ramakrishna Ayyar, T. V., 319, 480.
Rangaswami Iyengar, S., 382.
Rankin, W. H., 502.
Rao, Y. Ram(a)chandra, 269, 519.
Ratcliffe, F. N., 480.
Rawlins, W. A., 643, 644, 645.
Read, W. H., 144.
Readio, P. A., 563.
Reed, W. D., 302, 362, 365.

Reeks, W. A., 474, 531. Rees, D. M., 400. Regnier, R., 24. Reid jr., W. J., 158. Reinhard, H. J., 213. Rex, E. G., 368. Reyne, A., 380. Richards, O. W., 313. Richardson, C. H., 406, 483, 647. Richardson, H. H., 132. Ripley, L. B., 278. Ris Lambers, D. Hille, 27. Ritcher, P. O., 2, 261, 298, 362, 554. Rivero, I. La, 562. Rivnay, E., 73. Roark, R. C., 135, 592. Robbie, W. A., 328. Roberts, F. M., 26, 179. Robertson, W. H., 22, 330. Rodríguez, R., 497. Roesler, R., 232. Romanevich, B. V., 584. Romney, V. E., **533.** Roney, J. N., **47**. Roonwal, M. L., **518.** Rosa, J. S., 50. Roth, L. M., 592. Rubtzov, I. A., 484. Ruhmann, M. H., 23, 330. Ruiz Castro, A., 461. Rumyantzev, M., 487. Rusk, H. W., 206, 412. Russell, G. A., 398.

Saakyan, A., 452.
Sachtleben, H., 71.
Safronovich, B. M., 572.
St. John, J. L., 44.
Saito, H., 98.
Sakai, K., 98, 102.
Sakimura, K., 139, 274.
Salman, K. A., 312.
Salt, G., 340.
Sameshima, T., 101.
Sampson, W. W., 88.
Samuel, C. K., 523.
Sanderson, M. W., 2, 3.
Santis, L. De, 445, 655.
Sapra, A. N., 269, 271, 521.
Sardiña, J. R., 456.
Sauer, H. F. G., 623.
Sawamoto, T., 480.
Sayers, R. R., 444.
Sayre, J. D., 275.

Sazonov, P. V., 584, 654. Schneider, F., 105. Schneider, H., 528. Schneider-Orelli, O., 67. Schoof, H. F., 566. Schread, J. C., **161.** Schwardt, H. H., 198, 643. Schweizer, J., 492. Scott, L. B., 345. Searls, E. M., 42, 398. Seiferle, E. J., 406. Seixas, C. A., 511. Sekiya, H., 173. Sellers, W. F., **424.** Sellke, K., **70.** Severin, H. H. P., 301. Shabanova, H. P., 579. Shands, W. A., **404.** Shaw, F. R., **418, 549.** Shaw, H., 281. Shepard, C. E., 162. Shepard, H. H., 186. Shotwell, R. L., 603. Shumakov, E. M., 571. Shvetzova, O. I., 583. Siegler, E. H., 129, 241. Sievers, A. F., 88, 398. Silva, P., 392. Simmonds, F. J., **66.** Simmonds, H. W., **148.** Simpson, G. W., 188. Sitnikova, G. M., 15, 617. Slabaugh, R. E., **401**. Small, T., 63. Smee, C., **59, 60, 62, 153, 528,** 588. Smirnova, E. K., **584.** Smit, B., **615.** Smith, A. J., 184. Smith, C. E., 503. Smith, C. F., 352. Smith, C. M., 30, 397. Smith, C. W., 308. Smith, F. F., 204, 214, 256, 361, 532. Smith, G. L., 359. Smith, G. R., **161.** Smith, H. S., 397, 548. Smith, J. H., 465. Smith, K. M., 153, 542. Smith, M. R., 346. Smith, R. C., 224. Smith, R. H., 114, 209, 415, 564. Smith, S. G., 474. Snapp, O. I., 159, 197, 418.

Snelling, R. O., 187. Snipes, B. T., 10. Sohi, G. S., 523. Solodovnikova, O., 453. Sonan, J., **93, 94.** Sotola, J., **44.** Soukhov (see Sukhov). Spaeth, F., **331.** Spain, L. A., 327. Spencer, G. J., **394.** Spencer, H., **475.** Speyer, E. R., 144, 525. Speyer, W., 69. Staehelin, M., 66. Stahel, G., 169. Stanley, J., 450. Stanley, W. W., 534. Stearns, L. A., 508. Steer, W., 281, 282. Steiner, H. M., 191. Steiner, L. F., 193. Steinhaus, E. A., 285. Steinweden, J. B., 565. Stephenson, P. R., 230. Stirrett, G. M., 303. Stitt, L. L., 441. Stracener, C. L., 602. Stringfield, G. H., 529. Strong, L. A., 425, 499. Stultz, H. T., 311. Subramaniam, C. K., 268. Suenaga, H., 99. Sugihara, Y., 171. Suire, J., 25. Sukhov (Soukhov), K. S., 13. Sukhova, M. N., 13. Sullivan, W. N., 127, 361, 398, 555. Sundaram, C. V., **524.** Suneson, C. A., **215.** Swain, A. F., 410. Swain, R. B., 365. Swan, D. C., 547. Swank, G. R., 208, 554, 555. Sweetman, H. L., 63, 395, 420. Swenk, M. H., 277, 278. Swezey, O. H., 49, 274, 275, 330. Swingle, M. C., 241, 556, 557. Swingle, W. T., 241. Sy, M., 71.

Tabata, S., 95.
Takano, S., 92.
Takahashi, H., 104.
Takahashi, R., 88, 94, 622.
Takahashi, Y., 152.

Tamanuki, K., 95, 96. Tate, H. D., 277, 278. Tattersfield, F., 29, 483. Taylor, T. H. C., 175, 229. Telenga, N. A., 455, 480, 582. Tessier, G. A., **392.** Theobald, F. V., **590.** Thomas, A. S., 175. Thomas, C. A., 299. Thomas, E. L., 186. Thomas, F. L., 499. Thomas, I., 27, 180. Thomas, J. G., **341.** Thomas, K. M., **542.** Thompson, R. W., 303. Thomssen, E. G., **592.** Thorne, G., 187. Thorold, C. A., 373. Timoshenko, M. G., 586. Todhunter, E. N., 44. Toledo, A. A. de, 624. Tomlinson jr., W. E., 133. Tompkins, C. M., 234, 535. Tothill, J. D., 175. Townes jr., H. K., 220, 275. Townsend, C. H. T., 469. Townsend, J. F., 161, 533. Trägårdh, I., 324. Transhe, 583. Travis, B. V., **476.** Trehan, K. N., 374. Trelles, C. A. Lizer y, 343, 538. Tuck, J. B., 224. Tucker, R. W. E., 3, 5, 7, 160. Turner, N., 162. Turnipseed, G.F., 597. Twinn, C. R., 328. Tyler, L. J., 563. Tyumeneva, V. A., 581.

Uchida, T., 33. Udine, E. J., 400. Ullah, G., 266, 523. Underhill, G. W., 299. Urquijo, P., 463. Utsu, T., 97. Uvarov, B. P., 322.

Vaduzhev, 583. Valleau, W. D., 2, 166. van der Goot, P., 182. van der Laan, P. A., 318, 319. van der Vecht, J., 32, 328. van der Veen, R., 493. van Emden, F. I., 480.

Van Ess, M. W., 407. van Zwaluwenburg, R. H., 50. Varley, G. C., **490.** Vasser, R. E., **575.** Vecht, J. van der, 32, 328. Veen, R. van der, 493. Vel'tishchev, P. A., 570. Venkatsubba Iyer, C. S., 270. Verano, A., 8. Vesey-Fitzgerald, D., 82, 323, 340. Vevai, E. J., 180. Villalba, M. Mendizábal, 463. Vinson, C. G., 45, 160. Vinson, J., 609. Vinzant, J. P., 365. Vitzthum, H., 488. von Winning, E., 70. Vovk, A. M., 13. Vukasović, P., 71, 72.

Wainwright, C. J., 328, 592. Wakeland, C., 134. Walkden, H. H., **506.** Walker, H. G., 46, 121, 152, 249, 503, 504. Wallace, C. R., 236, 494, 608. Wallace, P., 163. Waloff, Z. V., 322. Walter, E. V., **250.** Walton, C. L., 109, 111. Walton, W. R., 300. Warburton, C., 146. Ward, J. F., **498.** Ward, K. M., 238. Watanabe, C., 34, 40, 445, 480. Waterhouse, G. A., 436. Waters, H. A., 39. Waterston, J. M., **138, 376.** Watkins, T. C., **260.** Watson, J. R., 475. Watson, M. A., 26, 65. Webs jr., J. E., 192. Webster, R. L., 376, 411. Weigel, C. A., 214. Weiland, G. S., 123, 152. Weiss, F., **636.** Weiss, H. B., 45. Wellman, F. L., 57. West jr., A. S., **551.** Wheeler, W. M., 499. Whelan, D. B., 277. White, R. T., 124, 125. Whitehead, F. E., 356. Whiting, P. W., 474.

Whitnall, A. B. M., 614. Whitten, R. R., 256. Whittington, F. B., 125. Wigglesworth, V. B., 239, 483. Wilford, B. H., **348.** Wilkinson, D. S., 79. Wilkinson, H., 338. Wille, J. E., 11, 40, 139, 509, 539. Williams, C. B., **152, 224.** Wilson, C. C., **354.** Wilson, G. F., 374, 448. Wilson, H. F., 255, 569. Wilson, J. W., 476. Winburn, T. F., 506. Wing, M. W., 88. Winning, E. von, 70. Winter, G., 234. Wishart, G., 88, 310. Witman, E. D., **39.** Woke, P. A., **481.** Wolcott, G. N., 157, 217, 552. Wolfenbarger, D. O., 221. Wood, A. A., 303. Wood, J., 374. Wood, M., 192. Wormald, H., 282. Worthley, H. N., 191, 371. Wright, D. W., 109, 383. Wright, J. A., 54. Wylie, W. D., **199, 643.**

Yablokova, V. A., **579.**Yago, M., **96.**Yaĭtzevskaya, E., **577.**Yamazaki, T., **171, 173.**Yashiro, H., **99, 103.**Yetter jr., W. P., **130.**Yothers, M. A., **216, 360, 409, 559.**Young, M. T., **358, 359.**Yuasa, H., **173.**Yvon, F. Durocher-, **610.**

Zařtzeva, A. Ya., 583.
Zaritsky, S. N., 15.
Zazhurilo, V. K., 15, 617.
Zeck, E. H., 467.
Zeuner, F. E., 489.
Zimmerman, E. C., 649.
Zinkernagel, R., 275.
Zirngiebl, L., 339.
Zolotarev, E. Kh., 487.
Zorina, L. M., 479.
Zorsch, C. P., 643.
Zotzenko, L. N., 580.
Zwaluwenburg, R. H. van, 50.

GENERAL INDEX.

In the case of scientific names, the page reference is cited only under the heading of the generic name.

When a generic name is printed in brackets, it signifies that the name is not the one adopted.

A.

Abacá (see Musa textilis).

abbreviata, Aphis (see A. rhamni).

abbreviatus, Diaprepes.

Abies, Prociphilus oriens on, in Japan, 173; pests of, in Quebec, 36,626; Hemerocampa pseudotsugata on, in U.S.A., 49.

Abies balsamea (Balsam Fir), Harmologa fumiferana on, in Minne-

sota, 507.

abietis, Chermes (Adelges); Hylobius; Neodiprion.

abnormis, Leptomastidea (Tanaomastix).

abrupta, Empoasca.

abruptorius, Exenterus. Abutilon, Prionobrachium fuscum on, in Argentina, 512; cotton pests on, in Sudan, 386, 387.

abyssinialis, Hypena.

Acacia, pests of, in S. Africa, 18, 19, 278; Pseudococcus on, in Argentina, 344; pests of, in Australia, 142, 335; Xyleborus dispar on, in Br. Columbia, 328.

Acacia arabica, Phenacoccus hirsutus on roots of, in Egypt, 30. Acacia catechu (Lac Host-tree),

pests of, in India, 263.

Acacia mollissima, Euproctis terminalis reared on, in S. Africa, 18.

Acaena, Antholcus varinervis imported into New Zealand against, 334.

acajutlae, Eutermes (Nasutitermes). Acalla (see Peronea).

Acalypha wilkesiana, Cryphalus jatropha on, in Fiji, 170.

Acanthocoris dentatus, on cacao and Solanum in Nigeria, 516.

Acanthocoris sordidus, bionomics and control of, in Japan, 171.

Acanthopsyche, on tea in Nyasaland, 588.

Acanthopsyche junodi, aeroplane dusting against, on wattle in S. Africa, 18.

Acanthoscelides (see Bruchus).

Acanthotermes, measures against, attacking tung trees in Nyasaland, 588.

Acarapis vagans, sp. n., distribution of, infesting honey bees, 528.

acasta, Melittobia.

Acaulona peruviana, parasite of Dysdercus ruficollis in Peru, 510.

accepta, Nacoleia (Omiodes).

Acemya dentata (parasite of Melanoplus), parasitised by Brachymeria tegularis in Illinois, 401.

Acer (see Maple).

Acer negundo, grasshoppers damaging, in U.S.A., 603.

Acer pseudoplatanus, Drepanosiphum platanoides associated with, in Eire, **329**.

Aceratagallia sanguinolenta, races of, in relation to virus disease in potato and clover, 500.

aceris, Trioxys.

Acerophagoides triangularis, gen. et sp. n., parasite of Coccids in Argentina, **344**.

Acerophagus nubilipennis, parasite of Pseudococcus adonidum in Porto Rico, **356**.

Acetic Acid, in baits, 414, 612; soil treatment with, against Lachnosterna, 400.

Acetone, as a solvent, 244, 360, 398. Acetyl Methyl Carbinol, in baits for Cydia pomonella. 414.

Cydia pomonella, 414. Achaea, damaging fruits in Gold Coast, 106, 107, 493.

Achaea catocaloides, in Gold Coast, 106, 107.

Achaea ezea, in Gold Coast, 106, 107.

Achaea faber, in Gold Coast, 106, 107.

Achaea janata, food-plants of, in Burma, 78, 79.

Achaea lienardi, defoliating trees in S. Africa, 19; in Gold Coast, 106, 107.

Achaetoneura archippivora, parasite of Lepidoptera in Hawaii, 34, 35, 650.

achatina, Dasychira (Olene).

Achillea millefolium, Galeruca tanaceti on, in Germany, 68.

Achrysocharella ritchiei (parasite of Leucoptera), bionomics of, in Tanganyika, 176.

Achrysopophagus, possibly hyperparasitic on Pseudococcus adonidum in California, 355.

aciculatus, Protanagyrus.

acinaciformis, Coptotermes. Aclypea opaca (see Silpha).

Aconitum, Plusia moneta on, in Britain, 448.

acontistis, Leucania.

Acrobasis caryae (Pecan Nut Casebearer), in U.S.A., 117, 133, 159, 211, 426; developing on foliage of pecan, 117; thrips predacious on, 211; measures against, 133, 426.

Acrocercops, species of, on chestnut and other trees in Japan, 99.

Acrocinus longimanus, bionomics and control of, in Brazil, 392.

Acroclisoides africanus, sp. n., parasite of Antestia in Belgian Congo and Kenya, 80.

Acrolyta mesochori (see Hemiteles). Acrosternum hilare, on Passiflora edulis in New South Wales, 144.

Actinomyces scabies, relation of Arthropods to, in potato in U.S.A., 501, 644.

aculeata, Sarcophaga. acuminatus, Neoclytus. acuta, Cantheconidea.

Adalia bipunctata, predacious on Aphids in Chile, 512; hibernating on apple in Germany, 70; effect of insecticides on, 70.

adamsoni, Porotermes.

Adansonia digitata, relation of Dysdercus fasciatus to, in Sudan, 387. Adapsilia, parasite of Schizonycha in Zanzibar, 83.

Adapsilia latipennis, in S. Africa, 609; imported into Mauritius against Lachnosterna smithi, 609.

Adelges (see Chermes).

Adelphocoris rapidus, tests of insecticides against, on cotton in U.S.A., 257, 258, 359.

Adesmus borgmeieri, on kapok in Brazil, 292.

Adhesives, banding with, 74, 269, 464, 623.

Adlay (see Coix lachryma-jobi).

Adonia variegata, value and suggested utilisation of, against Aphids on cotton in Russian Union, 486, 582; hibernation of, 486.

adonidum, Pseudococcus.

lablab Adoretus, on Dolichos Sudan, 388.

Adoretus sinicus, measures against, on beans in Hawaii, 650.

adspersus, Exenterus.

advena, Ahasverus (Cathartus).

Aëdes aegypti, effects of constant and varying temperatures 131.

Aegeria exitiosa (Peach Borer), bionomics of, in Ontario, 225 measures against, in U.S.A., 159.

aegopodii, Cavariella. aegypti, Aëdes.

aegyptiaca, Icerya.

aegyptium, Xanthogramma (Ischiodon).

Aelia, bionomics and importance of, on cereals in Russia, 574.

aenea, Halticoptera.

aeneoviridis, Catolaccus.

aeneus, Chalcodermus; Corymbites (Selatosomus); Meligethes.

Aeolarchis sphenotoma, on Pandanus in Fiji, 496.

Aeolus mellillus (dorsalis) (see Drasterius).

aequalis, Euaresta; chus; Pimpla. Heterobostry-

aerivora, Diplogaster.

Aerobacillus, use of, in baits for Cydia pomonella, 414.

Aerobacter, use of, in baits for Cydia pomonella, 414.

Aeroplanes (see Aircraft).

Aeropus sibiricus, in Asiatic Russia, **484 486**, **570**; climate affecting outbreaks of, **484 486**; natural enemies of, 485, 570.

Aerosol OT, as a wetting agent for lead arsenate, 595.

Aerua javanica, relation of Schistocerca gregaria to, in India, 519.

aeruginosa, Potosia.

Aesculus parviflora, reported toxicity of, to Popillia japonica, 118. aestivus, Rhizotrogus.

Aethina tumida, bionomics and control of, in bee-hives in S. Africa, 611; in Nigeria, 611.

affinis, Dibrachys; Exenterus: Pempherulus (Pempheres).

Afghanistan, Schistocerca gregaria in, 520.

Africa, French West, new parasite of Geometrid in, 514.

Africa, Portuguese East, Nomadacris septemfasciata in, 156.

Africa, South, cowpea Bruchids in, 19, 58, 321; forest pests in, 18, 19, 30, 278, 341; fruit pests in, **18, 227, 320, 341, 415, 435**; pests of turf in, 614; miscellaneous pests in, 19, 320, 615; foodplants of Heliothis armigera in, 80, 81; locusts in, 19, 20, 143, 144, 156, 322, 435, 612, 613, 614 termites in, 19, 338, 615; thrips and virus disease of plants in, 140, 491, 608; insect enemies of bees in, 611; natural enemies and biological control of insects in, 18, 19, 30, 320, 322, 385, 514, 611; utilisation of insects against Opuntia spp. in, 17, 18; Vanessa cardui on Xanthium spinosum in, 19; development of economic entomology in, 320; beneficial introduced into insects countries from 385, 609.

africana, Gryllotalpa. africanus, Acroclisoides.

afrum, Atoposoma variegatum. Agallia sticticollis (see Agalliana).

Agalliana ensigera, virus disease of beet transmitted by, in Argentina, 608; recorded as A. sticticollis, 608 (note).

Agalliana sticticollis, A. ensigera recorded as in Argentina, 608 (note). Agar, in mixtures for feeding

insects, 148, 395, 407.

Agathiella meridionalis, parasite of Tortrix postvittana in Tasmania, 151.

Ageneotettix deorum, aggravating soil erosion in Michigan, 205.

Ageniaspis pyrillae, parasite of Pyrilla in India, 183, 184.

Ageratum, Nezara on, in Fiji, 279. Ageratum conyzoides, Myzus persicae on, in India, 523; reservoir of tobacco virus diseases in Sumatra, 318.

agilis, Eulachnus; Microdus (Bassus); Pristomeridia (Pristomerus). Agonoderus lecontei (pallipes, auct.),

morphology of, 592.

Agral, as a wetter for sprays, 145; as an emulsifier for oil, 540.

agrili, Monogonogastra.

Agrilus anxius, bionomics of, in Canada, 347; in U.S.A., 347.

Agrilus hyperici, attempted introduction of, into Australia against Hypericum, 143.

Agriotes lineatus, measures against, damaging hops in Britain, 282; ecology of, in Russia, 573.

Agriotes mancus, bionomics of, damaging potato in New York, 644, 645.

Agriotes obscurus, in Russian Union, 572, 573, 577; bionomics of, 572, 573.

Agromyza, on beans in Nyasaland, 154.

Agromyza braziliensis, sp. n., on potato in Brazil, **36**; parasite of, **36**.

Agromyza coerulea, bionomics of, on sweet potato in Porto Rico, 261. Agromyza huidobrensis, on broad beans in Chile, 512.

Agromyza lantanae, on Lantana in India, **74**, **289**; parasites of, **289**. Agromyza obtusa, parasite of, on

Cajanus in India, 267.

Agromyza oryzella, bionomics of, in Japan, 98; sprays against, 99. Agromyza simplex, bionomics of,

on asparagus in Washington, 642. Agropyrum elongatum, resistant to Cephus cinctus in Canada, 393; wheat crossed with, 393.

Agropyrum repens, cutworms on, in Germany, 449; Coccid infesting, in Queensland, 466; reservoir of mosaic of cereals in Siberia, 14. Agrostis, Crambus spp. damaging, in California, 408.

Agrothereutes slossonae (see Spilocryptus).

Agrotis exclamationis, bionomics of, in Germany, 449, 450.

Agrotis orthogonia, protection of cereals against, in U.S.A., 428.

Agrotis segetum, bionomics of, in Germany, 449, 450; in Russia, 576, 583, 584; measures against, 576; experiments with fungus and bacterium against, 583, 584. Agrotis tritici (see Euxoa).

Agrotis vestigialis, bionomics of, in

Germany, 449, 450.

Agrotis vetusta, on asparagus in

Washington, 642.

Agrotis ypsilon, on tobacco in Fiji, 390; on flax in Formosa, 93; on beet in Sakhalin, 100; baits for, 100, 390; rearing of, for insecticide tests, 199.

Ahasverus advena, distribution of, in stored cacao, 616.

Aircraft, for broadcasting grass-hopper baits, 223, 428; methods of applying dusts from, against locust swarms, 156, 538; uses of, for applying insecticides against other pests, 152, 256, 300, 412, 438, 504, 655; surveys of damage by Toxoptera from, 356; for studying aerial distribution of insects, 262; danger of spread of pests by, 331, 544.

Alabama, Anticarsia gemmatalis in, 251; miscellaneous pests in, 48, 404, 494; pests of satsuma

orange in, 597.

Alabama argillacea (on cotton), in Argentina, 169; in Brazil, 10, 404, 469; in Colombia, 53; in Peru, 404, 510; in U.S.A., 3, 404; parasites of, 53, 169, 469, 510; migrations of, 404, 543; arsenicals against, 3, 10, 53.

alabamae, Winthemia (Hemimasi-

poda).

alaskensis, Pikonema.

alba, Lepidosaphes.

Alberta, Cephus cinctus in, 112; grasshoppers and their natural enemies in, 112, 308, 309, 428; Paratrioza cockerelli in, 330.

albicapitana, Rhyacionia (Petrova).

albicoxa, Elasmus.

albifrons, Decticus; Spilochalcis. albimaculata, Moonia.

albipes, Metallus; Technomyrmex (see T. detorquens).

Albizzia, Icerya seychellarum on, in Seychelles, 82; lac cultivated on, in Siam, 94.

albohirtum, Dermolepida (Lepido-

derma).

albolineatus, Dendrolimus (see D. superans); Microbracon (Bracon). albomaculatus, Elasmus.

albomarginatus, Chorthippus.

albotibialis, Eurytoma.

Alcohol, insecticides injected into apple trees in, against Saperda, 243, 244; and methyl bromide, soil treatment with, against Coleoptera, 203, 208, 568; use of, in spray containing rotenone, 360; factors affecting permeability of insect cuticle by, 240, 241; (sulphated), as a wetting and spreading agent for sprays, 134, 136, 277.

Alesia discolor (see Verania).

Aleuas lineatus, damaging pastures in Argentina, 512.

Aleurites spp., pests of, in Burma, 78, 79; pests of, in Nyasaland, 60, 62, 588.

Aleurocanthus woglumi, parasite and control of, on Citrus in Bahamas, 293.

Aleurodes brassicae (see A. proletella).

Aleurodes carpini (see Asterobemisia).

Aleurodes fragariae, doubtful status of, 374.

Aleurodes lonicerae, on strawberry in Britain, 374.

Aleurodes proletella, on cabbage and strawberry in Britain, 374; destroyed by cold, 374; parasites reared on, 374; synonymy of, 374.

Aleurodes rubicola (see Asterobemisia carpini).

Aleurodids (see Whiteflies).

Aleuroplatus citri, sp. n., on lemon in New South Wales, 88.

Alfalfa (see Lucerne).

Alfalfa Caterpillar (see Colias eury-theme).

Alfalfa Snout Beetle (see Otiorrhynchus ligustici).

Alfalfa Weevil (see Hypera variabilis).

Algeria, natural enemies of pests in, 83, 514.

Algerian Sahara, Ceratitis capitata in, 616.

alianta, Niphadolepis. alias, Autographa.

Alkylphenylbenzenesulphonic Acid, spreader for sprays containing, 361.

Allodontermes morogorensis, measures against, attacking tung trees in Nyasaland, 588.

Allodontermes schultzei, damaging grape-vines in S. Africa, 19.

Allotropa (parasite of Pseudococcus comstocki), in Japan, 425; introduction of, into U.S.A., 425.

allyni, Eupelmus.

Almond, mites on, in California, 211, 536; pests of, in India, 271, 523; Scolytus rugulosus on, in Peru, 539.

Alopecurus pratensis, Contarinia merceri on, in Britain, 314, 315. Alorco, a form of cryolite (q.v.), 356, 534.

Alsophila pometaria, tests of lead arsenates against, on shade trees in U.S.A., 119.

alternata, Mycodiplosis.

INDEX. 673

Althaea rosea (see Hollyhock). altiscuta, Pachyneuron.

Aluminium Acetate, as an adhesive for lead arsenate, 162.

Aluminium Arsenate, experiments with, in sprays, 67, 624.

Aluminium Silicate, as a carrier for derris dusts, 162.

Aluminium Sulphate, action of, in sprays, 162, 411, 412.

alvearium, Platybolium.

amabilis, Eublemma.

Amarantus, Hypolixus truncatulus on, in Madras, 522; Hymenia recurvalis on, in Virginia, 46.

Amathes c-nigrum, rearing of, for insecticide tests, 199.

Amazon Fly (see Metagonistylum minense).

Ambari Hemp (see Hibiscus cannabinus).

Ambeodontus tristis, damaging timber in New Zealand, 333.

ambitalis, Psara.

Amblymerus, parasite of Mayetiola destructor in Russia, 618.

Amblypelta cocophaga, causing nutfall of coconuts in Solomon Is., 325: natural enemies of, 325.

Amblyteles armatorius, parasite of Dendrolimus in Sakhalin, 95.

Ambrosia artemisifolia, insects on, in Peru, 511. Amelanchier, Saperda candida on, in

New York, 242. America, Gonia of, 280.

America, North, Buprestis of, 655; Chalcidini of, 189; Myzus of, 45; Pimplini of, 220.

American Hellebore (see Veratrum viride).

Malacosoma;

americana, Larra; Periplaneta.

amethystinus, Hylotrupes.

amicus, Sympherobius.

Ammonia, in baits for Trypetids, 91, 462, 465. Ammonium Alginate, improving

calcium-arsenate sprays, 584. Ammonium Caseinate, uses of, in sprays, 113, 419, 420.

Ammonium Fluoride, in baits for Trypetids, 462.

Ammonium Phosphate, in bait for Dacus oleae, 462.

Ammonium Polysulphide, in spray against Scirtothrips citri, 296.

Ammonium Sulphate, increasing susceptibility of oats to Halotydeus destructor, 141.

Amniscus polyraphoides, on apple in Peru, 511.

Amorbia emigratella, food-plants of, in Hawaii, 651.

amorphophalli, Rhizoecus.

Amorphophallus campanulatus, Galerucida bicolor on, in India, 319.

Amorphophallus variabilis, new Coccid on, in Java, 88.

ampelophaga, Haltica; mima (Theresia).

Ampelopsis quinquefolia (see Parthenocissus).

Amphicordus improportionalis, not infesting barked logs in Burma, 78. Amphimallon (see Amphimallus).

Amphimallus majalis, damaging turf, etc., in France, 24, 83; bionomics and control of, 24.

Amphimallus semivillosus, parasite of, in Morocco, 83.

Amphimallus solstitialis, damaging turf in France, 24, 83; parasite of, in Russia, 572.

Amphorophora rubi, possible vector of raspberry mosaic in Britain, 287, 288.

Amphorophora sonchi (see Macrosiphum).

amphytus, Rhizotrogus.

Amyl Acetate, tests of, in locust baits, **612.**

Anabasine, toxicity of, to eggs of Lygaeus kalmii, 352; tests with, against Mayetiola destructor, 576.

Anabasine Sulphate, tests with, against Aphids, 360.

Anabrus simplex (in U.S.A.), estimation of damage to forage crops by, 365; unusual type of injury to

Pinus ponderosa by, 48; temperature affecting feeding and migrations of, 428, 562; natural enemies of, 187, 223, 537; measures against, 326, 537.

Anacardium occidentale, damaging fruits of, in Gold Coast, 107.

Anachaetopsis nitidula, parasite of Galerucella luteola in Rumania, 13.

Anagrus armatus var. nigriventris, establishment of, against Typhlocyba froggatti in Tasmania, 291.

Anagyrus (parasite of Pseudococcus comstocki), in Japan, 425; introduction of, into U.S.A., 425.

Anagyrus coccidivorus (parasite of Pseudococcus brevipes), rearing and distribution of, in Porto Rico, 472.

Anagyrus fusciventris (parasite of Pseudococcus adonidum), in Australia and Hawaii, 298; establishment of, in California, 298, possibly parasitised by Achrysopophagus, 355; synonymy of, 355.

Anagyrus nigricornis (see A. fusci-

ventris).

Anagyrus pseudococci, parasite of Pseudococcus comstocki in Palestine, 339.

Anaitis plagiata, attempted establishment of, against Hypericum in Australia, 143.

analis, Bruchus.

Anaphoidea nitens (parasite of Gonipterus scutellatus), utilisa-tion of, in S. Africa, 18, 30; spread of, into S. Rhodesia and Nyasaland, 30.

Anaphothrips fragariae, on strawberry in Queensland, 466.

Anarhopus sydneyensis, in Australia, 297: establishment of, against Pseudococcus adonidum in California, 297, 298.

Anarsia lineatella, in California, 113, 210; thrips predacious on, 210: fumigation of packed plums and peaches against, 113.

Anasa tristis (on cucurbits U.S.A.), bionomics of, 120, 442-444; measures against, 163, 201, 596, 603.

Anastatoidea brachartonae, hyperparasite of Artona catoxantha in

Java and Malaya, 79.

Anastatus, parasite of Promecotheca papuana in New Guinea Territory, 378; parasite of Amblypelta cocophaga in Solomon Is., 325.

Anastatus antestiae, parasite of Antestia in Kenya and Uganda, status, synonymy and varieties of, 80.

Anastatus antestiae var. hancocki, 80. Anastatus antestiae var. longiclava, 80.

Anastatus bifasciatus, parasite of Antestia in Uganda, 230; status of supposed varieties of, 80.

Anastatus brevipennis (see A. antestiae).

Anastatus colemani, parasite of Tessaratoma javanica in India, 263.

Anastrepha, Dirhinus giffardi introduced into Dominican Republic against, 471; quarantine against introduction of, into U.S.A. from Porto Rico, 444; Tetrastichus giffardianus not reproducing on, 344.

Anastrepha chiclayae, on mango and cherimoya in Peru, 539.

fraterculus, fruits Anastrebha by, in Peru, 539: attacked species recorded as, in Porto Rico, 444.

Anastrepha mombinpraeoptans, in Porto Rico, 444, 471; rearing of Dirhinus giffardi against, 471.

Anastrepha suspensa, in Porto Rico, 444, 471; rearing of Dirhinus giffardi against, 471.

anatipennella, Coleophora.

ancoralis, Coccinella

Ancylis comptana (Strawberry Leafroller), in U.S.A., 190, 192, 193, 549, 633; parasites of, 190, 192, 193, 549; measures against, 633. Ancylis fragariae (see A. comptana). ancylivorus, Macrocentrus.

Ancylorrhynchus trapezicollis, sp. n., on Cocos spp. in Brazil, 10, 468.

ancylus, Aspidiotus.

Andaman Islands, new Calandra in, 79.

Andrena, parasitised by Myopa in Europe, 474.

angelicus, Pseudaphycus.

Angitia (parasite of Cydia nigricana), in Britain, 311; introduction of, into Canada, 311; (parasite of Nephantis serinopa), bionomics of, in Ceylon, 620, 621.

Angitia cerophaga (parasite of Plutella maculipennis), established in S. Africa, 19; imported into Australia, 142; establishment of, in New Zealand, 64, 333; oviposition habits and relation to other parasites of, 64, 65.

Angitia fenestralis (parasite of Plutella maculipennis), imported into Australia from New Zealand, 142.

Angitia | gardneri, utilisation of, against Hyblaea puera in Burma, 77; in India, 77.

Angitia molestae (parasite of Cydia molesta), in Japan and Korea, 87, 372; importation, rearing and colonisation of, in U.S.A., 87, 161. 295, 425.

Angoumois Grain Moth (see Sito-

troga cerealella).

Anguillonema, parasite of Dendroctonus frontalis in Virginia, 211. angulata, Perisierola. angustus, Cynaeus.

INDEX. 675

Anisandrus pyri (see Xyleborus dispar).

Anisoplia austriaca, parasite of, in Russia, 572.

Anisota senatoria, tests of aircraft spraying against, in U.S.A., 256. Annona, insecticidal properties of

species of, 29.

Annona cherimolia, Trypetids on, in Peru, 539.

Annona muricata, 29; Dacus ferrugineus on, in Ceylon, 594; Phytorus lineolatus on, in Guam,

Annona palustris, 29.

Annona reticulata, 29.

Annona squamosa, 29; Bryocoropsis laticollis on, in Gold Coast, 492.

annubilata, Nacoleia. annulata, Myzocallis.

Anobiids, digestive enzymes of, 340. punctatum (striatum), Anobium damaging timbers in Germany and Sweden, 233, 324; bionomics of, 233, 234.

Anogeissus acuminata, Coleoptera in timber of, in Burma, 78.

Anomala, parasites of, in Zanzibar,

Anomala binotata, in Kentucky, 299. Anomala delagoa, in Zanzibar, 83. Anomala flavipennis, in Kentucky, 299.

Anomala horticola, bionomics and control of, damaging grassland

in Britain, 110.

Anomala innuba, in Kentucky, 299. Anomala lucicola, in forest nurseries in New York, 421, 422.

Anomala nigropicta, in Kentucky,

Anomala oblivia, in forest nurseries

in New York, 421.

Anomala orientalis, on pineapple and sugar-cane in Hawaii, 49; in U.S.A., 203, 370; Neoaplectana glaseri parasitic in, 370; methyl bromide against, on azaleas, 203.

Anomala plebeja, in Zanzibar, 83. Anomala vetula, infesting golf courses

in S. Africa, 614.

Anomalon relictum, parasite Datana integerrima in Missouri, 167.

Anomis, damaging Citrus fruits in Gold Coast, 106.

Anomis luridula, dusts against, on cotton in Peru, 510; parasites of, **510.**

Anomis sabulifera, bionomics and control of, on jute in Bengal, 447. Anomis texana (see A. luridula).

Anomocaulus fulvovestitus, on pine-

apple in Fiji, 389, 390.

Anoplocelaeno, attacking Brontispa longissima in New Guinea Territory, 379.

Anoplocnemis phasiana, on cacao

in Malaya, 513.

Anoplolepis custodiens, attacking Cactoblastis cactorum in S. Africa,

Anoplolepis longipes, attacking Amblypelta cocophaga in Solomon Is., **325.**

Anoplonyx laricis, on larch Quebec, 37.

Anoxia, parasite of, in Morocco, 83. Anoxia pilosa, parasite of, Russia, 572.

Antestia, parasite and incidence of, on coffee in Tanganyika, 177, 589; Nematospora associated with, 177; measures against, 230, 589, 590.

Antestia faceta, ecology and importance of, on coffee in Uganda, 229, 230: natural enemies of, 229, 230.

Antestia lineaticollis (on coffee), in Belgian Congo, 80; in Kenya, 80; in Uganda, 80, 230; ecology and importance of, 230; sites of, 80, 230.

antestiae, Anastatus; Corioxenos.

Anthia sexguttata, liberation of, against Lepidoptera on Dalbergia sissoo in India, 74.

Antholous varinervis, imported into New Zealand from Chile against Acaena, 334.

Anthonomobsis mixtus. jumping buds of Prunus due to, in Florida,

Anthonomus bisignifer, new parasite of, in Japan, 40.

Anthonomus cinctus (see A. piri).

Anthonomus grandis (Cotton Boll Weevil), in U.S.A., 3, 257, 258, 262, 358, 359, 599, 600; caught on aircraft at high altitudes, 262; insecticides against, 3, 257, 258, 358, 359, 599, 600.

Anthonomus piri, measures against, on pear in Germany and Switzer-

land, 617.

Anthonomus pomorum, bionomics and control of, on apple in Germany, 69, 70.

Anthonomus vestitus, parasite of, on cotton in Peru, 510.

(1349)

Anthores leuconotus, value of shade and mulch against, on coffee in Nyasaland, 154.

anthracinus, Coccophagus.

Anthrenus (Carpet Beetles), in New Zealand, 335; Typhlodromus tineivorus attacking, 376; survey of data on, 335.

Anthrenus scrophulariae, traps for, in U.S.A., 255.

Anthrenus vorax, parasite of, in U.S.A., 137.

Anticarsia gemmatalis, outbreaks of, on leguminous crops in U.S.A., 251, 599; measures against, 252; temperature affecting toxicity of insecticides to, 356.

Antigastra catalaunalis, parasite of, in India, 184; in Russian Union, 478; larva of, 478.

Antigua, failure to establish Metagonistylum minense in, 4.

Antimonelle, 296. antiopa, Nymphalis. antipodum, Goniozus. antiqua, Hylemyia.

Antonina, damaging grasses in Queensland, 466.

Antrocephalus, parasite of Homona coffearia in Ceylon, 174.

Antrocephalus melitarae, parasite of Cactoblastis cactorum in Australia, 587.

Antrocephalus stokesi, parasite of Cactoblastis cactorum in Australia, 587.

Ants, associated with Aphids, 332, 430, 475, 607; associated with Coccids, 82, 168, 332, 346, 475; associated with other insects, 346, 467, 475, 601; destroying other ants, 325, 476; destroying other insects, 17, 62, 74, 82, 93, 105, 143, 172, 292, 325, 378, 442, 499, 517, 522; measures against, 40, 142, 168, 169, 175, 332, 430, 588.

Ants, Argentine (see *Iridomyrmex humilis*).

Ants, Small Fire (see Wasmannia auropunctata).

Anuraphis bakéri, on clover in Montana, 537.

Anuraphis crataegi, bionomics of, on apple in Britain, 282.

Anuraphis padi, on strawberry in Britain, 28.

Anuraphis roseus, measures against, on apple in U.S.A., 1, 127, 128, 164, 191, 194, 195, 248; technique for testing ovicides against,

195; transmitting mosaic of narcissus, 415.

anxia, Lachnosterna (Phyllophaga). anxius, Agrilus.

Anychus ricini, sp. n., food-plants of, in Punjab, 271.

Aonidiella, measures against, on Citrus in Transcaucasia, 653.

Aonidiella aurantii, in Australia, 144, 336; in California, 222, 336, 398, 399, 410, 426, 535, 561, 564; in Fiji, 389; bionomics of, in Japan, 102; on plants other than Citrus, 102, 222, 564; foodplant affecting parasitism of, by Habrolepis rouxi, 220; HCN fumigation against, 102, 336, 398, 410, 564; strains of, resistant to HCN, 399, 410, 535; other insecticides against, 102, 426, 561; A. citrina compared with, 144.

Aonidiella citrina (on Citrus), habits of, in New South Wales, 144; measures against, in Transcaucasia, 654.

Aonidiella orientalis, measures against, on lac host-trees in India, 263, 264.

aonidum, Chrysomphalus (see C. ficus).

Apanteles, parasite of Hyblaea puera in Burma, 77; parasite of Thosea spp. in Ceylon, 174; parasite of Heliothis armigera in Fiji, 148; imported into Hawaii against H. armigera, 496; parasitising various Lepidoptera in India, 74, 524; parasite of Laphygma exempta in Nyasaland, 589; in Uganda, 514; parasites of, 514, 524.

Apanteles artonae, parasites of, 79. Apanteles bordagei, parasite of Leucoptera in Tanganyika, 176.

Apanteles carpatus, parasite of Tineola biselliella in Switzerland, 376.

Apanteles congregatus, bionomics of, in N. Carolina, 218.

Apanteles dictyoplocae, sp. n., parasite of Dictyoploca japonica in Japan, 480.

Apanteles diparopsidis, parasite of Diparopsis castanea in Nyasaland, 588.

Apanteles expulsus, parasite of Cirphis unipuncta in Fiji, 495.

Apanteles flaviconchae, parasite of Colias eurytheme in Louisiana, 600.

INDEX. 677

Apanteles flavipes, parasite of mothborers in India and Ceylon, 184, 610.

Apanteles glomeratus (parasite of Pieris spp.), in Britain, 314; in Hawaii, 34, 35, 650; failure to establish in New Zealand, 314, 333; killed by wilt disease infecting P. rapae, 650; parasites of, 314.

Apanteles guamensis, parasite of Susumia exigua in Guam, 331.

Apanteles kurosawai, sp. n., parasite of Margaronia pyloalis in Japan, 34.

liparidis, Apanteles hosts and natural enemies of, in Japan, 172. Apanteles machaeralis, parasite of

Hapalia machaeralis in Burma,

Apanteles militaris, parasite of Cirphis unipuncta in U.S.A., 300.

Apanteles minor, parasite of Margaronia pyloalis in Japan and Formosa, 34.

Apanteles molestae, parasite of Cydia molesta in Japan and Korea, 372.

Apanteles obliquae var. niger, new parasite of, in India, 514.

Apanteles ordinarius, parasite of Dendrolimus in Sakhalin, 95.

Apanteles plutellae (parasite of Plutella maculipennis), proposed introduction of, into New Zealand, 64; oviposition habits and relation to other parasites of, 64, 65.

Apanteles praepotens, identity and hosts of, in Britain and Germany,

rubecula (parasite of Apanteles Pieris rapae), in Britain, 314; probable value of, in New Zealand, 333; parasites of, 314.

Apanteles taragamae, parasite Nephantis serinopa in India, 620; parasite of Cydia molesta Japan and Korea, 372. in

Apanteles tasmanica, parasite of Tortrix postvittana in Tasmania

and New Zealand, 151.

Apanteles thurberiae (parasite of Sacadodes pyralis), breeding and utilisation of, in Colombia, 8, 53.

Apanteles tischeriae, parasite of Parornix prunivorella in Kansas,

apantelophagus, Hypopteromalus. Apate indistincta, on Aleurites in

Nyasaland, **62.**

Apechthis, considered distinct from Pimpla, 220.

Aphelenchoides, associated with Dendroctonus frontalis in ginia, 211.

Aphelinus chrysomphali (see Aphy-

Aphelinus flavus (parasite of Myzocallis annulata), introduction and natural occurrence of, in Australia, **142.**

Aphelinus mali (parasite of Eriosoma lanigerum), value of, in New South Wales, 493; killed by cold in New Zealand, 333; utilisation of, in Peru, 511; natural dispersion of, in U.S.A., 362; effect of sprays against E. lanigerum on, 493.

Aphidius cardui, parasite of cotton

Aphids in Russia, 582.

Aphidius laricis (see Coelonotus). Aphidius pini (see Coelonotus).

Aphidius platensis, parasite of Toxoptera graminum in Argen-

tina, 36.

Aphids, relation of, to plant virus 503, 523, 532, 542, 608, 638; ants associated with, 332, 430, 475, 607; natural enemies of, 3, 34, 36, 42, 54, 92, 93, 95, 164, 165, 171, 184, 210, 211, 250, 266, 303, 331, 333, 362, 431, 445, 471, 486, 489, 510, 512, 582, 650, 651; tests of insecticides on, 29, 127, 132, 195, 353, 556, 585, 591; climatic factors affecting migration of, in Britain, 180; of Delhi, 266; classification and new species of, 27, 45.

Aphis, transmitting virus disease of banana in New South Wales,

Aphis abbreviata (see A. rhamni). Aphis citricidus, measures against,

on Citrus in Fiji, 389.

Aphis gossypii, in Bermuda, 138; in China, 446; in Colombia, 54; in Fiji, 171, 496; in Hawaii, 35, 140, 141, 650, 651; in India, 266; in New South Wales, 57; in Peru, 510; in Philippines, 317, 648; in U.S.A., 2, 359, 503, 603; on cotton, 2, 54, 359, 446, 496, 510, 603, 648; on other plants, 171, 650, 651; relation of, to virus diseases of plants, 35, 57, 138, 140, 141, 317, 503; natural enemies of, 3, 54, 171,

650, 651; effects of temperature and sunlight on, 3,651; measures against, 54, 171, 359, 603, 648, 650: dusts increasing infestation of cotton by, 3, 359.

Aphis idaei, possible vector of raspberry mosaic in Britain, 287,

Aphis laburni, Coccinellids predacious on, on cotton in Uzbekis-

tan, 486.

Aphis maidis, natural enemies of, in Guam, 331; in Hawaii, 35, 651; in India, 266, 523; in Queensland, 466; in U.S.A., 165, 187, 250, 276, 529; on maize, 276, 331, 523; resistance of strains of maize to, 187, 250, 529; on papaya and tomato, 651; on sorghum, 165, 466, 523; on wheat and oats, 523; on barley, transmitting sugar-cane 523; mosaic, 35, 523; not transmiting virus disease of papaya, 35; plants selenium-treatment of against, 165, 276.

Aphis medicaginis,* food-plants of, in Hawaii, 35, 650, 651; transmitting virus disease of

papaya, 35.

Aphis middletoni, on papaya and

tomato in Hawaii, 651.

Aphis pawneepae, bionomics and control of, on Cercis canadensis in Kansas, 635.

Aphis persicae (see Myzus).

Aphis pomi (on apple), in Britain, 281; in Germany, 231; Russia, 585; in U.S.A., 164, 194, 195, 360; bionomics of, 164; tests of sprays against, 194, 231, 281, 360, 585; technique for testing insecticides against, 195.

Aphis rhamni (abbreviata) potato), in Canada, 303; in Eire, 329; transmitting virus disease of potato in Maine, 188; winter food-plants of, 188; nomencla-

ture of, 188.

Aphis rumicis,* not transmitting virus disease of tobacco and tomato in Argentina, 608: tests of insecticides on, 29, 127, 353, 483, 556, 591.

Aphis, Apple (see Aphis pomi).

Aphis, Apple Grain (see Rhopalosiphum prunifoliae).

Aphis, Black Pecan (see Melanocallis caryaefoliae).

Aphis Aphis, Buckthorn (see rhamni)

Aphis, Cabbage (see Brevicoryne brassicae).

Aphis, Corn Leaf (see Aphis maidis).

Aphis, Cotton (see Aphis gossypii). Aphis, Foxglove (see Macrosiphum solani).

Aphis, Green Cereal (see Toxoptera graminum).

Aphis, Green Peach (see Myzus persicae).

Aphis, Hop-damson (see Phorodon humuli).

Aphis, Pea (see Macrosiphum onobrychis).

Aphis, Pink and Green Potato (see Macrosiphum solanifolii).

Aphis, Rosy Apple (see Anuraphis roseus).

Aphis, Spruce Trunk (see Chermes pineoides).

Aphis, Strawberry (see Capitophorus fragariae).

Aphis, Sugar-cane (see Sipha flava). Aphis, Tobacco (see Myzus persicae).

Aphis, Turnip (see Rhopalosiphum pseudobrassicae).

Aphis, Walnut (see Chromaphis juglandicola).

Aphis, Woolly Apple (see Eriosoma lanigerum).

Aphodius, bionomics and identity of species of, damaging pastures in Australia, 335, 547; in lawns, 548; measures against, 548.

Aphodius howitti, 335, 547. Aphodius tasmaniae, 335, 547.

Aphomia gularis, in stored grain in Japan, **97.**

Aphthona euphorbiae, ecological factors affecting, on flax in Russian Union, 181, 182.

Aphycus nigrivarius, parasite of Eriococcus coriaceus in Australia.

Aphytis, parasite of Chrysomphalus ficus in Brazil, 625.

Aphytis chrysomphali, parasite of Aonidiella aurantii in Japan, 102. apicalis, Hyperaspis festiva; Nepho-

tettix (see N. bipunctatus). apicicornis, Brachymeria.

apicipennis, Exora.

Apion corchori, bionomics of, on jute in India, 446, 522.

Apion martinezi, on Zanthoxylum flavum in Porto Rico, 427.

^{*} In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonymy.

Apion ulicis, utilisation of, against Ulex europaeus in New Zealand and Tasmania, 334.

Apis dorsata, Microbracon hebetor bred on, in India, 264.

Apis indica, new Tenebrionid in hives of, in Madras, 343.

aplanatideclivis, Xyleborus.

Aplanobacter stewarti, host range and insect vectors of, in U.S.A., 244.

Aplastomorpha calandrae, ecology of, parasitising Pempherulus affinis in Madras, 522.

Aplomyiopsis galerucellae (see Syna-

plomyia).

Apogonia, on teak in Burma, 77.

Apogonia cribricollis, bionomics and control of, on cacao, etc., in Malaya, 513.

appendigaster, Eurytoma.

Apple, 603; Ceratitis capitata on, in S. Africa, 18; pests of, in Argentina, 445, 496, 497; pests of, in Australia, 55, 150-152, 216, 238, 279, 291, 335, 465, 493; pests of, in Austria, 232; pests of, in Britain, 108, 281, 282, 285, 286, 287; pests of, in Canada, 51, 224, 226, 330, 508, 548; Cydia pomonella on, in France Switzerland, 66; pests of, in Germany, 68, 69, 71, 231; pests of, in Br. Guiana, 22, 23; pests of, in India, 267, 341, 518; pests of, in Japan, 96, 622; C. pomonella on, in Jugoslavia, 72; Lachnosterna morosa on, in Korea, 171; pests of, in New Zealand, 150, 151, 333; pests of, in Peru, 511, 539; pests of, in Russian Union, 580, 581, 582, 585, 653; *C. pomonella* on, in Spain, 457; pests of, in U.S.A., 1, 2, 41, 45, 48, 71, 113, 118, 119, 127, 128, 129, 134, 160, 161, 164, 190, 191, 192, 193, 194, 195, 206, 210, 242–244, 248, 276, 300, 351, 360, 361, 362, 363, 364, 371, 406, 411, 415, 425, 536, 537, 549, 559, 562, 563, 632, 633-635, 639, 640, 641, 642, susceptibility of 643, 645; varieties of, to Psylla mali, 232: Eriosoma lanigerum associated with canker on, 112; insecticides and injury to, 55, 119, 127, 162, 190, 192, 238, 300, 351, 406, 559, 585, 633, 639, 640, 641, 643, 652, 653; effect of fruit growth of, on spray deposits, 206.

Apple (Fruit), fumigation of, against pests, 18, 113, 364; problem of spray residues on, 44, 45, 128, 330, 351, 411, 412, 633, 634, 639; in baits for Tyloderma fragariae, 362; juice of, in bait-sprays for Dacus, 337.

Apple, Flowering (see Pyrus specta-

Apple Aphis (see Aphis pomi).

Apple Aphis, Rosy (see Anuraphis roseus).

Apple Aphis, Woolly (see Eriosoma lanigerum).

Apple Blossom Weevil (see Anthonomus pomorum).

Apple Budmoth (see Spilonota ocellana).

Apple Grain Aphis (see Rhopalosi-phum prunifoliae).

Apple Leaf-roller (see Tortrix argy-rospila).

Apple Maggot (see Rhagoletis pomonella).

Apple Sawfly (see Hoplocampa testudinea).

Apple Sucker (see Psylla mali).

Apple-tree Borer, Round-headed (see Saperda candida).

approximatus, Calotermes (Kalotermes); Pissodes.

Apricot, Cydia pomonella on, in S. Africa, 415; Ceratitis capitata on, in Algerian Sahara, 616; Tortrix sarcostega on, in Baluchistan, 518; pests of, in China, 446; Bostrychid boring in, in New South Wales, 335; pests of, in U.S.A., 210, 405, 536.

Aprostocetus roseveari, probably hyperparasitic on Phytolyma lata in Gold Coast, 492.

Aptesis tsugae (see Microcryptus). Aquilegia, Papaipema purpurifascia on, in New York, 644.

Arab Literature, references to insect pests in, 157.

Arachis hypogaea (see Ground-nuts). Araecerus fasciculatus, on soy beans in Ceylon, 594; in stored products in Venezuela and Gold Coast, 292, 293, 616.

Araucaria cunninghami, Ceramby-cid on; in Queensland, 467.

Archichauliodes diversus, bionomics of, in New Zealand, 334.

archippivora, Archaetoneura (Frontina).

Archips (see Tortrix).

Archytas, parasite of Alabama argillacea in Colombia, 53.

arcuata, Coccinella (Harmonia). Arcyptera microptera (see Pararcyptera).

areator, Hemiteles.

Areca catechu, Brontispa longissima on, in New Guinea Territory, 379. areolatus, Neoplectrus.

Arge pectoralis, on birch in Quebec,

argentina, Tachardiella.

Argentina, pests of apple in, 445, 496, 497; Toxoptera graminum on cereals in, 36; pests of Citrus in, 263, 445, 512; miscellaneous pests in, 445, 498, 512, 608; insects and virus diseases of plants in, 141, 608; catalogue of Coccids of, 343; Hylotrupes bajulus in, 497; locusts and grasshoppers in, 308, 322, 497, locusts and 511, 512, 528, 538; catalogue of Thysanoptera of, 655; exchange of natural enemies of grasshoppers between Canada and, 308; other beneficial insects in, 36, 169, 344, 445.

Argentine Ant (see Iridomyrmez humilis).

argillacea, Alabama.

Argyresthia ephippella (pruniella, auct.), tests of tar distillates against, on cherry in Germany, 231.

Argyria sticticraspis (see Diatraea).
Argyroploce illepida, on Inocarpus
edulis in Fiji, 496; in India,
137; bionomics of, on leguminous plants in Java, 182;
parasites of, 137, 496.

Argyroploce leucotreta, on cotton in E. Africa, 227, 228; on Citrus, etc., in S. Africa, 18, 227; bionomics of, 227, 228; other species possibly confused with,

227.

Argyroploce urticana, food-plants of, in Britain, 283; derris against, on strawberry, 283.

Argyroploce variegana, on apple in New York, 1.

argyrospila, Tortrix (Cacoecia).

Argyrotoxa (see Tortrix).

Aristotelia, measures against, on soy beans in Ceylon, 594.

Arizona, beet pests in, 429, 441, 533, 631; pests of lucerne in, 403, 429, 441, 628, 629; miscellaneous pests in, 562, 596, 629; beneficial insects in, 190, 442; Trichogramma introduced into Louisiana from, 601.

Arkansas, miscellaneous pests in, 2, 3, 201; Epicauta pennsylvanica destroying grasshopper eggs in, 566.

Arma custos (see Auriga). armatorius, Amblyteles.

armatus, Anagrus.

armigera, Heliothis (Chloridea).

Armyworm, Fall (see Laphygma frugiperda).

Armyworm, Southern (see Laphyg-ma eridania).

Armyworms (see Cutworms). arontes, Pelochyta.

arrogans, Trigonophymus (Dichro-

plus).

Arsenic, in baits, 136, 300, 354, 355, 357, 612, 613; tests of sheep dips containing, against Calandra oryzae, 540, 541; in spray against thrips, 414; uses of, against termites and Atta, 142, 169, 338, 610; effect of, on midgut wall of Laphygma eridania, 481; question of danger to bees and pheasants of sprays and baits containing, 136, 205, 550; problem of residues of, on sprayed apples, 330, 351, 412, 633; Ephestia killed by residues of, in stored tobacco, 365.

Arthrocnodax, 315.

Artichoke, new Jassid on, in India, 265.

articulatus, Selenaspidus (Pseudaonidia).

Artificial Silk (see Rayon).

Artocarpus communis (incisa) (Breadfruit), Margaronia caesalis on, in Ceylon, 594; Phytorus lineolatus on, in Guam, 332; Xylothrips religiosus on, in Samoa, 170; Icerya seychellarum on, in Seychelles, 82.

Artocarpus integrifolia, pests of, in Brazil, 391, 392; Dacus umbrosus on, in Malaya, 90; Icerya seychellarum on, in Seychelles, 82.

Artona catoxantha, parasites of, on coconut in Java and Malaya, 79. artonae, Apanteles; Euplectromorpha.

arundinis, Hyalopterus.

Ascogaster quadridentata (carpocapsae), introduced into Canada against Cydia nigricana, 310, 311; introduced into Spain and Peru against C. pomonella, 457, 511; parasite of C. pomonella in U.S.A., 193; introduction of, into U.S.A. against C. molesta, 295; parasite of, 193.

Ascotis (see Boarmia).

Aserica castanea, destroying maize seedlings in Connecticut, 164.

Ash (Fraxinus), Prociphilus oriens on, in Japan, 173; Anobium punctatum damaging timber of, 233.

Ash, Green (see Fraxinus lanceolata).

Asia, Central, wheat insects in. 655: Coleoptera in stored products in, 453, 454, 455; beneficial insects and suggested biological control of pests in, 486, 582.

Asiatic Beetle (see Anomala orient-

alis).

asozanus, Gelis (Pezomachus). asparagi, Crioceris; Zacerata.

Asparagus, Zacerata asparagi on, in S. Africa, 19; Aphis medicaginis on, in Hawaii, 651; pests of, in U.S.A., 48, 430, 642.

Asparagus Beetle (see Crioceris

asparagi).

Aspen, American (see Populus tre-

muloides).

Aspergillus flavus, infesting Gilpinia polytoma and Microcryptus basizonus, 637.

Aspidiotiphagus citrinus, parasite of Coccids in Japan and Brazil,

102, 625.

Aspidiotiphagus lounsburyi, parasite of Coccids in Nyasaland and Brazil, 62, 625.

ancylus. Aspidiotus measures against, on cranberry in Washington, 642.

Aspidiotus camelliae, auct. (see A.

rapax).

Aspidiotus destructor, in Florida, 472; in Guam, 331; in Siam, 95; in W. Indies, 263, 471, 498; natural enemies and biological control of, on coconut, 263, 331, 471, 498; on other plants, 95, 331.

Aspidiotus hederae, parasite of, on Aleurites in Nyasaland, 62; not attacked by Comperiella bifasciata in Spain, 457; Parlatoria oleae associated with, in U.S.A., 629.

Aspidiotus lataniae, on fuchsias and Camellia in California, 223.

Aspidiotus orientalis (see Aonidiella). perniciosus, in Aspidiotus Columbia, 23, 330; carriage of, by birds and bats in India, 272; in Mauritius, 610; in Russian Union, 582, 585, 652, 653;

U.S.A., 1, 223, 409, 629, 640, 642, 647; on cranberry, 640, 642; on coconut, 610; on fruit trees, 1, 23, 223, 272, 330, 409, 585, 647, 652, 653; utilisation of Chilocorus against, 582, 610; other measures against, 585, 640, 642, 647, 652, 653; lime-sulphur sprays preventing reproduction in, 223, 409.

Aspidiotus rapax, Parlatoria oleae associated with, in U.S.A., 629. Aspongopus viduatus, on melon in

Sudan, 388.

Assam, new Calandra in, 79. assimilis, Ceuthorrhynchus;

Gryllulus (Gryllus); Muscina.

assulta, Heliothis.

Aster Leafhopper (see Macrosteles divisus).

Asterobemisia, gen. n., 374.

Asterobemisia (Aleurodes) carpini synonymy of, **374.**Asterolecanium bambusae, natural

enemies and biological control of, on bamboo in tropical America, 471, 472.

Asterolecanium miliaris, natural enemies and biological control of, on bamboo in tropical America,

471, 472.

Asterolecanium variolosum, utilisation of Habrolepis dalmanni against, on oak in Australia, 142.

Asters (Callistephus), thrips transmitting spotted wilt to, in France, 25; pests of, in U.S.A., 301, 302, 361, 536; studies on vector, transmission and host-range of yellows disease of, 301, 302, 438, 502, 646.

Astomaspis (see Hemiteles).

astur, Opharus.

Astycus lateralis, on teak Aleuvites in Burma, 77, 79. Asynonychus (see Pantomorus).

atalantae, Theronia.

ater, Hylastes; Isaniris; genes.

atlanis, Sarcophaga.

atlanticus, Tetranychus.

Atoposoma variegatum var. afrum (parasite of Leucoptera), bionomics of, in Tanganyika, 176.

atra, Phyllotreta.

atrellus, Prosapegus. atricornis, Phytomyza.

atropurpureus, Eupelmus.

Atta cephalotes, measures against, on coffee and Citrus in Dutch Guiana, 169.

Atta sexdens, in Brazil, 40; on cassava in Dutch Guiana, 169; measures against, 40, 169.

Attagenus byturoides, bionomics and control of, in stored products in

Central Asia, 453.

Attagenus piceus, in U.S.A., 255, 363, 644; experiments with baits for, 255; effect of paradichlorobenzene on feeding of, 363.

Attalea spp., pests of, in Brazil, 9,

468.

atymnius, Castnia.

Aulacaspis madiunensis (on sugarcane), in Formosa, 622; in Queensland, 389, 622; susceptibility of varieties of sugar-cane to, 389; on grasses, 622.

Aulacaspis pentagona, on peach in Peru, 511; in W. Indies, 471, 498; on papaya, 498; natural

enemies of, 471, 511.

Aulacorthum solani (see Macro-siphum).

Aulocara elliotti, in Montana, **536.** aurantii, Aonidiella; Pulvinaria; Scirtothrips; Toxoptera.

aurata, Cetonia. auratus, Dirhinus. aurea, Lonchaea. auricilia, Diatraea. auricularia, Forficula.

Auriga custos, doubtful value of, against Leptinotarsa decemlineata

in Germany, 70. auritus, Conoderus.

auropunctata, Wasmannia.

aurulentus, Epitragus.

Austracris guttulosa, on cotton in Queensland, 466.

australasiae, Periplaneta.

Australia, Ctenolepisma longicaudata in, 55-57; miscellaneous pests in, 141-143, 355; orchard pests in, 141, 150–152, 238, 279, 494; pests of stored grain in, 380, 436, 540; insects and virus diseases in, 149, 608; beneficial insects and biological control of insects and plants in, 17, 141, 142, 143, 587, 608; parasites of Cactoblastis cactorum in, 587; introduction of beneficial insects into other countries from, 18, 297, 545; Gonipterus scutellatus probably introduced into S. Africa from, 30.

Australia, South, Aphodius howitti in, 547; Coccids and their natural enemies on Citrus and Eucalyptus in, 280, 290, 545, 546.

Australia, 'Western, Austroicetes cruciata and its egg-parasites in, 169; new Empusa infesting Halotydeus destructor in, 587; miscellaneous pests in, 141, 142. australicum, Trichogramma.

australis, Eucelatoria; Gryllotalpa; Teleonemia; Tenuipalpus.

Austria, orchard pests in, 232, 304; new Acarapis on honey bee in, 528. austriaca, Anisoplia; Eurygaster.

Austroicetes cruciata (jungi), eggparasites of, in W. Australia, 169; measures against, 169.

Autogiro (see Aircraft).

Autographa alias, bionomics of, on conifers in Quebec, 626.

Autographa brassicae (see Plusia). Autographa chalcites (see Plusia). Autographa rectangula, bionomics

Autographa rectangula, bionomics of, on conifers in Quebec, 626.

Autographa selecta, bionomics of,

conifers in Quebec, 626.

autographae, Meteorus. Autoserica (see Aserica).

autumnalis, Systoechus.

avellanae, Eriophyes. avenae, Macrosiphum.

Averrhoa, moths damaging fruits of, in Gold Coast, 107.

Averrhoa carambola, Dacus spp. on in Malaya, **90, 91;** juice of, in baits for D. ferrugineus, **91.**

avidus, Protostrophus.

Avocado, Pseudococcus adonidum on, in California, 297, 298, 355; Dacus ferrugineus on, in Ceylon, 594; pests of, in Fiji, 170; pests of, in Guam, 332; Pseudococcus spp. on, in Porto Rico, 346, 472; Icerya seychellarum on, in Seychelles, 82.

Avrosia translucens, probably on begonia in California, 536.

Axionicus insignis, on Brachychiton populneus in New South Wales, 467.

Axonopus compressus, Blissus insularis damaging lawns of, in Trinidad, 373.

ayyari, Tetrastichus.

Azadirachta indica (see Melia azadirachta).

Azalea, pests of, in U.S.A., 86, 203; methods of fumigating, with methyl bromide, 203.

Azochis gripusalis, measures against, on fig in Brazil, **392.**

Azya, introduced into Porto Rico from Brazil against Asterole-canium spp., 471.

Azya luteipes, 471; predacious on Chrysomphalus ficus in Brazil,

Azya trinitatis (predacious on Coccids), in Trinidad, 263, 471; introduction of: into Porto Rico, 471; into Hawaii, 472; into St. Lucia, 263.

B.

Baboons, destroying Cactoblastis cactorum in S. Africa, 17.

bacchus, Rhynchites.

Bacillus lentimorbus and popilliae, spp. n., causing milky disease in Popillia japonica in U.S.A., 369: temperature affecting

development of, 369.

Bacteria, review of relations between insects and, 285; (infecting insects), 24, 37, 106, 369, 379, 504; utilisation of, 124, 125, 504, 583, 584; 583, 584; (infecting plants), relation of insects to, 15, 53, 112, 239, 244, 245, 288, 647; (from fermenting syrup), effect of, in baits for Cydia pomonella, 414.

Bacterial Wilt of Maize (see Aplano-

bacter stewarti).

Bacterium prodigiosum, experiments with, against Lepidoptera, 583,

Bacterium salicis, experiments on relation of Xiphydria prolongata to, in willow in Britain, 288.

badiipennella, Coleophora.

badius, Termes.

Baeoanusia minor, hyperparasite of Saissetia oleae in Africa, 386.

Bahamas, parasite and control of Aleurocanthus woglumi on Citrus in, 293.

bahianum, Himatidium. baileyi, Calepitrimerus.

Baits, ineffective against Antestia, 230; for ants, 142, 332, 430, 475; for Coleoptera, 126, 252, 255, 357, 362, 548, 571, 577, 586, 596, 642, 643, 645; for crickets, 173, 237; for Forficula auricularia, 304, 538; for Julus moreleti, 138; for locusts and grasshoppers, 20, 136, 144, 169, 205, 223, 241, 269, 354, 355, 393, 394, 428, 461, 529, 612, equipment for 613, 614, 648; mixing and distributing, 144, 169; distributed from aircraft, 223, 428; costs of other methods of locust control compared with, 459; for Lepidopterous larvae, 31, 100, 138, 170, 171, 252, 300, 390, 429, 534, 542, 576, 598, 615, 631, 635; for moths, 107, 193, 238, 255, 414, 435, 465, 583, 584; for termites, 19, 387; for Thermobia domestica, 406; for Trypetids, 91, 94, 99, 183, 461, 462, 465, 563, 594; factors vitiating tests of, against Ceratitis capitata, 464, for other Diptera, 158, 283; formulae for, 91, 99, 100, 126, 142, 183, 237, 252, 269, 300, 304, 354, 355, 362, 406, 428, 430, 435, 461, 462, 465, 538, 542, 577, 612, 613, 614; types of containers for, 430, 462, 563.

Bait-sprays, for Antestia, 589, 590; for beet weevil, 586: for Trypetids, 54, 55, 337, 494.

Bajra (see Pennisetum typhoideum).

bajulus, Hylotrupes.

bakeri, Anuraphis.

Balanephagus gica (see Balanoecus). Balaninus (see Curculio).

Balanoecus gica, sp. n., on Cocos coronata in Brazil, 468.

Balsa (see Ochroma).

Balsam Fir (see Abies balsamea). Baluchistan, Tortricids on fruit

trees in, 267, 518.
Bamboo, Coccids on, in tropical America, 471, 472; Dinoderus minutus in, in Fiji, 170; Dinoderus in, in India, 268; susceptibility of species of, to D. minutus in Porto Rico, 472.

Bamboo Scales (see Asterolecanium bambusae and A. miliaris).

Bambusa (see Bamboo).

bambusae, Asterolecanium. Banak (see Virola merendonis).

Banana, Aphids and virus diseases of, in Australia, 57, 58, 149; other pests of, in Queensland, 436, 465, 466; Dacus ferru-Dacus ferrugineus on, in Ceylon, 594; Cast-niid on, in Colombia, 8; fruit-flies not pests of, in Fiji, 495; moths damaging fruit of, in Gold Coast, 107; pests of, in Guam, 331; pests of, in Jamaica, 445; Cosmopolites sordidus on, in New Caledonia, 391; Icerya sey-chellarum on, in Seychelles, 82; locusts damaging, in Uganda, 228, 229; restrictions on importation of fruit of, into U.S.A., from Fiji, 495.

Banana Rust Thrips (see Scirto-

thrips signipennis).

Banana Weevil Borer (see Cosmo-polites sordidus).

Bandicota, destroying Rhynchophorus ferrugineus in India, 270.

Banding, against Cydia pomonella, 71, 279, 580; against other Lepidoptera, 75, 372; against Pseudococcus adonidum, 280; treatment of materials for, 279. (See Adhesives.)

banksi, Dirhinus.

Barbados, termites of, 5–7; biological control of sugar-cane pests in, 3–5, 7, 160.

Barbasco, pH value of diluents affecting toxicity of dusts containing, 301.

Barium, tests of compounds of, in baits for *Thermobia* and locusts,

269, 406, 614.

Barium Chloride, spraying with,

479, 575, 584.

Barium Fluosilicate, dusting with, 162, 163, 318, 430, 561; in baits, 237, 269, 406; spraying with, 190, 561; tests of toxicity of, to Lepidopterous larvae, 481, 600; formulae containing, 162, 237, 269, 318.

Bark-beetles, of conifers in Japan and Sakhalin, 96, 480; effects of felling and logging on incidence of, 323, 357, 382, 383; relation of, to Ophiostoma ulmi, 116, 119, 166, 212, 364, 365, 502, 550, 563, 606, 644, 645, 646; other fungi associated with, 96, 97, 116, 165, 357; digestive enzymes of, 340; new species of, 528, 592.

Barley, Toxoptera graminum on, in Argentina, 36; Coccid on roots of, in Egypt, 30; pests of, in India, 81, 266, 523; Criocerid experimentally feeding on, in Japan, 93; pests and virus diseases of, in Russian Union, 13, 14, 16, 574, 576, 617; pests of, in U.S.A., 251, 254, 356, 603.

Barley (Stored), pests in, and their control, 72, 73, 436, 576; suitability of types of, for Calandra granaria, 454, 455; olfactory reaction of Rhizopertha dominica to, 451.

Barriers, against Anabrus simplex, 537; against Blissus leucopterus, 245, 277; against cutworms, 428; against locusts, 289, 498, 529.

basalis, Coccophagus; Microctonus (Perilitus).

basizonus, Microcryptus.

Bassus [Braconid] (see Microdus).
batatae, Euscepes (see E. postfasciatus).

Bathyplectes curculionis, bionomics and establishment of, against Hypera variabilis in California, 401, 409, 410.

Batrachedra perobtusa, bionomics and control of, on coconut, etc.,

in Brazil, 9, 10.

Bats, carriage of Aspidiotus perniciosus by, 272; destroying Polyphylla perversa, 509.

Baumspritzmittel, 70, 231. (See Tar Distillates).

Bay Oil, 90.

Bean Beetle, Mexican (see Epilach-

na varivestis).

Beans, pests of, in S. Africa, 320, 615; pests of, in Australia, 31, 466, 495; pests of, in green-houses in Britain, 145, 146, 525; pests of, in Canada, 90, 330; in Canada, 90, 330; pests of, in Australia, 31, Agromyza huidobrensis on, in Chile, 512; Coccid on roots of, in Egypt, 30; pests of, in Hawaii, 35, 273, 650; new Jassid on, in India, 266; Bruchus obtectus on, in Jugoslavia, 72; Epilachna varivestis on, in Mexico, 114; Agromyza on, in Nyasaland, 154; Pyralids on, in Porto Rico, 345; locusts damaging, in Uganda, 228; pests of, in U.S.A., 39, 114, 122, 158, 162, 208, 214, 223, 247, 248, 252, 431, 439, 476, 477, 599, 601, 603; Thrips tabaci transmitting virus disease to, 140; susceptibility of varieties of, to pests, 273, 650; survival of Pyrausta nubilalis on, 43; insecticides and injury to, 162, 208, 249, 260, 441, 584, 599.

Beans (Stored), Bruchids in, 72, 148, 220; (green), fumigation against introduction of Trypetids into U.S.A. from Hawaii, in, 499.

Beans, Lima (see Lima Beans).
Beans, Mungo (see *Phaseolus mungo*).

Beans, Soy (see Soy Beans).

Beans, Velvet (see Stizolobium).

Beauveria, possibly infesting Dendroctonus frontalis in Virginia, 116. Beauveria bassiana, experimental

utilisation of, against insects, 368, 583.

beckii, Lepidosaphes.

Beech, Xyleborus germanus on, in New Jersey, 551.

Beer Must, infected with Beauveria bassiana in baits for Agrotis segetum, 583.

Bumble, parasitised

Myopa in Europe, 474.
Bees, Honey, new Acarapis on, 528; insect enemies of, 343, 473, method of protecting, 611; against Bufo marinus, 8; fumigation of chimney against, 418; transmitting fire-blight, 112; risk of poisoning, by insecticides, 136, 407, 549, 550, 642.

Bees, Wild Rock (see Apis dorsata).

beesoni, Calosoma.

Beet, Jassid on, in Argentina, 608 (and note); pests of, in Australia, 31, 237; Cleonus mendicus on, in Italy, 316; pests of, in Russian Union, 455, 456, 569, 577, 583, 584, 586, 654; Agrotis ypsilon on, in Sakhalin, 100; pests of, in U.S.A., 46, 47, 122, 134, 136, 161, 429, 441, 533, 537, 563, 631; insects and virus diseases of, 65, 66, 134, 179, 180, 438, 501, 608, 631; Pyrausta nubilalis not surviving on, 44; preparation of extracts of, for use with insecticides against Cleonus punctiventris, 586, 587.

Leafhopper (see Eutettix

tenellus).

Beet Webworm (see Loxostege sticticalis).

Beet Webworm, Hawaiian (see Hymenia recurvalis).

Beet Weevil (see Cleonus punctiventris).

Begonia, mite on, in California, 536.

Bell Pepper (see Capsicum). bellicosus, Macrotermes (Termes).

Bemisia gossypiperda (see B. tabaci). Bemisia inconspicua, on cotton in

Philippines, 649.

Bemisia tabaci, food-plants and relation to virus diseases of, in India and Sumatra, 183, 318, 523; food-plants of, in Sudan, 387. benefica, Pseudoleucopis.

beneficiens, Telenomus (Phanurus). Bengal, pests of jute in, 446.

Bent Grass (see Agrostis).

Bentonite, as a diluent for dinitroo-cyclohexylphenol dust, 249 uses of, in sprays, 45, 113, 129, 160, 239, 351, 352, 425, 633, 634, 638, 639, 645; study of types of, for preparation of nicotine-bentonite sprays, 558; formulae containing, 634, 638, 639.

Benzene, ineffective for treating tobacco plant-beds against Epitrix, 404; and paraffin wax, as a dispersing agent in sprays, 112; as a solvent, 147, 586.

Benzyl Alcohol, as a solvent for

rotenone, 237.

bergi, Neomaskellia. bergrothi, Helopeltis.

Bermuda, miscellaneous pests in, 138, 139; insects and virus diseases of plants in, 138, 646; list of insects in, 376; Macrodactylus subspinosus intercepted on roses in, 139.

Bermuda Grass (see Cynodon dacty-

lon).

Bermuda Palmetto (see Sabal bermudiana).

Berseem (see Trifolium alexandrinum).

Bessa selecta (see Ptychomyia).

β, β-dichloroethyl Ether, in sprays against gladiolus thrips, 136. Betula (see Birch).

betulae, Trypodendron.

bezziana, Stomatomyia. Bhindi (see Hibiscus esculentus).

bianchii, Organothrips. bicarinatus, Neoplectrus.

bicolor, Callitula; Echthrodelphax; Galerucida; Leefmansia.

Bidens pilosa, mite on, in S. Africa, 320; flea-beetle on, in Nyasaland, 588.

bidens, Picromerus.

bidentatus, Ips (Pityogenes).

bifasciata, Comperiella.

bifasciatus, Anastatus; Hylotrupes. Big-bud Mite, Filbert (see Eriophyes avellanae).

bigsbyana, Calligrapha multipunctata.

billoti, Hyperaspis.

bimaculata, Sympiesis.

bimaculatus, Neoanisotylus; Tetranychus (see T. telarius).

binotata, Anomala; Empoasca.

bioculatus, Paratetranychus (Tetra-

nvchus).

Biological Control, review of work on, in Canada, 307; of greenhouse pests, review of work on, 309, 310; general papers relating to 219, 362, 397; of noxious plants, 17, 141, 143, 289, 334, 467.

bipunctalis, Psara (Pachyzancla). bipunctata, Adalia (Coccinella). bipunctatus, Nephotettix.

Birch (Betula), Argyroploce urticana on, in Britain, 283; pests of, in Canada, 37, 347, 348; pests of, in Latvia, 85, 86; Melolontha hippocastani on, in Russia, 571; Coleoptera on, in U.S.A., 165, 347; - fungi associated with Scolytid on, 165; Anobium punctatum in timber of, 234.

Birch Borer, Bronze (see Agrilus

anxius).

Birds, destroying noxious insects, 24, 110, 212, 217, 242, 243, 244, 271, 312, 313, 314, 374, 380, 395, 497, 507, 509, 563; carriage of Aspidiotus perniciosus by, 272; dispersing milky diseases of Popillia japonica, 125.

birmanicus, Pachydissus.

biselliella, Tineola.

bisignifer, Anthonomus.

Bitoma palmarum, sp. n., on Attalea funifera in Brazil, 468.

Bittacus occidentalis, predacious on Crambus vulvivagellus in Kansas, 505, 506.

Bitter Melon (see Momordica charantia).

bivittatus, Melanoplus.

bizonatus, Phymatioderus.

Black Leaf 155, 22, 117, 128, 563,

Black Scale, application of name, 564.

Blackberry, pests of, in Britain, 283, 287, 544.

Blaesoxipha (see Sarcophaga).

blanda, Terias; Zenillia.

Blaps spp., humidity and temperature reactions of, in Russia, 571, 573.

Blattella germanica, 562.

Blattella vaga, habits of, in U.S.A.,

Blissus insularis, measures against, damaging lawns and rice in Trinidad, 373.

Blissus leucopterus (Chinch Bug), on cereals in U.S.A., 245, 254, 277; factors affecting resistance of Sorghum to, 258, 299; measures against, 245, 277; tests of dinitro-orthocresol dusts on, 326.

Blister Beetle, Black (see Epicauta pennsylvanica).

Blood (Dried), in baits for Hylemyia cilicrura, 158.

Blood Albumin, uses and effects of, in sprays, 113, 349, 350, 411.

Blueberry, experiments with forms of Rhagoletis pomonella and, in Nova Scotia, 226; pests of, in

U.S.A., **165, 259.** Blueberry Thrips (see *Frankliniella* vàccinii).

Boarmia imparata (see B. selenaria). Boarmia infixaria, on Burma, 77.

teak Boarmia selenaria, on in Burma, 77; on Melia azedarach in India, 74.

boetica, Cosmolyce (Lycaena).

Boletus tropicus, symbiosis of Pseudococcus comstocki and, Citrus in Brazil, 167, 168.

Bombay Locust (see Patanga succincta).

Bombus (see Bees, Bumble).

Bombyx mori (Silkworm), Dipterous parasites and mite attacking in Japan, 88, 98, 172; Trogoderma versicolor damaging cocoons of, 453; tests of insecticides on, 206, 207, 242.

bondari, Derelomus; Himatidium; Tonesia.

Books, insects damaging, and their control, 536.

Borax, as a timber preservative against Lyctus, 147, 447, 448; in mixture against termites, 610; in baits for Trypetids, 183, 462; unsatisfactory against Calandra in stored wheat, 541.

bordagei, Apanteles.

Bordeaux Mixture, against Empoasca spp., 148, 163, 325, 326, 441; study of action of, on E. fabae, against Haltica ampelo-440; phaga, 458; as a repellent for Lamellicorn beetles, 98, 171; against Pyralids, 10, 392; effect of, on Coccid populations on Citrus, 648; formulae for, 191, 248, 325, 441, 503, 558, 602, 605, 634; formulae containing, 148, 325, 503, 558, 580, 602, 605, 634; and arsenicals, 10, 163, 171, 325, 326, 412, 426, 503, 558, 580, lessening risk of foliage 634; injury by arsenicals, 426; relation of, to lead-arsenate spray residues on apple, 412; nicotine sulphate, 127, 148, 191, 381, 558; fixed nicotines incompatible with, 558; and oil emulsions, 441, 597, 634, 648; as an emulsifier for oils, 191, 248, 605; and sulphur, 597, 602.

borealis, Cyclocephala; Epilachna.

borgmeieri, Adesmus.

Borreria verticillata, relation of Larra americana to, in Porto Rico, 553.

Bostrychids, digestive enzymes of, 340.

Bostrychopsis eremita, on apple in Peru, 511.

Bostrychopsis jesuita, bionamics and control of, in New South Wales, 335.

Bostrychopsis uncinata var. orbignyi, on Eucalyptus in Argentina, 512. Bostrychulus (see Micrapate).

bothynoderis, Xenocrepis (Caeno-crepis).

botrana, Polychrosis.

Box (Buxus), pests of, in U.S.A., 636.

Box-elder (see Acer negundo). brachartonae, Anastatoidea.

brachialis, Bruchus.

Brachycaudus helichrysi (see Anu-

raphis padi).
Brachycerus undatus, parasitised by

Rondania cucullata in Italy, 316. Brachyckiton populneus (Kurrajong), value and pests of, in New South Wales, 467.

Brachycolus brassicae (see Brevicoryne).

Brachylybas variegatus, on tomato, pepper and taro in Fiji, 390.

Brachymeria, parasite of Dichomeris eridantis in India, 74; parasite of Tortrix postvittana in Tasmania, 151, 152.

Brachymeria apicicornis, hyperparasite of Artona catoxantha in Java,

79.

Brachymeria carinatifrons, parasite of Paradexodes epilachnae in Mexico, 115.

Brachymeria comitator, parasite of Alabama argillacea in Colombia, 53.

Brachymeria euploeae, parasite of Lepidoptera in Burma, 77; bionomics of, in Sumatra, 105, 106.

Brachymeria lugubris, hyperparasite of Artona catoxantha in Java and Malaya, 79.

Brachymeria nephantidis (parasite of Nephantis serinopa), bionomics and value of, in Ceylon, 621.

Brachymeria obscurata, parasite of Pieris rapae in Hawaii, 34, 35. Brachymeria ovata, parasite of Alabama argillacea in Colombia, 53.

Brachymeria punctiventris, hyperparasite of Artona catoxantha in Java and Malaya, 79. Brachymeria tegularis, hyperparasite of Melanoplus differentialis in Illinois, 401.

Brachyrrhinus (see Otiorrhynchus).
Bracon albolineatus (see Microbracon).

Bracon desertor, parasite of Hapalia machaeralis in Burma and India, 77.

Bracon hebetor (see Microbracon).
Braconids, of Russian Union, 480.
Bran (including shorts), in baits, 20,
100, 138, 170, 171, 223, 252, 269,
283, 300, 304, 354, 355, 357, 362,
390, 428, 429, 461, 538, 542, 548,
614, 615; baits for Trypetids
prepared with, 183, 462, 464.

brasiliensis, Gasterocercodes.

Brassica chinensis (Chinese Cabbage), pests of, in Hawaii, 651.

brassicae, Aleurodes (see A. proletella); Brevicoryne (Brachycolus); Hylemyia (Chortophila, Delia); Mamestra; Pieris; Plusia (Autographa).

Brassolis sophorae, on coconut in

Br. Guiana, 538.

Brazil, pests of field and stored cacao in, 391, 392, 616; pests of Citrus in, 167, 168, 392, 624; pests of coconut and other palms in, 9, 10, 468, 469; cotton pests in, 10, 11, 404, 469, 623; pests of fig and Artocarpus in, 391, 392; miscellaneous pests in, 40, 292, 345, 511; new Agromyzid on potato in, 36; thrips and virus disease of tobacco and tomato in, 141, 608; beneficial insects in, 4, 5, 36, 168, 292, 469, 470, 625; establishment of Tetrastichus giffardianus against Ceratitis capitata in, 344; introduction of beneficial insects into other countries from, 4, 298, 355, 431, 470, 471, 553.

braziliensis, Agromyza.

Breadfruit (see Artocarpus communis).

Bremia, 315.

brethesi, Leptomastidea.

brevicollis, Dociostaurus crucigerus.

brevicomis, Dendroctonus. brevicornis, Microbracon.

Brevicoryne brassicae (on crucifers), in Britain, 145, 381, 384, 590; in Fiji, 496; in Germany, 435; in India, 266; in Victoria, 546; in Wisconsin, 135, 136; probable relation of, to virus disease of

cauliflower, 235; certain Coccinellids not predactions on, 486; measures against 135, 136, 381, 384; tests of pyrethrum extracts on, 586.

brevipennis, Anastatus (see A. antestiae).

brevipes, Pseudococcus.

brevis, Calotermes (Cryptotermes, Kalotermes); Phaneroptera.

brevistigma, Tetrastichus.

breviuscula, Rodolia

breweri, Podagrica (Nisotra).

Brinjal (see Egg-plant).

British Columbia, pests of forest and shade trees in, 328, 415; orchard pests in, 22, 23, 328, 330, 508; vegetable pests in, 22, 310, 330, 508; grasshoppers in, 241, 308, 309, 394; Polyphylla perversa in, 508; beneficial insects and biological control in, 308, 309, 311, 415, 417, 509; introduction of natural enemies of grasshoppers into Argentina from, 308.

British Empire, memorandum on storage of foodstuffs in, 139.

British Isles, forest pests in, 323, 382, 383, 593, 628; pests and diseases of willows in, 146, 288, 312; pests of grasses and turf in, 110, 314, 315, 375, 381, 524, 526; pests of greenhouse and ornamental plants in, 27, 28, 63, 144-146, 381, 489, 490, 525, 628; legislation against Diarthronomyia chrysanthemi in, 64; pests of hops and nuts in, 282; orchard pests in, 108, 281, 282, 284, 285, 286, 287, 304, 590; pests of small fruits in, 27, 28, 146, 281, 283, 287, 375, 544; pests of vegetable and root crops in, 28, 63, 109, 111, 144, 145, 146, 310, 313, 314, 329, 374, 375, 381, 383– **385, 525, 590**; pests of stored grain and cacao in, 186, 616, 617; beetles damaging timber in, 147, 178, 234, 284, 315, 447, 448, 593; Aleurodids of, 374; key to Lamellicorn larvae of, 480; leafmining Lepidoptera in, Locusta migratoria in, 322, 323 key to Ptinids of, 340; Aphids and virus diseases in, 27, 287, 329, 375; climatic factors affecting migration of Aphids in, 180; correlation of insect outbreaks with sun spots in, 63; beneficial insects in, 79, 80, 142, 282, 310, 313, 314, 315, 334, 374, 417, 489, 490, 526; supplements to revision of Tachinids of, 328, 592; beneficial insects imported into other countries from, 142, 310, 334, 417, 609.

Broccoli, pests of, in Britain, 144, 145, 384; pests of, in Hawaii, 651; wireworms on, in New Jersey, 122; effect of dichloroethyl ether on, 122.

bromeliae, Diaspis.

bromeliae, auct., Pseudococcus (see P. brevipes).

Bromus inermis, reservoir of mosaic of cereals in Siberia, 14.

Brontispa froggatti (see B. longis-sima).

Brontispa longissima (on coconut), in New Caledonia, **391**; bionomics and biological control of, in New Guinea Territory, **379**.

Brontispa mariana, on coconut in Saipan, 331; precautions against introduction of, into Guam, 331. brontispae, Tetrastichus (Tetrastichodes).

Brotolomia meticulosa, bionomics and control of, on greenhouse plants, etc., in Britain, 145, 525, 526.

Brown Solution (and derivatives), preparation and use of, in dust insecticides, 247, 248.

Bruchidius (see Bruchus).

Bruchocida orientalis, hosts and ecology of, in S. India, 522.

Bruchoctonus senex, parasite of Bruchus pisorum in Russia, 575. Bruchophagus, parasite of Lasioderma serricorne in Fiji, 391.

Bruchus, identity, bionomics and control of species of, infesting cowpeas in S. Africa, 19, 58, 321.

Bruchus analis (in cowpeas), possibly in S. Africa, 321; intercepted in Fiji, 148.

cepted in Fiji, 148.

Bruchus brachialis (Vetch Bruchid), in U.S.A., 47, 48, 430; hibernation quarters of, 48; introduction of parasites against, 48, 430; fumigation of vetch seed against, 47.

Bruchus chinensis, fumigation of stored cowpeas against, in S. Africa, 19; Bruchid recorded as, in cowpeas in S. Africa, 58, 321; in Cajanus seeds in Fiji, 496; parasite of, in South India, 522; tests of atomised sprays on, 434.

Bruchus elegans, on Schinus molle in Chile, 512.

INDEX. 689

Bruchus impressithorax, in stored peas in Gold Coast, 492.

Bruchus maculatus (quadrimaculatus), in stored peas in Gold Coast, 492; in U.S.A., 215, 566; effects of humidity on, in stored cowpeas, 566; Pediculoides ventricosus infesting, 215.

Bruchus marshalli, 321.

Bruchus obtectus, intercepted in beans in Fiji, 148; on field and stored beans in Jugoslavia, 72; Pediculoides ventricosus destroying cultures of, in U.S.A., 215; egg production and longevity of, 220; attempted rearing of Triaspis thoracicus on, 48.

Bruchus pisorum (on peas), proposed introduction of Triaspis thoracicus into W. Australia against, 142; in Russia, 575; measures against, in U.S.A., 408, 630, 631, 640; bionomics of, 575, 630; intercepted in broad beans in Fiji, 148.

Prachas

Bruchus pruininus, intercepted in seeds of Desmanthus in Fiji, 148. Bruchus quadrimaculatus (see B. maculatus).

Bruguiera gymnorrhiza, Parendymia pilipes in timber of, in Fiji, 170.

brumata, Operophtera.

Brumus octosignatus, hibernation and value of, against Aphids in Uzbekistan, 486.

Brumus suturalis, predacious on Pyrilla perpusilla in Punjab, 82. Brunfelsia hopeana, Thyridia themis-

to on, in Brazil, 292.

brunnea, Crematogaster.

brunneipennis, Hypera. brunneus, Luperodes (see Monolepta varicornis); Lyctus.

Brush Box (see *Tristania conferta*). Brussels Sprouts, *Hylemyia brassicae* on, in Britain, **109**, **384**; virus disease of, in Germany, **234**.

Bryobia, on chrysanthemum in Punjab, **269**.

Bryobia praetiosa, in U.S.A., 164, 211, 561; thrips predacious on, on almond, 211; entering houses, 164; dusts against, 164, 561.

Bryocoropsis laticollis, on cacao and Annona in Gold Coast, **492.**

Bucculatrix canadensisella, on birch in New Brunswick, 348.

Buckthorn Aphis (see *Aphis rhamni*).

Budmoth, Apple (see Spilonota ocellana).

Budworm, Spruce (see Harmologa fumiferana).

Buffalo Carpet Beetle (see Anthrenus scrophulariae).

Buffalo Grass (see Stenotaphrum). Bufo marinus, distribution and utilisation of, against noxious insects, 7, 93, 377, 389, 496, 603; artificial breeding of, 7; method of protecting bee-hives against,

Building Timbers, Coleoptera in, 86, 103, 139, 179, 316, 324, 333, 385, 497, 593; Siricids in, 385; termites in, 6, 19, 142, 333, 400, 473, 495, 537, 610.

Bulgaria, parasites of Lepidoptera in, 71, 514.

buoliana, Rhyacionia (Evetria).

Bupalus piniarius, unimportant on pine in Latvia, 86.

Buprestis, revision of N. American species of, 655.

Burma, forest pests and their biological control in, **75–79**; new *Calandra* in, **79**; new parasite of Pyralid in, **514**; parasites of teak pests introduced into India from, **75**, **77**.

bursarius, Pemphigus.

Bush Locust (see Phymateus leprosus).

Busseola fusca (on maize), rainfall favouring, in S. Africa, 19; in Nyasaland, 154.

Bussothrips claratibia, on cotton in Philippines, 649.

Butea frondosa, cultivation of lac on, in Siam, 94.

Buttons, *Coccotrypes dactyliperda* attacking imported, in Bermuda, **139.**

Butyl Carbitol Thiocyanate, 56, 108, 111, 434. (See Thiocyanates.)

Butylphenoxy Ethanol, Paratertiary, intermediary solvent for derris resins when used with oils, **350.**

buxi, Monarthropalpus; Pinnaspis; Psylla.

Buxus (see Box).

Bythoscopus cedaranus, dusts against, on wattle in S. Africa, 19, 278.

byturoides, Attagenus.

Byturus tomentosus (Raspberry Beetle), tests of pyrethrum sprays against, in Britain, 281.

C

c-nigrum, Amathes (Graphiphora). Cabbage, Plutella maculipennis on, in S. Africa, 19; pests of, in Australia, 142, 290, 436, 494, 546; Feltia subterranea on, in Bermuda, 138; pests of, in Br. Isles, 313, 375, 381, 383-385, 590; pests of, in Germany, 68, 231, 232, 450; pests of, in Hawaii, 34, 35, 650, 651; pests of, in India, 266, 271, 519; pests of, in Japan, 100, 173; Plutella maculipennis on, in New Caledonia, 391; Pieris rapae on, in New Zealand, 546; pests of, in Russian Union, 478, 585; pests of, in U.S.A., 122, 123, 135, 199, 214; Myzus persicae overwintering on, 329, 590; mosaic disease in, 234; effect of dichloroethyl ether on, 122; derris residues on, **555.**

Cabbage Aphis (see Brevicoryne brassicae).

Cabbage Moth (see Plutella maculipennis).

Cabbage Root Maggot (see Hylemyia brassicae).

Cabbage White Butterflies (see *Pieris brassicae* and *P. rapae*).

Cabbage Whitefly (see Aleurodes proletella).

Cables, Electric, Coleoptera boring in lead sheathing of, 335, 345; treatment against termites of tabric covering of, 142.

Cacao, pests of, in W. Africa, 492, 515, 516; pests of, in Brazil, 391, 392; Coccid on, in Dominican Republic, 498; locust attacking, in Formosa, 94; pests of, in Malaya, 512, 513; pests of, in Uganda, 175.

Cacao (Stored), pests in, 103, 493, 616, 617.

Cacoecia (see Tortrix).

Cacoecia fumiferana (see Harmologa).

cacoeciae, Euplectrus.

cacti, Chilocorus.

Cactoblastis cactorum (destroying Opuntia), establishment, utilisation and value of, in S. Africa and Australia, 17, 587.

cactorum, Cactoblastis.

Cadurcia (parasite of Artona catox-antha), parasites of, 79.

Cadurcia vanderwulpi, parasite of Dichomeris eridantis in India, 74.

Caenocrepis bothynoderis (see Xenocrepis).

caesalis, Margaronia. caffeina, Leucoptera.

Cages, for rearing and studying insects, 8, 308, 311, 344, 557.

Cajanus cajan (indicus), pests of, in Fiji, 390, 496; pests of, in India, 265, 267, 523; Helopeltis bergrothi on, in Nyasaland, 589; Empoasca lybica on, in Sudan, 387.

calacte, Macrocentrus.

Calamagrostis pseudophragmites, relation of Locusta migratoria to, in Russia, 570.

Calandra, in stored grain in Britain, 186; key to Indian species of, 79; review of data on, in stored wheat, 480; proposed retention of name, 79.

Calandra conicollis, sp. n., in Anda-

man Islands, 79.

Calandra erosa, sp. n., in Burma and Assam, 79.

Calandra granaria (in stored cereals), 434; in Australia, 380, 540; in Russian Union, 454, 455, 576; in Spain, 457; in U.S.A., 209, 477; suitability of different cereals for, 454, 455; effects of temperature and humidity on, 454; measures and experiments against, 209, 380, 457, 477, 540, 576.

Calandra oryzae (in stored cereals), in S. Africa, 19; in Australia, 380, 540; in Japan, 172; in Korea, 92; in U.S.A., 209, 477; factors affecting infestation of field and stored maize by, in S. Rhodesia, 184; in imported ground-nuts in Fiji, 496; in dried sweet potato, 173; effects of temperature and humidity on, 92, 172; measures and experiments against, 19, 185, 209, 380, 477, 540; tests of insecticides on, 434, 540.

Calandra vateriae, sp. n., in seeds of Vateria indica in India, 79. calandrae, Aplastomorpha.

Calcid, **336.**

calcitrans, Stomoxys.

Calcium Antimony Tartrate, in sprays against thrips, 204, 277, 296, 413.

Calcium Arsenate, against cereal pests, 99, 148, 300, 464, 600; against Sexava on coconut, 91, 92; against cotton pests, 3, 10,

11, 53, 257, 258, 358, 359, 428, 430, 510, 539, 599, 600, 624, 649; effect of, on infestation of cotton by Aphis gossypii, 3, 359; against forest pests, 36, 86; against pests of leguminous crops, 39, 252, 534; against orchard pests, 129, 171, 281, 406, 550, 580, 585, 641; against pests of solanaceous plants, 31, 138, 163, 330, 429, 503, 504, 534, 598, 631, 639, 640, 641; against Polychrosis viteana on vines, 117,558; against pests of other plants, 46, 165, 426, 547, 647, 649, 654; tests of toxicity of, to Lepidoptera, 356, 481, 584, 600; particle size affecting toxicity of, to honey bees, 550; in baits, 100, 170, 534, 598, 613; formulae for, in dusts, 10, 31, 252, 258, 358, 359, 504, 510, 547, 600, 640, 641, 647; carriers for, 53, 148, 252, 504, 598, 640, 641, 647; conditioning agents for dusts of, 600, 654; aeroplane dusting with, 300; spraying with, 10, 11, 36, 39, 86, 99, 117, 129, 148, 171, 281, 406, 426, 503, 534, 539, 558, 580, 585, 624, 641, 649, 654; adhesives for, 117, 406; ammonium alginate stabilising suspensions of, 584; and Bordeaux mixture, 163, 171, 426, 503, 558, 580; and ferrous sulphate, 281, 406; and lime, 10, 148, 406; and limesulphur, 171, 281; and oil emulsions, 585; and Paris green, 300; and sulphur, 10, 258, 358, 359, 510, 600, 639; forms of, 257, 359; and injury to plants, 252, 259, 281, 406, 426, 584, 599. Calcium Arsenite, in baits for

Calcium Arsenite, in baits for locusts, 612, 613; and lime, dusting with, against Longitarsus spp., 479; effect of, on midgut wall of Laphygma eridania, 481.

Calcium Bromacetate, in baits for locusts, 614.

Calcium Carbonate, electrostatic charges of dusts of, 569.

Calcium Caseinate, as a spreader and adhesive for sprays, 32, 117, 249, 441.

Calcium Chloracetate, in baits for locusts, 612, 613, 614.

Calcium Chloride, test of, in spray against thrips, 296; reducing ovicidal action of nicotine, 556; increasing oil deposits from sprays, 411.

Calcium Cyanamide, soil treatment with, against pests, 418, 512.

Calcium Cyanide, as a greenhouse fumigant, 32; form of, used for tent fumigation of Citrus against Aonidiella aurantii, 336; against pests of stored products, 19, 377, 610; against termites, 338; methods of using, against Anasa tristis, 201; in dust against Empoasca devastans, 517; in paint against Saperda candida, 243; chimney fumigated with, against honey-bees, 418.

Calcium Pitchate, action of dusts containing, on Epilachna vari-

vestis, 247.

Calcium Sulphate, not affecting toxicity of dusts containing rotenone, **301**.

Calepitrimerus baileyi, thrips predacious on, on apple in California, 210.

calidum, Calosoma.

California, Pseudococcus adonidum on avocado in, 297, 355; pests of Citrus in, 209, 222, 295, 297, 336, 349, 355, 386, 398, 399, 407, 410, 425, 535, 553, 561, 564, 565; pests of forest and shade trees in, 49, 357, 371, 416, 551; pests of lucerne in, 222, 223, 354, 401, 402, 403, 409, 441, 628, 629; pests of nut trees in, 211, 222, 223, 445, 425, 549, 561, 565; pests of ornamental plants in, 45, 114, 223, 536, 564, 565; pests and diseases of vegetables in, 222, 223, 224, 234, 301, 302, 429, 500, 501, 565, 608; miscellaneous pests in, 215, 221, 222, 416, 562, 629; Crambus spp. damaging lawns in, 408; Dinapate wrighti in, 137; grasshoppers in, 222, 223, 354; Rhagoletis spp. in, 222, 565; beneficial insects and biological control in, 86, 190, 210, 220, 280, 297, 310, 355, 385, 386, 401, 406, 409, 416, 417, 425, 442, 565; introduction of beneficial insects into other countries from, 9, 172, 310, 320, 457, 601, 651.

California Red Scale (see Aonidiella

aurantii).
californica, Melanophila.
californicus, Tenuipalpus.

Caliroa limacina, cherry defoliated by, in Japan, 96.

Calliceras (see Ceraphron).

C2

(1349)

Callidium violaceum, damaging building timbers in Sweden, 324.

Calliephialtes (see Ephialtes).

Calligonum polygonoides, relation of Schistocerca gregaria to, in India, 519.

Calligrapha multipunctata bigsbyana, bionomics of, in Quebec, 626.

calligraphus, Ips.

Callimome, parasite of Contarinia medicaginis in Hungary, 12.

Calliphora erythrocephala, permeability of cuticle of, by insecticides, 240.

Calliptamus italicus, in Russia, 570; in Spain, 529; measures against, in Turkey, 289.

Callistephus (see Aster).

Callitula bicolor, parasite of Mayetiola destructor in Russia, 618.

Callosobruchus (see Bruchus).

Caloglyphus julidicolus, on locusts in cages in S. Africa, 322. Calomel (see Mercurous Chloride).

Calosoma beesoni, predacious on Ectropis deodarae in India, 74.

Calosoma calidum, predacious on Cirphis unipuncta in U.S.A., 300. Calosoma maderae indicum, libera-

tion of, against Lepidoptera on Dalbergia sissoo in India, 74. Calotermes, attacking trees and

timber in Barbados, 6; treatment of timber against, Bermuda, 138.

Calotermes approximatus, timbers resistant to, in Bermuda, 138.

Calotermes brevis (attacking timber), in Bermuda, 138; in Florida, 158; in W. Indies, 5, 6, 157; resistance of different timbers to, 157; measures against, 6, 138.

Calotermes domesticus, investigation of damage to sepetir wood by, in Malaya, 543.

Calotermes minor, in Utah, 400.

pubescens, attacking Calotermes trees and timber in Porto Rico, 473.

Calotermes repandus, in timber in Fiji, 495, 496.

Calotermes snyderi, attacking trees on Mona Island, 473.

Calotropis gigantea, question of repellence of, to Schistocerca gregaria, 519.

Calystegia (see Convolvulus). cambriensis, Sitodiplosis.

Camellia, Coccids on, in nurseries in California, 223.

camelliae, auct., Aspidiotus (see A. rapax).

camelliae, Lepidosaphes.

Camnula pellucida, in Canada, 241, 309, 393, 394; in U.S.A., 205, 223, 604; bionomics of, 604; effect of hailstorms on, 394; aggravating soil erosion, 205; measures against, 241, 393; introduced parasite reared on, 309. Campaea margaritata, habits of,

on apple in Britain, 282.

Campsomeris, 84.

Campsomeris ciliata, parasite of Melolonthids in Algeria, 83. Campsomeris klugi, parasite of Poly-

phylla fullo in Russia, 572.

Campsomeris phalerata, liberation of, against Lachnosterna smithi in Mauritius, 609.

Campsomeris sexmaculata, parasite of Anoxia pilosa in Russia, 572. Camptothlipsis furtifica, parasite of Dichomeris eridantis in India, 74.

Campylomma nicolasi, apparently causing "crazy top" on cotton

in Sudan, 387.

Canada, forest pests in, 37-39, 293, 294, 307, 312, 328, 347, 348, 395, 474, 530, 531, 532; grasshoppers in, 144, 308, 309; Lachnosterna spp. in, 89; miscellaneous pests in, 161, 310, 328, 393; pests and disease of rose in, 532; key to Ptinids in dwellings and warehouses in, 152; beneficial insects and biological control in, 36, 37, 38, 39, 90, 293, 294, 307, 308, 309, 310, 530, 531, 532, 552; review of use of aircraft for insect control in, 152; directory of insect pest control in, 438; beneficial insects introduced into other countries from, 308, 637.

canadensis, Exenterus; Spathius.

canadensisella, Bucculatrix.

Canary Fly (see Typhlocyba frog-

gatti). Canary Islands, Ceratitis capitata in. 459.

Canavalia spp., Pyralids on, in Porto Rico, 345. candida, Saperda.

canella, Diatraea; Paria.

Canna, locust attacking, in Formosa, 94; relation of species of, Aphid-borne virus disease of Musa spp. in Australia and Philippines, 57, 317.

Cantaloupe (see Melon)

Cantheconidea acuta, C. gaugleri recorded as, in Sumatra, 105.

Cantheconidea gaugleri, sp. n. (predacious on Oreta carnea), bionomics of, in Sumatra, 105.

Weed (see Cryptostemma Cape

calendulaceum).

capensis, Coccophagus; Microphthalma (see M. sejuncta); Nudaurelia cytherea.

capitata, Ceratitis.

Capitophorus, identity of species of, on strawberry, rose and Potentilla, 27: Pentatrichopus considered distinct from, 27.

Capitophorus fragaefolii, status of,

27.

Capitophorus fragariae, on straw-berry in Britain, 27, 283, 375; transmitting virus diseases strawberry, 27; experiments with, on other plants, 28; bionomics and systematic position of, 27; measures against, 283, 375.

Capitophorus potentillae, on Potentilla anserina in Britain, 28; experiments with, on other plants, 28; question of syno-

nymy of, 27.

Capitophorus tetrarhodus, on roses in Britain, 27, 28; experiments with, on other plants, 28.

Capnodis tenebrionis, on stonefruit trees in Jugoslavia, 72.

Pseudococcus com-Capnodium, stocki associated with, on Citrus in Brazil, 168.

Capparis aphylla, relation of Schistocerca gregaria to, in India, 519. Capparis sandwichiana, Murgantia histrionica on, in Hawaii, 35.

Capsicum, Gnorimoschema operculella on, in Bermuda, 138; Tarsonemus latus on, in California, 223; Dacus cucurbitae on, in Malaya, 90; Lineodes integra on, in Peru, 539; insects and virus disease in, 140, 456; virus disease of egg-plant not transmitted to, 542; fumigation against introduction of Trypetids into U.S.A. from Hawaii in fruits of, 499.

Caragana, grasshoppers damaging,

in U.S.A., 603.

Carambola (see Averrhoa carambola). Carbolic Acid, in dust against mites, 236; in paint against Scolytids, 170.

Carbolineum, types of, 231, 232. (See Tar Distillates.)

Carbon Bisulphide, against ants and termites, 169, 338; tool for use in fumigating ants' nests with, 40; against other pests in soil, 24, 111, 145, 168, 199, 203, 237, 299, 300, 348, 374, 428, 498, 568, **579, 580, 614;** diffusion of, in soil, 580; against Blissus insularis, 373; injected into apple trees against Saperda candida, 243; causing injury to plants, 145, 300, 428; fumigation with, against pests of seeds and stored products, 47, 59, 95, 139, 182, 278, 327, 390, 429, 454, 537, 611; effect of, on germination of seed, 59, 95; in spray against Dermestids, 454; use of, for dipping eggs of Sitotroga, 193; effect of ether on toxicity of, to Tribolium confusum, 327; and naphthalene, 111; and paradichlorobenzene, 243, 579; emulsions of, 111, 145, 168, 299, 300, 348, 373, 498.

Carbon Dioxide, effects of, on loss of water and respiration in insects, 302, 452; uses of, to destroy pests of stored grain, 185, 380; relation of time, concentration and toxicity in tests of, 434; fumigants used with, 85, 452, 536.

Carbon Tetrachloride, effect of ether on toxicity of, to Tribolium confusum, 327; and ethyl acetate, against Dermestids in floorboards, 393; and ethylene dichloride, against pests of stored grain, 278, 429; unsatisfactory against Otiorrhynchus in soil, 198; for preparation of Tephro-

sia extracts, 586. Carcelia (see Zenillia).

Cardamine hirsuta, Pieris rapae on, in Japan, 173.

Cardamom (see Elettaria cardamomum).

Poecilips (Thamnurcardamomi, gides).

cardinalis, Rodolia (Novius, Veda-

cardui, Aphidius; Vanessa.

carduorum, Rhizotrogus.

Carex, Septis finitima on, in Maine, 560.

caridei, Sarcophaga.

carinatifrons, Brachymeria.

carinatus, Tetrastichus.

Carissa ovata, Hypsid defoliating, in Queensland, 467.

Carnation, Tortricid on, in greenhouses in Britain, 146; pests of, in U.S.A., 207, 214, 413, 641; low absorption of selenium from nutrient solutions by, 207.

carnea, Oreta.

Carneocephala flaviceps, on rose in

U.S.A., 256. Carolina, North, new mites on blueberry and strawberry in, 259, miscellaneous pests in, 48, 49; tobacco pests in, 201, 218, 365, 404; pests of vegetables in, 158, 201, 218, 560; pests of stored products in, 365, 429, 566; beneficial insects and biological control in, 48, 218, 400, 560.

Carolina, South, miscellaneous pests in, 49, 253, 257, 425; tobacco pests in, 253, 404; pests of vegetables in, 158, 199, 253.

carolina, Dissosteira.

carpatus, Apanteles.

Carpet Beetles (see Anthrenus and Attagenus piceus).

Carpets (see Textiles).

carpini, Asterobemisia (Aleurodes). Carpocapsa (see Cydia).

carpocapsae, Ascogaster (see A. quadridentata).

Carpoglyphis lactis, in dried fruits and wine, 488; deutonymph of,

Cryptophlebia carpophaga, (see Argyroploce illepida).

Carpophilus, habits and control of species of, attacking damaged pineapples in Fiji, 390.

Carpophilus dimidiatus, 390; stored cacao in Brazil, 616,; in imported ground-nuts in Fiji, 496.

Carpophilus hemipterus, 390. Carpophilus humeralis, 390. Carpophilus mutabilis, 390.

Carrot, Feltia subterranea on, in Bermuda, 138; susceptibility of, to Lachnosterna in Canada, new Jassid on, in India, 266; insects on, in U.S.A., 46, Listroderes obliquus on, in Victoria and Hawaii, 31, 35. Carthamus tinctorius, pests of, in India, 266, 523.

caryae, Acrobasis; Halisidota. caryaefoliae, Melanocallis.

caryana, Enarmonia (Laspeyresia). Casein, as a spreader for sprays, 145, 238; as an emulsifier for oils, 241. (See Ammonium and Calcium Caseinates.)

Cashew (see Anacardium occidentale).

Cassava, pests of, in S. America, 9, 169; Lepidosaphes alba on, in Dominican Republic, 498; L. dispar on, in Gold Coast, 492; locusts damaging, in Uganda, 228; experiments with Zeuzera coffeae and, 271.

Cassia, pests of, in Burma, 78. Cassia fistula, pests of, in India,

265, 271.

Cassia occidentalis (see Ditremena). Castalius rosimon, on Zizyphus jujuba in Siam, 94. castanea, Aserica (Autoserica);

Diparopsis; Pentilia.

castaneum, Tribolium. castanoptera, Xylopsocus.

Castnia atymnius r. humboldti, bionomics of, on banana and sugar-cane in Colombia, 8.

Castor, Tarsonemus latus on, in S. Africa, 320; Xyleborus fornicatus on, in Ceylon, 381; pests of, in India, 266, 271, 516; Parasa vivida on, in Nyasaland, 60; reported toxicity of, to Popillia japonica, 118; of no value against grasshoppers, 327. Castor Oil (see Oils, Vegetable).

Casuarina, Icerya purchasi on, in Dominican Republic, 498; I. seychellarum on, in Seychelles, 82; distillate of leaves of, in baits for locusts, 612.

catalaunalis, Antigastra.

Catalpa, pests of, in U.S.A., 195, 218.

catalpae, Ceratomia.

Catascopus facialis, predacious on Oryctes rhinoceros in Malaya, 148; introduced into Samoa, 148.

catenatus, Orthocraspeda.

Cathartus advena (see Ahasverus). catocaloides, Achaea.

Catolaccus aeneoviridis, parasite of Parornix prunivorella in Kansas, 191.

Catolaccus endonis, sp. n., parasite of Anthonomus bisignifer Japan, 40.

Catolellerus strangulatus, on coconut in Brazil, 469.

Catopsilia crocale, on Cassia in Burma, 78.

Catorama herbarium, in upholstery in Fiji, 496.

catoxantha, Artona.

catoxanthae, Eupelmus.

caudata, Lepidiota.

caudatus, Dacus (Chaetodacus). caulicola, Rhaconotus.

Cauliflower, pests of, in Australia, 436, 494, 496, 546; pests of, in Britain, 109, 145, 381, 384; Brevicoryne brassicae imported into Fiji on, 496; Aphids and virus disease of, in Germany, 234, 235; Plusia brassicae on, in Hawaii, 651; B. brassicae on, in India, 266; wireworms on, in New Jersey, 122, 123; experiment with virus disease of other plants and, 140; effect dichloroethyl ether on, 122. effect of

cautella, Ephestia.

Cavariella aegopodii, on strawberry

in Britain, 28.

Cecidomyiids, studies on, affecting seed-production of grasses in Britain, 314, 315, 375; of West Indies, 468.

Cecidostiba dendroctoni, parasite of Dendroctonus frontalis in Vir-

ginia, 211.

Cedar, Spanish (see Cedrela).

cedaranus, Bythoscopus.

Cedrela, Hypsipyla grandella on, in

Porto Rico, 427. Cedrela toona, billets of, for trapping Pagiophloeus longiclavis, 75. Cedria paradoxa, utilisation against *Hapalia machaeralis* in Burma, **77**.

Cedrus deodara, Ectropis deodarae on, in India, 74.

celerio, Hippotion.

Celery, virus diseases in, 57, 140.

Cellophane, penetration of, by larvae of Lasioderma, 493.

celsissimus, Phytodietus. celtis, Selepa (Plotheia).

Centaurea, Trypetid on, 490.

Centrodora speciosissima, parasite of Mayetiola destructor in Russia,

Cephaleia, on spruce in Quebec, 36.

cephalonica, Corcyra.

Cephalosporium lecanii, experiments with, against Ceroplastes sinensis in Russia, 582.

cephalotes, Atta.

Cephus cinctus (in Canada), cultural practices affecting, on wheat, 112; factors in resistance of food-plants to, 393.

cera, Xanthopimpla.

Cerambycids, digestive enzymes of, 340.

ceramicus, Xyleutes.

Ceraphron, parasite of Perisierola in Ceylon, 621; parasite of Mayetiola destructor in Russia,

Ceraphron kamiyae, hyperparasite of Lymantria dispar in Japan,

Ceraphron parasite vagans, Mayetiola destructor in Russia, 618, 619; male of, 618.

cerasi, Myzus; Rhagoletis. cerasivorana, Macrocentrus;

trix (Cacoecia).

Cerastium arrense, Galeruca tanaceti on, in Germany, 68.
Ceratitis capitata, in S. Africa, 18; in Algerian Sahara, 616; in Brazil, 344; in Canary Islands, 459; in Hawaii, 458, 459; in New South Wales, 55; in Spain, 457, 459, 461, 462, 464; legislation against importation of, into U.S.A. from Hawaii, 499; on coffee, 344; on citrus, 462, 616; on other fruits, 18, 462, 464, 616; bionomics of, 18, 459; parasites and biological control of, 344, 457, 458, 459; tests of baitsprays, baits and traps for, 55, 461, 462; factors vitiating tests of baits and traps against, 464; bags for protecting peaches against, 464; fumigation of harvested fruit, etc., against, 18, 499.

Ceratoma trifurcata, relation of, to cowpea mosaic in U.S.A., 503.

Ceratomegilla fuscilabris, predacious on Colias eurytheme in Louisiana, 600.

Ceratomegilla maculata, predacious on noxious insects in U.S.A., 251, 534.

Ceratomia catalpae, parasite and hyperparasites of, on Catalpa in N. Carolina, 218.

Ceratosmicra, hyperparasite of Protoparce sexta in N. Carolina, 219. Ceratostomella piceae, bark-beetles associated with, on spruce in Sakhalin, 96.

Ceratostomella ulmi (see Ophiostoma).

Ceratovacuna lanigera (see Oregma). cercerisella, Fascista (Gelechia).

Cercis canadensis, pests of, Kansas, 632, 635.

Cercopis sanguinea (see Tomaspis). Cerdana alliodora, Monanthia monotropidia on, in Porto Rico, 51. cerealella, Sitotroga.

cerealellae, Habrocytus. Cereals, Sitotroga cerealella on, in Jugoslavia, 72; Lema melanopa on, in Spain, 464; pests of, in Germany, 68, 449; pests of, in Russian Union, 13, 15, 569, 571, 574, 576, 582, 618; pests of, in U.S.A., 245, 254, 255, 277, 300, 327, 428, 537, 605; type of injury to, by Eurygaster integriceps, 574; insects and mosaic diseases of, 13, 15, 16. Maize, Wheat, etc.)

Cereals (Stored), pests in, and their control, 19, 72, 97, 102, 139, 173, 184, 186, 209, 222, 254, 277, 278, 288, 289, 380, 429, 436, 451, 453, 454, 457, 477, 493, 506, 537, 540, 553, 569, 576; book on control of pests of, in U.S.A., 605; absorption of methyl bromide by,

444.

cerigoides, Dasychira.

Ceromasia lepida, considered correct name for C. senilis (q.v.),

Ceromasia senilis, establishment of, against Pyrausta nubilalis Guam, 331; correct name for, 331.

cerophaga, Angitia.

Ceroplastes, new parasite of, in Argentina, 344; on Citrus in Transcaucasia, 653; resistant to oil emulsions, 653.

Ceroplastes rubens (on Citrus), parasites and attempted biological control of, in Japan, 172; measures against, in Queensland, 437.

Ceroplastes sinensis, experiments Cephalosporium with against, on Citrus in Russia, 582.

ceroplastis, Cheiloneurus.

cervina, Thosea.

cervinatus, Oxycanus.

Cetonia aurata, Scoliids reared on, in Russia, 572.

Ceuthorrhynchus assimilis. parasite of, on turnip in Washington, 569.

Ceuthorrhynchus picitarsis, on rape in Germany, 235.

Ceuthorrhynchus quadridens, on rape

in Germany, 235.

Ceylon, Nephantis serinopa on coconut in, 594, 619-622; miscellaneous pests in, 594; tea pests laneous pests in, 594; in, 60, 173, 174, 175, 381; probable virus disease of tea in, 175; pests of stored cacao in, 616;

subspecies of Xyleborus fornicatus in, 381; cultivation and pests of lac in, 594; beneficial insects and biological control in, 174, 514, 594, 610, 619, 620, 621; Mermithids infesting 621; Homona coffearia in, 173; beneficial insects introduced Mauritius from, 610; pests imported into Fiji in ground-nuts from, 496.

ceylonica, Fornicia. ceylonicus, Dactylopius.

Chaetocnema spp., relation of, to bacterial wilt of maize in U.S.A., 244, 245.

Chaetodacus (see Dacus).

Chaetolixophaga laspeyresiae, gen. et sp. n., parasite of Cydia molesta in Argentina, 169.

Chalcidini, revision of N. American, 189.

Chalcis, parasite of Dichomeris eridantis in India, 74. Chalcis, auct. (see Brachymeria).

chalcites, Plusia (Autographa, Phytometra).

Chalcodermus aeneus (Cowpea Curculio), bionomics of, in U.S.A., 405, 598; measures against, 598.

Chalepus dorsalis, bionomics and control of, on soybeans in U.S.A., 354.

Cheese, absorption of methyl bromide by, **444**.

Cheiloneurus ceroplastis, hyperparasite of Ceroplastes rubens Japan, 172.

Cheiloneurus elegans, parasite of Mayetiola destructor in Russia, 618.

Cheiloneurus longisetaceus sp. n., parasite of Coccus hesperidum in Argentina, 445.

Cheiloneurus pyrillae, parasite of Pyrilla in India, 183, 184.

Chelonella (see Chelonus).

Chelonus cycloporus, parasite of Holcocera pulverea in India, 264. Chelonus ritchiei, parasite of Platy-

edra gossypiella in Nyasaland, 153.

Chelonus tabonus, parasite of Margaronia pyloalis in Japan, 34. Chenopodium, pests on, in U.S.A.,

46, 441. Cherimoya (see Annona cherimolia).

Chermes abietis, on spruce Quebec, 36.

Chermes cooleyi, on Picea spp. in Montana, 537.

Chermes nordmannianae, 67.

Chermes pineoides, on spruce in Switzerland, 67: derris dust unsatisfactory against, 67.

Chermes pinifoliae, on Picea engelmanni in Montana, 537.

Chermes strobi, characters distinguishing C. pineoides and, 67. chermesina, Rodolia.

Cherry, pests of, in Australia, 149, 291, 335; Phyllocoptes fockeui on, in Austria and France, 304; Tortricids on, in Baluchistan, **267**, **518**; pests of, in Canada, 328, 508, 625; pests of, in Germany, 231; pests of, in Japan, 96, 622; Rhagoletis cerasi on, in Jugoslavia, 72; Lachnosterna morosa on, in Korea, 171; Scolytus rugulosus on, in Peru, 539; pests of, in U.S.A., 2, 48, 190, 196, 222, 415, 565, 639, 641; spray schedule for, in Kentucky, 2; susceptibility of species of, to Parornix prunivorella, 190; Rhagoletis pomonella developing on, 226; arsenicals causing injury to, 196; problem of residues on fruit of, 196.

Cherry Fruit Worm (see Cydia

packardi).

Cherry Fruit-flies (see Rhagoletis cerasi, R. cingulata and R. fausta). Cherry Tent Caterpillar (see Tortrix

cerasivorana).

Chestnut, pests and fungus disease of, in Japan, 97, 99; Cydia latiferreana on, in U.S.A., 405.

Chick Pea (see Cicer arietinum).

Chickweed (see Stellaria media).

chiclayae, Anastrepha. Chicory, Thrips tabaci transmitting

virus disease to, 140.

Chile, miscellaneous pests in, 428, 512; beneficial insects and biological control in, 512; summary of plant quarantines in, 349; Antholcus varinervis imported into New Zealand from, against Acaena, 334.

Chillies (see Capsicum).

Chilo, new parasites of, in India,

Chilo dichromellus, on rice in Ceylon, 594.

Chilo infuscatellus, result of rearing Telenomus beneficiens on, Formosa, 92.

Chilo simplex, on Coix lachryma-jobi in Ceylon, 594; in Japan, 97; in Spain, 458; bionomics of, on rice, 97, 458; measures against, 458.

Chilo trypetes, susceptibility varieties of sugar-cane to, in Punjab, 272.

Chilo zonellus, parasites of, in India and Ceylon, 268, 610.

Chilocorus, experiments against Aspidiotus perniciosus in Russian Union, 582.

Chilocorus cacti, establishment of, against Coccids in Porto Rico,

471.

Chilocorus distigma, introduced into Seychelles against Coccids, 324.

Chilocorus kuwanae, predacious on Aonidiella aurantii in Japan, 102. Chilocorus nigritus, imported into

Mauritius from Ceylon, **610.** Chilocorus politus, utilisation of, against Aspidiotus destructor in Mauritius, 610.

Chilocorus wahlbergi, introduced into Seychelles against Coccids, 324.

Chilomenes lunata (see Cydonia).

Chilomenes sexmaculatus, predacious on Pyrilla perpusilla in Punjab, 82.

China, new Trypetid on Citrus in, **524**; Lepidopterous forest pests in, 544; miscellaneous pests in, 153, 446; utilisation of Ericerus pela in, 153; parasite of Cydia molesta in, 33; insecticidal plant in, 241.

Chinch Bug (see Blissus insularis

and B. leucopterus).

Chinch Bug, False (see Nysius ericae).

chinensis, Bruchus (Callosobruchus); Oxya; Pristomerus.

Chionaspis citri (see Prontaspis).

Chionaspis furfura, on Cotoneaster in California, 223.

Chionaspis madiunensis (see Aulacaspis).

Chirozetes scutellaris, systematic position of, 265.

chittendeni, Dialeurodes.

Chives, Micromyzus formosanus on, in Hawaii, 651.

Chloridea (see Heliothis).

Chloric Acid, tests with, against Hylemyia brassicae on cauliflowers, 110.

Chlorine, tests of, as a fumigant against Coleoptera, 454, 477.

Chloris gayana, pests of, in Queensland, 466.

Chlorochroa ligata, bionomics of, on cotton and flax in U.S.A., 499,

500.

Chlorochroa sayi (in U.S.A.), on cereals, 537; types of injury to beet and potatoes by, 429, 596; measures against overwintering populations of, 537.

Chloroform, as a solvent, 352, 397,

398, 586.

Chlorophora excelsa, Phytolyma lata

on, in Gold Coast, 492.

Chloropicrin, fumigation of date offshoots with, against Hypera brunneipennis, 629; as a soil fumigant, 199, 203; against pests of stered products, 97, 209, 429, 454; methods of fumigating sacked grain with, 209, 210; moisture content of maize affecting injury by, **429**.

Chlorops oryzae, in Japan, 152, 173; effect of ear-formation of grasses on development of, 173; mor-

phology of, 152.

Chocolate Factories, Ephestia spp. in, in Britain, 617.

Cholam (see Sorghum).

Chondracris rosea, bionomics of, in Formosa, 94.

albomarginatus, Chorthippus Russian Union, 485, 486, 570; effect of climate on outbreaks of, 485, 486; natural enemies of, 485.

Chortoicetes terminifera, in Australia, 141, 144, 466; outbreaks of, 141, 466; effect of temperature and rainfall on, 141; organisation of control of, 144.

chortoicetes, Scelio.

Chortophaga viridifasciata, experimentally parasitised by Muscina stabulans in Texas, 477.

Chortophila (see Hylemyia).

Chromaphis juglandicola, thrips predacious on, on walnut in California, 211.

chrysanthemi, Diarthronomyia.

Chrysanthemum, pests of, in Br. Isles, 63, 381, 525; legislation against Diarthronomyia chrysanthemi on, in N. Ireland, 64; Bryobia on, in Punjab, 269; pests of, in U.S.A., 207, 555; Macrosiphum sanborni reared on, for testing insecticides, 591; injured by methyl bromide, 555; introduction of selenium tissues of, 207.

Chrysanthemum Midge (see Diarth-

ronomyia chrysanthemi).

Chrysis, parasite of Niphadolepis alianta in Nyasaland, 154, 588.

Chrysobothris femorata, bionomics and control of, on Cercis canadensis in Kansas, 635.

chrysocephala, Psylliodes.

Chrysocharis, parasite of Phytomyza

spicata in Fiji, 496.

Chrysocharis gemma (parasite of Phytomyza ilicis), in Britain, 417, 490; establishment of, in Br. 417; intrinsically Columbia, superior to Opius ilicis, 490.

Chrysocharis syma (parasite Phytomyza ilicis), in Britain, 417; introduced into Br. Columbia,

417.

Chrysomela (Chrysolina) geminata, introduced into Australia against Hypericum perforatum, 143.

chrysomphali, Aphytis (Aphelinus);

Xenocomys.

Chrysomphalus, measures against, on Citrus in Transcaucasia, 653.

Chrysomphalus aonidum, auct. (see C. ficus).

Chrysomphalus dictyospermi, attacked by Comperiella bifasciata

in Spain, 457.

Chrysomphalus ficus, in Brazil, 624; in Florida, 426, 474; known as "black scale" in Palestine, 564 (note); in Seychelles, 324; in W. Indies, 445, 471, 498; on banana, 445; on Citrus, 426, 471, 474, 498, 624; on coconut, 324, 498; bionomics of, 426, 474, 625; natural enemies and biological control of, 324, 471, 625.

Chrysomphalus lahillei, new parasite

of, in Argentina, 344.

Chrysomphalus obscurus, eradication campaign against, in California,

Chrysopa (predacious on noxious insects), in Connecticut, 443; in India, 184; in Rumania, 13.

Chrysopa irregularis, predacious on Pseudococcus brevipes in Fiji, 389.

Chrysopa plorabunda, predacious on Toxoptera graminum in Oklahoma, 251.

Chrysopa ramburi, predacious on Pseudococcus brevipes in Fiji, 389. chrysopae, Brischke, Hemiteles.

chrysopae, Uch., Hemiteles (Rhadiur-

gus) (see H. kichijoi).

Chrysoplatycerus splendens, parasite of Pseudococcus maritimus in California, 310; introduced into Canada, 310.

chrysorrhoea, auct., Euproctis (see Nygmia phaeorrhoea).

Cicadula, auct. (see Macrosteles).
Cicer arietinum, as a trap crop for
Heliothis virescens in Peru, 11,
510.

Cichorium endivia (see Endive). ciliata, Campsomeris.

cilicrura, Hylemyia.

Cimex lectularius, action of pyrethrum on, 484; permeability of cuticle of, by insecticide mixtures, 239.

Cinara hyalina (see C. pinicola). Cinara laricicola, parasites of, on larch in Japan, 445.

Cinara pini, measures against, on

pines in U.S.A., 607.

Cinara pinicola, measures against, on Picea abies in U.S.A., 607.

Cinara sabinae, bionomics and control of, on Juniperus virginiana in U.S.A., 607.

Cinara strobi, bionomics and control of, on Pinus strobus in U.S.A., 607.

cincticeps, Nephotettix bipunctatus. cinctipennis, Taeniothrips usitatus. cinctus, Anthonomus (see A. piri); Cephus.

Cineraria, Brotolomia meticulosa on, in Br. Isles, **525.**

cinerea, Zeuxia.

cingulata, Rhagoletis.

Circopes pilistriatus, on Brachychiton populneus in New South Wales, 467.

circumflexus, Myzus.

Cirphis unipuncta, on grasses in Fiji, 495; on flax in Formosa, 93; on Setaria italica in Japan, 97; bionomics and control of, on cereals, etc., in U.S.A., 254, 300, 356; natural enemies of, 98, 300, 495; laboratory rearing of, for insecticide tests, 199.

Cirrospilus nigrovariegatus, parasite of Parornix prunivorella in Kan-

sas, 190.

cistipennis, Fundella.

citri, Aleuroplatus; Dialeurodes; Halticus; Mellesis; Paratetranychus; Prontaspis (Chionaspis); Pseudococcus.

citricidus, Aphis. citrina, Aonidiella.

citrinus, Aspidiotiphagus.

Citripestis sagittiferella, measures against, on Citrus in Malaya, 91. Citron, pests on, in Transcaucasia, 654.

Citronella Oil, 183, 594.

citrulli, Itonida.

Citrullus vulgaris (see Watermelon). Citrus, pests of, in S. Africa, 18, 227, 320, 341, 435; Ceratitis capitata on, in Algerian Sahara, 616; pests of, in Argentina, 263, 344, 445, 512; pests of, in Australia, 88, 144, 280, 335, 336, 437, 465, 480; pests of, in Brazil, 167, 168, 392, 624; moths damaging fruits of, in Ceylon, 594; Mellesis citri on, in China, 524; Trypetids on, in Cyprus, 544; pests of, in Fiji, 169, 170, 389, 390; pests of, in Formosa, 94, 103, 622; moths damaging fruits of, in Gold Coast, 16, 106, 107; pests of, in Guam, 331, Atta cephalotes on, in Dutch Guiana, 169; pests of, in Japan, 98, 102, 172; pests of, in Malaya, 91, 513; mite on, in Nyasaland, 589; Pseudococcus comstocki on, in Palestine, 73, 339; Icerya purchasi on, in Peru, 511; new mite on, in Punjab, 271; Icerya seychellarum on, in Seychelles, 82; pests of, in Spain, 462, 463; pests of, in Transcaucasia, 582, 653, 654; pests of, in U.S.A., 135, 209, 214, 222, 295-297, 298, 336, 349, 355, 386, 398, 399, 405, 407, 410, 426, 474, 475, 535, 553, 561, 564, 565, 597; pests of, in W. Indies, 51, 262, 293, 471, 498, 648; preferred food-plant of Pseudococcus citri, 155; factors affecting parasitism of Coccids on, 220; evaluation of results of fumigating, against Coccids, 463; insecticides and injury to, 98, 280, 296, 437, 561, 652, 653, 654; methods of determining oil deposits on foliage of, 410, 411; fruit of, in baits for fruit-piercing moths, 107. Citrus Fruit Borer (see Citripestis

Citrus Fruit Borer (see Citripestis sagittiferella).

Citrus Red Scale (see Aonidiella aurantii).

Citrus Rust Mite (see Phyllocoptruta oleivorus).

Citrus Thrips (see Scivtothrips aurantii).

civiloides, Tachina (Eutachina).

Clania crameri, on Lantana camara in India, 289.

claratibia, Bussothrips. claripalpis, Theresia.

claripennis, Exenterus; Phorocera.

Clary Sage (see Salvia sclarea).

clausa, Trichopsidea (Parasymmictus).

Clausenia purpurea (parasite of Pseudococcus comstocki), in Japan, 425; in U.S.A., 196, 425.

clavicornis, Parechthrodryinus.

clavinervis, Mesostenoideus.

Clay Coloured Weevil (see Otiorrhynchus singularis).

Clays, as carriers for dusts, 22, 46, 165, 555, 569; effect of electrostatic charges of dusts containing, 569; for emulsifying oils, 654. (See Bentonite.)

cleanthes, Rhaconotus.

cleghorni, Diploneura (see D. cornuta).

clematifoliae, Myzus (see M. varians).

Clematis, Myzus varians migrating from peach to, in California, 45.

Clensel, in baits for fruit-flies, 183, 462.

Cleome monophylla, Myzus persicae on, in Nyasaland, 154.

on, in Nyasaland, 154.
Cleonus mendicus, natural enemies

of, on beet in Italy, 316. Cleonus punctiventris (on beet in Ukraine), hibernation studies on, 577; movements of, on soil, 577, 578; value of trap ditches for, 577, 578; other measures against, 584, 586; agricultural practices affecting egg parasite of, 455, 456.

cleonyma, Opogona.

Clerodendron, Arctiid on, in Formosa, 94.

Clerodendron infortunatum, Xyleutes ceramicus on, in Burma, 76.

Cletomorpha lancigera, on cacao and Solanum in Nigeria, 516.

Clidemia hirta, distribution of Liothrips urichi against, in Pacific Islands, 495, 496; Xyleborus compactus on, in Fiji, 170.

Closterocerus splendens, parasite of Promecotheca papuana in New

Guinea Territory, 378.

Clothes Moths, penetration of cracks by larvae of, 216; baits and traps for, 255; paradichlorobenzene against, 363; toxicity of thiourea to, 408. (See *Tineola biselliella*.)

Clove, Melolonthids not important pests of, in Zanzibar, 83.

Clover (*Trifolium*), pests of, in Australia, **236**, **547**, **587**; resistant to *Lachnosterna* in Canada,

90; Tipulids on, in Germany, 68; Odontria zealandica on, in New Zealand, 541; relation of wireworms to, in Russia, 572, 573; pests of, in U.S.A., 164, 357, 530, 537; in crop rotation against Blissus leucopterus, 245.

Clover, Crimson (see Trifolium

incarnatum).

Clover, Sour (see Melilotus indica). Clover, Subterranean (see Trifolium subterraneum).

Clover, Sweet (see Melilotus).

Clover Leafhopper (see Aceratagallia sanguinolenta).

Clubiona, effect of insecticides on, hibernating on apple in Germany, 70.

Clubiona jucanda, destroying Phaenacantha marcida in Formosa, 105. clytusalis, Sylepta.

Cob Nuts (see Corylus).

Coccidaphycus nigricans, gen. et sp. n., parasite of Coccids in Argentina, 344.

coccidivorus, Anagyrus.

coccidophaga, Neocopidosoma.

Coccidoxenus niloticus, sp. n., parasite of Saissetia oleae in Kenya, 385; introduced into California, 385, 386.

Coccinella ancoralis, predacious on Toxoptera graminum in Argentina, 36.

Coccinella arcuata, predacious on Aphis maidis in Guam, 331; attempted establishment of, in Hawaii, 331.

Coccinella bipunctata (see Adalia). Coccinella metallica (see Plagiodera).

Coccinella septempunctata, predacious on Pyrilla perpusilla in Punjab, 82; predacious on other insects in Russian Union, 486, 582; hibernation of, 486.

Coccinella undecimpunctata, predacious on Pyrilla perpusilla in Punjab, 82; predacious on Aphids in Uzbekistan, 486; hibernation of, 486.

Coccoloba ilhense, Lamiid on, in Brazil, 292.

Coccomytilus (see Lepidosaphes).

Coccophagus (parasitising Saissetia oleae), African species of, and their introduction into California, 385, 386; key to, 386.

Coccophagus anthracinus, 385. Coccophagus basalis, 385. Coccophagus capensis, 385, 386. Coccophagus coracinus, 385.

Coccophagus cowperi, 385, 386. Coccophagus eleaphilus, 385. Coccophagus flavidus, sp. n., 385. Coccophagus gurneyi, hosts of, in California, 297, 355.

Coccophagus japonicus, parasite of Ceroplastes rubens in Japan, 172.

Coccophagus nigritus, 385.

Coccophagus ochraceus, 385, 386. Coccophagus pulvinariae, 385, 386.

rusti Coccophagus (parasite Saissetia oleae), 385, 386; factor effecting parasitism by, 220. Coccophagus saintebeauvei, 385.

Coccophagus spectabilis, 385. Coccophagus trifasciatus, 385, 386. Coccophagus tschirchii, parasite of

Laccifer lacca in India, 263.

coccorum, Pachyneuron.

Coccosterphus tuberculatus, suspected vector of spike disease of sandal in India, 75.

Coccotrypes dactyliperda in imported buttons in Bermuda, 139.

Coccus hesperidum (on Citrus), new parasite of, in Argentina, 445; in Fiji, 389; measures against, in Transcaucasia, 654.

Coccus pseudomagnoliarum, measures against, on Citrus in Cali-

fornia, 349.

Coccus viridis, in New Guinea Territory, 377; in W. Indies, 262, 498; on Citrus, 262, 498; on coffee, 377, 498; factors affecting attacks of, 262.

coccus, Dactylopius. cockerelli, Paratrioza.

Cockroaches, in greenhouses in Britain, 489; damaging plants, etc., in U.S.A., 562; tests of insecticides on, 352, 353, 556.

Cocksfoot Grass (see

glomerata).

Coconut, pests of, in Brazil, 9, 10, 468; Nephantis serinopa on, in Ceylon, 594, 619-622; Rhynin, chophorus palmarum in, in Colombia, 9; pests of, in Fiji, 169, 495; Aspidiotus destructor on, in Florida, 472; pests of, in Guam, 331, 332; Brassolis sophorae on, in Br. Guiana, 538; pests of, in Hawaii, 274, 331; pests of, in India, 270, 620; pests of, in Java, 79, 148; 148; Melittomma insulare on, in Madagascar, 341; pests of, in Malaya, 79, 148; A. destructor on, in Mauritius, 610; pests of, in New Caledonia, 391; pests of, in New

Guinea Territory, 91, 377, 378, 379; Brontispa mariana on, in Saipan, 331; Oryctes rhinoceros on, in Samoa, 148; pests of, in Seychelles, 82, 323, 340, 341; Amblypelta cocophaga causing nutfall of, in Solomon Is., 324, 325; pests of, in W. Indies, 263, 471, 472, 498; methods of treating seedlings of, against Strategus quadrifoveatus, 472; Ephestia cautella damaging stored nuts of, 9.

Coconut Caterpillar (see Brassolis sophorae and Nephantis serinopa). Coconut Leaf Hispid (see Brontispa

longissima).

Coconut Leaf Miner (see Promecotheca papuana).

Coconut Oil (see Oils, Vegetable). Coconut Scale (see Aspidiotus des-

tructor).
Coconut Treehoppers (see Sexava). cocophaga, Amplypelta; Graeffea. Cocos spp., pests of, in Brazil, 9,

10, 468. (See Coconut.)

Codling Moth (see Cydia pomonella). Codling Moth, False (see Argyroploce leucotreta).

Coeloides pissodis, parasite of Dendroctonus frontalis in Virginia,

Coelonotus laricis, parasite of Cinara laricicola in Japan, 445.

Coelonotus pini, parasite of Cinara laricicola in Japan, 445.

Coelophora inaequalis (predacious on Aphids), in Guam, 331; in Hawaii, 651; establishment of, in Porto Rico, 471; introduced into U.S.A., 431.

Coelophora saucia, introduced into Formosa against Oregma lanigera,

coenosulus, Nysius. coerulea, Agromyza.

coerulescens, Tetrastichus.

coffeae, Helorimorpha; Saissetia; Zeuzera.

coffearia, Homona.

Coffee, pests of, in Brazil, 292, 344; Stephanoderes hampei on, in Ceylon, 594; pests of, in Belgian Congo, 80, 514; Coccids on, in Republic, Dominican original record of Leucoptera coffeella on, in Guadeloupe, 35; Atta cephalotes on, in Dutch Guiana, 169; Pseudococcus citri on, in Java, 155; Antestia lineaticollis on, in Kenya, 80;

pests of, in Malaya, 512, 513; Coccids on, in New Guinea Territory, 377; pests of, in Nyasaland, 59, 60, 154; pests of, in Tanganyika, 175-177, 514, 589; pests of, in Uganda, 80, 175, 229, 230, 341; effects of shade and altitude on susceptibility of, to Antestia, 229, 230; Nematospora associated with Antestia on, 177. Coffee (Stored), Araecerus fascicu-

latus in, in Venezuela, 292.

Coffee Berry Borer (see Stephanoderes hampei).

Coffee Leaf-miners (see Leucoptera caffeina and L. coffeella).

coffeella, Leucoptera.

coimbatorensis, Dinarmus; Euplectrus.

Coix lachryma-jobi, pests of, in Ceylon, 594; susceptible Aplanobacter stewarti in U.S.A., 245.

Colasposoma sellatum, on Citrus in Queensland, 465.

Cold Storage, uses of, against pests, 138, 426, 544, 605; insects kept in, 84, 151, 265; rendering apples susceptible to injury by methyl bromide, 113.

Cold-hardiness, studies on, in insects, 69, 123.

colemani, Anastatus.

Coleophora, keys to Palaearctic species of, feeding on fruit trees, 25.

Coleophora anatipennella, bionomics of, in France, 26.

Coleophora badiipennella, on fruit trees in France, 26.

Coleophora currucipennella, on fruit trees in France, 26.

Coleophora fuscedinella, on fruit trees in France, 26.

Coleophora hemerobiella, on fruit trees in France, 26; tests with tar distillates against, in Germany, 231.

Coleophora malivorella, parasites of, on apple in W. Virginia, 363.

Coleophora nigricella, bionomics of, in France, 26.

Coleopterous Larvae, digestive enzymes of wood-boring species of, 340.

Coleoxestia spinipennis, on fig and orange in Brazil, 392.

Colias electo, Microbracon brevi-·cornis associated with, in S. Africa, 17.

Colias eurytheme (in U.S.A.), on ground-nuts, 599; on lucerne, 223, 600; bionomics of, 600; tests of insecticides against, 600. Colias marnoana, on lucerne in Sudan, 388.

collaris, Thyraeella (Diadromus). Colletes, parasitised by Myopa in

Europe, 474.

Colobogaster cyanitarsis, bionomics of, on Ficus spp. in Brazil, 391.

Colobogaster quadridentata, on Ficus spp. in Brazil, 391.

Colocasia (Taro), pests of, in Pacific Is., 170, 171, 273, 280, 377, 390.

Colombia, cotton pests in, 8, 53, 54; miscellaneous pests in, 8, 9; beneficial insects and biological control in, 9, 53, 54; thrips intercepted in New York on Eugenia from, 86.

colonus, Xylotrechus.

coloradensis, Sarcophaga.

Colorado, Dendroctonus ponderosae on pines in, 426; Paratrioza cockerelli in, 21, 596; miscellaneous pests in, 533, 596; beneficial insects in, 190, 294. Colorado Potato Beetle (see Lep-

tinotarsa decemlineata). columbicus, Dysdercus.

comariana, Peronea (Acalla). comitator, Brachymeria.

Commelina nudiflora, vector and identity of virus disease of, in Hawaii, 140.

communis, Melanotus.

compactus, Pyrilloxenos; Xyleborus. comperei, Euryischia.

Comperiella bifasciata, introduced into Spain from California, 457.

completa, Rhagoletis suavis. composita, Persectania.

compressus, Cylas (see C. puncticollis).

Compsilura concinnata, parasite of Pieris rapae in Britain, 314. comptana, Ancylis.

comstocki, Ephialtes (Calliephialtes); Pseudococcus

comstockiana, Rhyacionia.

Conchyloctenia parummaculata, parasite of, on sweet potato in Nyasaland, 589.

concinnata, Compsilura.

Condy's Crystals, slight toxicity of solution of, to Calandra oryzae, 541.

confusa, Diacrisia obliqua.

Confused Flour Beetle (see Tribolium confusum).

confusum, Tribolium. confusus, Dactylopius.

Congo, Belgian, Helopeltis on cotton in, 516; locusts in, 228, 388; lists of termites and timbers resistant to them in, 611; new parasites of coffee pests in, 80, 514.

congregatus, Apanteles. conicollis, Calandra.

Coniophora cerebella, favouring Xestobium rufovillosum in willow timber, 315.

conjunctus, Pristogonatopus.

Connecticut, Anasa tristis on, cucurbits in, 163, 442-444; forest pests in, 161, 163, 164, 424, 552; miscellaneous pests in, 161-164; orchard pests in, 87, 161, 425, 533; beneficial insects and biological control in, 87, 160, 424, 442, 443, 533, 552.

connexa, Eriopis.

Conocephalus saltator, on potato in Hawaii, 650.

Conoderus auritus, bionomics of, on tobacco in Kentucky, 482; cultural measures against, 483.

Conopia exitiosa (see Aegeria). conopleura, Echthromorpha. Conorrhynchus (see Cleonus).

Conotrachelus denieri, sp. n., in Argentina, 445; bionomics of,

on cotton in Paraguay, 445.
Conotrachelus nenuphar, bionomics of, on peach and apple in U.S.A., 41, 196, 197, 418; measures against, 42, 196, 198.

conquisitor, Pimpla (Ephialtes). consobrina, Phyllodes.

conspersa, Narosa. "Contact," in spray formula against Dermestids, 454.

Contarinia, periodicity in oviposition of species of, on grasses in Britain, 314 315.

Contarinia dactylidis, on Dactylis glomerata in Britain, 314, 315.

Contarinia festucae, sp. Festuca in Britain, 315. n., on

Contarinia juniperina, binomics of, on Juniperus spp. in U.S.A., 432, 433.

Contarinia lolii, on Lolium perenne in Britain, 315.

Contarinia medicaginis, bionomics and control of, on lucerne in Hungary, 12.

Contarinia merceri, on Alopecurus pratensis in Britain, 314, 315.

Contarinia sorghicola, factors affecting injury to Sorghum by, in Queensland, 466.

Contarinia tritici, periodicity in oviposition of, 315.

continentalis, Tetramorium sericiventre.

contraria, Ochrogaster.

convergens, Hippodamia; Omocena. convexa, Pulvinaria.

Convolvulus spp., insects on, in Russia, 479.

cooleyi, Chermes (Adelges).

Copidosoma, parasite of Autographa spp. in Quebec, 626.

Copper Ammonium Silicate (Coposil), effect of, on lead-arsenate spray residues on apple, 412.

Copper Arsenate, tests of toxicity of, to Colias eurytheme, 600; in baits for locusts, 612, 613; (basic), experiments with, in dusts and sprays, 39, 119, 561, 632, 647; temperature affecting toxicity of, to insects, 356.

Copper Carbonate, mixed with cowpeas against Bruchus chinensis,

Copper Oxychloride, addition of, to derris dust against bacterial wilt of cucurbits, 647; cryolite compatible with, 599.

Copper Pitchate, action of dusts containing, on insects, 247.

Copper Sulphate, of little value against Calandra oryzae, 540, 541; protecting fruit trees from fungus-growing termites, against snails on watercress, 111 toxicity of, to Empoasca fabae, 440, 441; in mixed dusts, 250, 441,

504. (See Bordeaux Mixture.) Copper Sulphate, Basic, cryolite compatible with, 599; (as an emulsifier), decreasing injury to apple by oils, 585.

Copra, Ephestia cautella damaging, 9, 377.

Coptopterus decoratus (see Strongylurus).

Coptotermes acinaciformis, laboratory rearing of, in Australia, 142; damaging building timbers in New Zealand, 333.

Coptotermes havilandi, in Barbados, 6; in Mauritius, 6; possibly in

Br. Guiana, 6.

Coptotermes testaceus, imported into Barbados in wood from Br. Guiana, 6.

coracinus, Coccophagus.

corallipes, Xanthippus.

corchori, Apion.

Corchorus acutangulus, Pempherulus affinis on, in Madras, 521.

Corchorus capsularis, pests of, in Bengal, 446.

Corchorus olitorius, pests of, in India, 446, 521, 522.

Corcyra cephalonica, in stored cacao in Gold Coast and Venezuela, 616; probably migrating from stored rice, 616.

Cordyceps, infesting insects in Br. Honduras, 152.

Cordyceps ravenelii, infesting Lachnosterna in Canada, 90.

coriaceus, Eriococcus.

Coriander (Coriandrum sativum), beetles imported into Fiji in seed of, 391, 496; Bemisia tabaci on, in India, 523; factors affecting infestation of, by Systole coriandri in Russia, 576; attractiveness of oil of, for Oscinella frit, 573.

. coriandri, Systole.

Corioxenos antestiae, parasite of Antestia in Tanganyika, 177, 590. Cork Oak (see Quercus ilex var. suber).

Corn Borer, European (see *Pyrausta nubilalis*).

Corn Earworm (see Heliothis armigera).

Corn Leaf Aphis (see Aphis maidis). Corn Rootworm, Southern (see Diabrotica duodecimpunctata).

Corn Seed Beetle (see Agonoderus lecontei).

Corn Silk Beetle (see Monolepta varicornis).

corni, Lecanium (Eulecanium).

Cornus, Cecidomyiid causing club gall of, in U.S.A., 606.

cornuta, Diploneura.

Corrosive Sublimate (see Mercuric Chloride).

corrupta, Epilachna (see E. varivestis).

corvinus, Zarhopalus.

Corylus (Filbert, Hazel), Curculio nucum on, in Britain and Switzerland, 282, 376; pests of, in U.S.A., 223, 405; basic copper arsenate causing injury to, 119.

Corymbites aeneus, observations on, in Russia, 572.

Corynbites inflatus, predacious on Gilpinia polytoma in Maine, 637. Corythuca gossypii, food-plants of, in Porto Rico, 51.

Cosmolyce boetica, Maruca testulalis confused with, on beans in Oahu, 35.

Cosmophila sabulifera (see Anomis).
Cosmopolites sordidus, on banana in New Caledonia, 391; predacious enemies introduced into Jamaica and Queensland against, 445, 466.

Cosmopteryx dulcivora, bionomics of, on sugar-cane in Java, 524. costalis, Encoptolophus sordidus;

Eutermes (Nasutitermes).

costimacula, Eublemma.

costirostris, auct., Listroderes (see L. obliquus).

Cotinis nitida, in Kentucky, 299. Cotoneaster, Coccid on, in California,

223. Cotton, factors affecting infestation of, by Heliothis armigera in S. Africa, 81; Helopeltis on, in W. Africa, 515, 516; pests of, in Argentina, 498, 512; pests of, in Brazil, 10, 11, 404, 623; pests of, in China, 446; pests of, in Colombia, 8, 53, 54; Platyedra gossypiella on, in Ecuador, 539; pests of, in Egypt, 30, 338; pests of, in Fiji, 147, 170, 496; leavest attacking in Ferrman 94. locust attacking, in Formosa, 94; pests of, in India, 183, 265, 266, 268, 515, 516, 517, 519, 521, 522, 523; Zeuzera coffeae on, in Malaya, 513; pests of, in Nyasaland, 60, 62, 153, 588; new weevil on, in Paraguay, 445: pests of, in Peru, 11, 40, 404, 509-511, 539; pests of, in Philippines, 648, 649; pests of, in Queensland, 149, 466; pests of, in Russian Union, 235, 486, 582, 584; pests of, in Sudan, 386, 387; Argyroploce on, in Tanganyika, 227; pests of, in Uganda, 175, 228, 230, 231; pests of, in U.S.A., 2, 3, 192, 198, 213, 252, 253, 257, 258, 258 198, 213, 252, 253, 257, 258, 358, 359, 404, 428, 430, 441, 500, 599, 600, 602, 603; pests of, in W. Indies, 11, 498; susceptibility of varieties of, to pests, 148, 230, 231, 266, 466, 517; Capsids associated with "crazy top" in, 387; Dysdercus spp. associated with boll rots in, 53; list of species of Empoasca attacking, 516; doubtful food-plant of Hymenia recurvalis, 46; unsuitable food-plant for Melanoplus differentialis, 3; Pedinus femoralis

possibly poisoned by, 571; in crop rotation against Euetheola rugiceps, 3; not infected with mosaic of Musa textilis, 317; insecticides causing injury to, 260, 358, 466.

Cotton Aphis (see Aphis gossypii). Cotton Boll Weevil (see Anthonomus

grandis).

Cotton Bollworm (see Heliothis armigera).

Cotton Bollworm, Pink (see Platvedra gossypiella).

Cotton Fleahopper (see

seriatus). Cotton Leafworm (see Alabama

argillacea). Cotton Stem Weevil (see Pempheru-

lus affinis).

Cottonseed, treatment of, against *Platyedra gossypiella* in Colombia, 53; products of, in baits for insects, 158, 300, 430, 534, 571.

Couch Grass (see Agropyrum repens and Cynodon dactylon).

Cowpea Curculio (see Chalcodermus

aeneus).

Cowpeas (Vigna), Bruchids on, in S. Africa, 19, 58, 59, 321; Nezara viridula on, in Fiji, 170; pests of, in India, 265, 266, 523; Pyralids on, in Porto Rico, 345: pests of, in U.S.A., 214, 252, 253, 254, 405, 503, 598, 599, 602; time of planting and varieties of, in relation to Chalcodermus aeneus, 598; insects and mosaic of, 502, 503.

Cowpeas (Stored), Bruchids in, and their control, 19, 58, 59, 148, 321,

cowperi, Coccophagus.

Crambus, on tobacco in Kentucky, 482; observations on, in Kansas, 505; measures against, damaging lawns, 408.

Crambus bonifatellus, bionomics of, in California, 408.

Crambus sperryellus, bionomics of, in California, 408.

vulvivagellus, natural Crambus enemies of, in Kansas, 505, 506. crameri, Clania (Cryptothelea).

Cranberry, pests of, in U.S.A., 45, 640, 642

Cranberry Fireworm (see Rhopobota naevana).

Cranberry Fruit Worm (see Mineola vaccinii).

Cranberry Rootworm (see Rhabdopterus picipes).

Craniotectus, on Derris in Sumatra, 319.

crassicornis, Meraporus.

crataegi, Anuraphis.

Crataegus, Typhlocyba froggatti on, in Britain and Tasmania, 284, 291; pests of, in U.S.A., 45, 242, 243; forms of Rhagoletis pomonella on, 226.

Creatonotus transiens, on Lantana

camara in India, 289.

Crematogaster brunnea matsumurai. destroying Apanteles liparidis in Japan, 172.

Crematogaster curvispinosa, destroying Stephanoderes hampei

Brazil, 292.

Cremnops (see Bracon).

crenulata, Lachnosterna (Phyllophaga).

Creolin, emulsion of, against Solenopsis moelleri, 168.

Creontiades pallidifer, food-plants of, in India, 523.

Creontiades pallidus, apparently causing "crazy top" on cotton in Sudan, 387.

Creosote, treatments of timber and logs with, against termites and Coleoptera, 7, 147, 312, 467; tests of emulsions of, against Calandra oryzae, 540, 541; and nicotine, in dust against Nysius vinitor, 31.

Cresylic Acid, and oil emulsions, in sprays against Aphids and Spilonota, 128, 194.

cretica, Sesamia.

cribricollis, Apogonia.

crinita, Lachnosterna (Phyllophaga). Crioceris asparagi, on asparagus in U.S.A., 48, 430, 642; bionomics of, 642; Meigenia mutabilis introduced from France against,

430.

duodecimpunctata, Crioceris asparagus in Minnesota, 48.

critolaus, Spathius.

crocale, Catopsilia.

Crossocosmia sericariae (see Sturmia).

Crossotarsus saundersi, on avocado and grapefruit in Fiji, 170, 639; measures against, 389.

Crotalaria, susceptibility of species of, to Pyralids in Porto Rico, 345. Crotalaria juncea, Creontiades pallid-

ifer on, in India, 523.

Crotalaria spectabilis, not attacked by wireworms in S. Carolina, 253.

Croton, relation of Psallus seriatus to, in Texas, 198.

P. crotonis, Pseudococcus (see lilacinus).

Crows, destroying Polyphylla perversa, 509.

cruciata, Austroicetes.

cruciferarum, Plutella (see P. maculipennis).

crucigerus, Dociostaurus.

Cryolite, in baits, 429, 534, 598, 631; in bait-sprays, 55; dusting with, 145, 247, 257, 345, 358, 428, 429, 430, 476, 510, 526, 561, 562, 598, 599, 600, 601, 631, 640, 641; carriers for, 145, 430, 510, 598, 631, 640, 641; conditioning agent for dust of, 358; aeroplane dusting with, 18; spraying with, 22, 55, 67, 134, 190, 197, 243, 330, 412, 534, 535, 559, 561, 562, 598, 599, 633, 644; tests of toxicity of, to insects, 55, 356, 481, 510, 550, 600; temperature affecting of, toxicity 356; particle size affecting toxicity of, to bees, 550; types of, 260, 356, 534, 561, 599, 633; formulae containing, 55, 134, 145, 197, 330, 429, 510, 534, 535, 633, 640, 641; and copper compounds, 599; and cubé, 562; and derris, 562; and oil emulsions, 134, 633; and Selocide, **559**: soaps as spreaders for, 330; and sucrose, 561, 562; and sulphur, 430, 510, 599; and tartar emetic, 562; and zinc sulphate, 134; and injury to plants, 197, 260, 358, 428, 599; other insecticides compared with, 55, 67, 197, 247, 257, 429, 430, 561, 599, 631, 640, 641.

Cryphalus jatropha, on wilkesiana in Fiji, 170. Acalypha

Cryptaspidiotus shastae, bionomics and control of, on Juniperus virginiana in Kansas, 632.

Cryptochetum iceryae, established against *Icerya purchasi* in Chile, 512.

Cryptognatha flaviceps, introduced into St. Lucia against Aspidiotus destructor, 263.

Cryptognatha nodiceps (predacious on Aspidiotus destructor), distribution and utilisation of, in W. Indies, 263, 471, 498; introduced into Florida and Hawaii, 471, 472.

Cryptognatha simillima, introduced into St. Lucia against Aspidiotus destructor, 263.

Cryptogonus orbiculus var. nigripennis, Telsimia nitida recorded as, in Guam and Hawaii, 331.

Cryptoideus luctuosus, parasite of Saperda candida in New York, 243.

Cryptolaemus montrouzieri, predacious on Dactylopius spp. in S. Africa, 17; utilisation of, against Pseudococcus spp.: in Portugal, 63; in U.S.A., 297, 425; in W. Indies, 52, 498.

Cryptophasa unipunctata, boring in trees in Australia, 335.

Cryptophlebia illepida (carpophaga) (see Argyroploce).

Cryptorrhynchus mangiferae, intercepted in mango seeds in Fiji, 148.

Cryptostemma calendulaceum, favouring Halotydeus destructor in W. Australia, 141.

Cryptotermes (see Calotermes).

Cryptothelea crameri (see Clania).

Cryptus, parasite of Dendrolimus in Sakhalin, 95.

Ctenolepisma longicaudata (Silverfish), bionomics of, in Australia, 55-57; measures against, 56.

Cuba, parasite of Laphygma eridania in, 538.

Cubé, dusting with, 118, 162, 202, 405, 408, 504, 562, 596, 601, 630, 631; effect of pH of carrier on toxicity of, 301; carriers for, 504, 630; conditioning agents for dusts of, 504; spraying with, 120, 196, 202, 426, 597, 606; tests of toxicity of, to insects, 118, 408, 562; formulae containing, 120, 196, 597, 606; and oils, 426; and resin residue emulsion, 606; rotenone contents of, 120, 162, 408, 504, 562, 596, 597, 601, 630, 631; other insecticides compared with, 120, 196, 202, 405, 426.

cucullata, Rondania.

Cucumber, Mycetophilids on, in greenhouses in Britain, 145; pests of, in Hawaii, 35, 273; Dacus ferrugineus on, in Malaya, 90; pests of, in U.S.A., 122, 162, 214, 599, 647; fumigation of fruits of, against introduction of U.S.A. Trypetids into Hawaii, 499; vectors, host range and transmission of virus diseases of, 57, 179, 438, 501; cowpea mosaic not transmitted to, 503;

insecticides causing injury to, 122, 214, 599.

Cucumber Beetle, Spotted Diabrotica duodecimpunctata).

Cucumber Beetle, Striped (see Diabrotica melanocephala).

cucumeris, Epitrix.

cucurbitae, Dacus (Chaetodacus); Manilabaris; Tetranychus.

Cultural Practices, influence of, on insect pests, 112, 204, 253, 254, 375, 393, 547, 643; effect of, on parasite of Cleonus punctiventris, 455, 456.

cunea, Hyphantria.

Cuprous Cyanide, uses of, against insect pests, 117, 561, 614.

Curculio nucum (on Corylus), in Britain, 282; bionomics of, in Switzerland, 376.

Curculio, Cowpea (see Chalcodermus aeneus).

Curculio, Plum (see Conotrachelus nenuphar).

curculionis, Bathyplectes; Eupelmus; Triaspis.

Curinus (predacious on Asterolecanium spp.), occurrence and utilisation of, in West Indies, 471. Currant, spray schedule for, in

Kentucky, 2.

Currant, Black, Eriophyes ribis on, in Britain, 281; lime-sulphur causing injury to, 281. currucipennella, Coleophora.

curta, Eurytoma.

curvipennis, Dacus (Strumeta).

curvispinosa, Crematogaster. custator, Thyanta.

custodiens, Anoplolepis. custos, Auriga (Arma).

Cutworms, baits for, 31, 100, 138, 170, 300, 390, 576, 615, 648; other measures against, 39, 300, 428, 450, 547, 576, 615; laboratory rearing of, 199.

cyanea, Haltica; Scutellista.

cyanella, Lema.

cyanimus, Torymus.

cyanitarsis, Colobogaster.

Cyanogas, 19, 336, 338, 377, 610.

Cycas, Aonidiella aurantii on, in Japan, 102; affecting parasitism of A. aurantii by Habrolepis rouxi, 220.

Cyclamen, Tortricid on, in greenhouses in Britain, 146.

Cyclocephala, in Arkansas, 3. Cyclocephala borealis, in Kentucky,

299.

Cyclocephala immaculata, in U.S.A., 3, 261, 299; longevity, and eggstage of, 3; winter rearing of, for soil insecticide tests, 261.

Cycloneda sanguinea, predacious on Aphids in S. America, 36, 54. cycloporus, Chelonus (Chelonella).

Cydia funebrana, experiments with Trichogramma spp. against, on plum in Russia, 581.

Cydia glycinivorella, parasites of, on soy beans in Far East, 33, 34, 98. Cydia latiferreana, bionomics of, in

U.S.A., 86, 405.

molesta (Oriental Cydia 549, 602, 632; on pip fruits, 48, 141, 372; parasites and biological control of, 33, 87, 129, 130, 141, 161, 169, 192, 193, 220, 294, 372, 373, 425, 549, 602; banding against, 373; spraying ineffective against, 141.

Cydia nigricana (Pea Moth), in Britain, 310; in Canada, 22, 310; parasites and biological control of, 310, 311; regulated area

against, 22.

Cydia packardi, on cherry in Wash-639, ington, 641; measures

against, 641.

Cydia pomonella (Codling Moth), in S. Africa, **415**, **435**; in Argentina, 497; in Australia, 151, 216, 238, 279, 291, 465; in Britain, 281; in Canada, 22, 51, 224, 330; in France, 66; in Germany, 71; in Jugoslavia, 72; in Peru, 511, 539; in Russian Union, 580, 581, 582, 585; in Spain, 457; in Switzerland, 66; in U.S.A., 1, 2, 45, 48, 71, 113, 127, 128, 129, 160, 190, 192, 193, 206, 220, 222, 351, 359, 361, 371, 406, 411, 414, 415, 535, 537, 549, 550, 559, 632, 415, 535, 537, 549, 550, 559, 632, 633, 635, 639, 640, 641, 643, 645; on pear, 66, 72, 113, 222, 279; spur-burrowing habit of, on pear, 359; on stone fruits, 48, 415, 416; apparent development of race of, attacking walnut, 549; studies on adult activity of, 66, 72, 193, 238, 425, 435, 580; effects of temperature and rainfall on, 1, 2, 131, 330; other bionomics of, 51, 66, 71, 72, 216, 224, 238; diapause in, covering

two winters, 559; parasites and biological control of, 66, 192, 193, 220, 457, 511, 581, 582; fungus infesting, 535; baits for, 414, 435: measures against, 1, 2, 22, 45, 67, 71, 72, 127, 128, 160, 190, 206, 238, 279, 281, 291, 330, 351, 371, 406, 411, 416, 465, 550, 580, 585, 632, 633, 639, 640, 641, 643; attractants for use in sprays against, 129; road dust decreasing efficiency of sprays against, 361; fumigation of packed fruit against, 113, 222; tests of toxicity of insecticides to, 206, 207, 242, 414, 645.

Cydia pomonella var. simpsoni, in Australia, 144, 216.

Cydia strobilella, on spruce in Latvia, 86.

Cydonia lunata, predacious on Myzus persicae in S. Rhodesia, 184.

Cylas compressus (see C. puncti-

Cylas formicarius (Sweet Potato Weevil), fungus associated with, in Louchoo Islands, 97; in Louisiana, 413, 601, 602; bionomics of, 601, 602; fumigation of seed sweet potatoes against,

413.

Cylas puncticollis, on sweet potato in Nyasaland, 154.

Cylindrocopturus deleoni, sp. n., on pines in U.S.A., 371.

Cylindrocopturus eatoni, sp. n., on pines in California, 371.

Cylindrocopturus furnissi, sp. n., on Pseudotsuga taxifolia in U.S.A., 370.

Cylindrocopturus longulus, bionomics of, on Pseudotsuga taxifolia in Washington, 427.

Cynaeus angustus, in stored cereals in U.S.A., 506.

Cynodon dactylon (Bermuda Grass), Pantomorus taeniatulus associated with, in Australia, 608; insects associated with in U.S.A., 2, 418, 598, 629.

Cynosurus cristatus, little attacked by Porina in New Zealand, 89.

Cyperus, Toxoptera piricola possibly migrating from pear to, in Japan, 96.

Cyperus arenarius, preferred by Schistocerca gregaria in India, 519.

Cyprus, miscellaneous pests in, **544**; new parasite of *Gnorimoschema* operculella in, **514**; establish-

ment of *Rodolia cardinalis* in, 544; Bruchids from, intercepted in Fiji, 148.

Cyrtogaster vulgaris (parasite of Phytomyza ilicis), in Britain, 417; introduced into Br. Columbia, 417.

Cyrtomorpha flaviscutellaris, eggparasite of Austroicetes cruciata in W. Australia, 169.

Cyrtopeltis varians (see Engytatus geniculatus).

Cyrtorhinus fulvus, predacious on Megamelus proserpina in Fiji, **170.** cytherea, Nudaurelia.

Čzechoslovakia, parasites and forms of Gilpinia polytoma in 294, 474. czizeki, Tipula.

D.

dactylidis, Contarinia; Dasyneura; Sitodiplosis.

dactyliperda, Coccotrypes.

Dactylis glomerata, Cecidomyiids on, in Britain, 314, 315; little attacked by Porina in New Zealand, 89.

dactylopii, Leptomastix.

Dactylopius spp., utilisation of, against Opuntia spp. in S. Africa, 17; Cryptolaemus montrouzieri predacious on, 17.

Dactylosternum hydrophiloides (predacious on Cosmopolites sordidus), establishment of, in Jamaica, 445; introduced into Queensland, 466.

Dacus, liberation of parasites against, in Fiji and Ceylon, 495, 496, 594; baits showing seasonal prevalence of, 594; measures against, 91.

Dacus caudatus, host-fruits of, in Malaya, 90.

Dacus cucurbitae, in Formosa, 99; in Hawaii, 273, 499; attempted introduction of Opius fletcheri into Loochoo Islands against, 99; in Malaya, 90; legislation against importation of, into U.S.A., 499; host fruits of, 90, 273; effect of low temperatures on pupae of, 104; baits for, 90, 99; fumigation of fruit, etc., against, 499.

Dacus curvipennis, scarce in Fiji, 495; not a pest of bananas, 495. Dacus ferrugineus, in Ceylon, 594; colour forms and varieties of, in India, 183, 524; in Malaya, 90;

host-fruits of, 90, 183, 594; baits for, 90, 91, 183.

Dacus ferrugineus dorsalis, in Formosa, 94, 104; on Solanum auriculatum in India, 524; effects of temperature and humidity on, 104; baits for, 94.

Dacus ferrugineus incisus, Solanum auriculatum in India,

Dacus ferrugineus tryoni (in Australia), bionomics of, 465; tests of bait-sprays and baits for, 54, 55, 337, 465, 494; effect of sublethal doses of poison on, 494, 495; other Diptera confused with, 237.

Dacus oleae, baits for, on olive in Spain, 462.

Dacus passiflorae (in Fiji), establishment of Syntomosphyrum indicum against, 170; Phaenocarpa leveri erroneously recorded as a parasite of, 496.

Dacus psidii, not occurring in Fiji, 495.

Dacus umbrosus, on Artocarpus integrifolia in Malaya, 90; baits for, 90.

dahlemica, Phlugiola.

Dahlia, Tarsonemus latus on, in S. Africa, 320; Thrips tabaci transmitting spotted wilt to, in France, 25.

Daikon, Murgantia histrionica on, in Hawaii, 651.

daimio, Scolytoplatypus.

Daintex, composition and use of, as wetting agent, 595.

Dakota, North, grasshoppers in, 254, 394.

Dakota, South, grasshoppers in, 394, 603; Ips spp. on Pinus ponderosa in, 49.

Dalbergia sissoo, pests of, in India, 74, 514, 523.

dalmanni, Habrolepis.

Danaus limniace, parasite of, on sugar-cane in India, 183.

daricella, Leucoptera.

Dasychira achatina, 119.

Dasychira cerigoides, on Dipterocarpus tuberculatus in Burma, 78. Dasychira mendosa, on Aleurites

spp. in Burma, 79.

Dasyneura dactylidis, oviposition peak of, on Dactylis glomerata in Britain, 315.

Dasyneura leguminicola, on clover in Montana, 537.

Dasyneura mangiferae, habits of, on mango in India, 270.

Dasyscapus parvipennis, imported into Florida from Porto Rico against thrips, 472.

Datana integerrima, bionomics and control of, in Missouri, 166; tests of basic copper arsenate against, 119.

datanae, Heteropelma.

Date Palm, Dinapate filifera on, in California, 137; mites on, in Sudan, 388; fumigation of offshoots of, against Hypera brunneipennis, 629.

Datura, Lema nigrovittata on, in Hawaii, 35.

Datura fastuosa, virus disease of, in Madras. 542.

Datura stramonium, Epitrix parvula on, in Virginia, 200; Thrips tabaci transmitting virus disease to, 140.

Datura stramonium var. tatula, Tarsonemus latus on, in S. Africa,

Death Watch Beetle, application of name, 543. (See Xestobium rufovillosum).

debilis, Micromalthus; Typhlocyba. Decadarchis flavistriata, plants attacked by, in Hawaii, 273, 274; parasite of, 274.

decemlineata, Leptinotarsa (Doryphora).

decempunctata Oides.

declivis, Scobicia.

decoratus, Strongylurus (Coptopterus) Decticus albifrons, in Spain, 529. defecta, Epilachna.

defoliaria, Evannis.

Deguelin, determination of, derris, 30; Tephrosia virginiana containing, 397.

dejeani, Scolia.

delagoa, Anomala.

Delaware, Pyrausta nubilalis on maize and potatoes in, 48, 508; Plagiodera versicolora on willow in, 438; fungus infesting Cydia pomonella in, 535.

deleoni, Cylindrocopturus.

Delia (see Hylemyia). deliva, Spilochalcis.

Delomerista diprionis (parasite of sawflies), in Canada, 552; in U.S.A., 552, 637, 638; utilisation of, 552.

Delonix regia (see Poinciana).

Delphacodes striatellus, relation of, to virus diseases of cereals in Russian Union, 13, 14, 16; bionomics and control of, 14.

Delphastus (predacious on Asterolecanium spp.), introduced into Porto Rico from Trinidad, 471.

Delphastus flaviceps (see Cryptognatha).

Delphax striatella (see Delphacodes). Delphinium, pests of, in Britain, 448.

deltae, Lecanium.

Deltocephalus dorsalis, on rice in Japan, 101.

Deltocephalus striatus (see Laevicephalus).

delusorius, Syrrhizus.

Dendrocalamus (see Bamboo).

dendroctoni, Cecidostiba.

Dendroctonus, fungi associated with, on Pinus ponderosa in California,

Dendroctonus brevicomis, on pines in U.S.A., 49, 422.

Dendroctorus frontalis, on pines in U.S.A., 49, 116, 211; natural enemies of, 116, 211, 212.

Dendroctorus micans, on spruce in Sakhalin, 96.

Dendroctonus monticolae, on pines in U.S.A., 49, 647.

Dendroctonus piceaperda, on spruce in Canada, 37, 532.

Dendroctonus ponderosae, on pines in U.S.A., 49, 426; effects of low temperatures on, 426, 427.

Dendroctonus pseudotsugae, on Pseudotsuga taxifolia in U.S.A.,

Dendrol, 563.

dendrolimi, Rhogas; Trichogramma. Dendrolimus albolineatus (see D. superans).

Dendrolimus pini, unimportant on pine in Latvia, 86.

Dendrolimus punctatus, bionomics of, on pine in China, 544, 545. Dendrolimus spectabilis, hyper-

parasites of, in Japan, 172. Dendrolimus superans (albolineatus), parasites of, on pine in Sakhalin, 95.

dendrolimusi, Telenomus.

Dendrosoter sulcatus, parasite of Dendroctonus frontalis in Virginia, 211.

denieri, Conotrachelus.

dentata, Acemya.

dentatus, Acanthocoris.

denticulata, Chaetocnema.

Deodar (see Cedrus deodara).

deodarae, Ectropis.

deorum, Ageneotettix.

depressella, Emmalocera.

Derelomus bondari, sp. n., on Cocos coronata and coconut in Brazil, 10, 468.

Dermaleipa, damaging Citrus fruits in Gold Coast, 106.

Dermatitis, caused by pyrethrum,

Dermestes, in timber in Germany,

lardarius, measures Dermestes against, boring in timber in Quebec, 392, 393; Typhlodromus timber in tineivorus attacking, 376.

Dermolepida, n. n., for Lepidoderma,

388.

Dermolepida albohirtum, increase and control of, on sugar-cane in Queensland, 388; susceptibility of varieties of sugar-cane to, 389; Bufo marinus ineffective against, 389.

derogata, Sylepta.

Derostenus (parasite of Promecotheca papuana), bionomics of, in New

Guinea Territory, 378.

Derris, against Aphids, 29, 67, 96, 136, 202, 250, 287, 318, 353, 492, 504, 544, 596, 597, 631; against Coccids, 349; against Jassids, 19, 325, 326, 647; against other Rhynchota, 19, 21, 105, 121, 171, 319, 373, 465, 615; against caddis fly on watercress, 111; against Coleoptera, 29, 120, 163, 354, 361, 408, 540, 600, 601, 630, 647; against Diptera, 99, 161, 165, 527, 562; in mixture for maize ears treating Euxesta stigmatias, 470; against Hoplocampa spp., 285, 286; against Lepidoptera, 46, 129, 135, 136, 145, 283, 290, 318, 373, 404, 405, 408, 600, 641, 647; against mites, 109, 237, 281, 285, 286, 287, 546, 637; against thrips, 204, 413, 492; tests of toxicity of, to insects, 19, 29, 99, 121, 165, 204, 233, 408, 540, 562, 600; duration of activity of, 233; review of effects of, on insects, 88; toxicity to insects of smoke from, 127; dusting with, 19, 21, 46, 67, 121, 135, 136, 163, 165, 202, 233, 237, 250, 283, 290, 318, 325, 326, 353, 373, 404, 405, 408, 504, 544, 546, 555, 562, 596, 600, 204, 620, 624, 647; carriers for 601, 630, 631, 647; carriers for, 135, 136, 162, 165, 250, 318, 325, 326, 353, 404, 405, 504, 555, 630;

pH of carrier affecting toxicity of, 301; conditioning agents for dusts containing, 161, 250, 353, 504; spraying with, 29, 96, 99, 105, 109, 120, 145, 171, 202, 204, 233, 281, 285, 286, 287, 318, 319, 349, 350, 361, 373, 408, 413, 419, 420, 465, 492, 504, 597, 637, 641; use of aircraft for applying sprays containing, 256; treatment of turf with, against pests, 408, 527, 615; formulae containing, 120, 136, 145, 237, 286, 319, 325, 349, 360, 420, 527, 546, 597, 637; and copper oxychloride, 647; and lime-sulphur, 285; and nicotine, 135, 136, 237, 287, 546; and oil emulsions, 285, 286, 287, 349, 350; intermediary solvents for resins of, when used with oils, 349, 350; and pyrethrum, 233, 361, 527; and soap, 105, 145, 171, 281, 318, 319; other spreaders and adhesives for, 145, 349, 350, 361, 419, 420, 637; and sulphur, 405, 647; residues of, on cabbage, 555; stabilised form of, 120; insecticides containing, rotenone contents of, 21, 29, 120, 121, 162, 163, 165, 250, 325, 349, 350, 353, 354, 361, 404, 405, 408, 414, 492, 504, 527, 555, 562, 596, 597, 600, 601, 630, 631, 637, 647; other constituents and evaluation of, 29, 30, 135; production and marketing of, 88; other insecticides compared with, 19, 21, 29, 120, 121, 165, 202, 204, 233, 250, 318, 325, 326, 405, 414, 544.

possible Derris, food-plant Niphadolepis alianta in Nyasaland, 59; Craniotectus on, in Sumatra, 319; insecticidal properties and constituents of species of, 29, 30, 483; effect of defoliation on constituents of, 470.

desarmata, Ladoria. deserticola, Geotrogus.

desertor, Bracon (Cremnops).

Desmanthus virgatus, Bruchid intercepted in seeds of, in Fiji, 148.

Desmidophorus hebes, on Hibiscus

in India, 263.

destructor, Aspidiotus; Halotydeus; Lepidoglyphus; Mayetiola (Phytophaga); Merisus; Phyllocoptes; Scolytus (see S. scolytus).

detorquens, Technomyrmex. detrimentosus, Pleurotropis. devastans, Empoasca.

devastator, Melanoplus.

Devil's Shoestring (see Tephrosia virginiana).

dewitzi, Erencyrtus.

Dexia rustica, imported into Mauritius from England against Lachnosterna smithi, 609.

Dexiomorpha picta (parasite exiomorpha picta (parasite of Lamellicorn larvae), bionomics of, in Morocco, 83; importation of, into Mauritius against Lachnosterna smithi, 84, 609.

Dextrin, and tartar emetic, in spray against mites, 215.

Dextrose (see Sugar).

Diabrotica, not transmitting virus disease of tobacco and tomato in Argentina, 608.

Diabrotica duodecimpunctata U.S.A.), bionomics of, on maize, 599; measures against, on china asters, 361; aerial distribution of, 262.

Diabrotica melanocephala U.S.A.), on china asters, 361; on squash, 163; relation of, to bacterial wilt of cucurbits, 647; not transmitting cowpea mosaic, 503; measures against, 163, 361, 647.

Diabrotica soror, relation of, to virus diseases of squash in California, 501.

Diabrotica trivittata, relation of, to virus diseases of squash in California, **501.**

Diabrotica vittata (see D. melanocephala).

Diacrisia obliqua, bionomics and control of, on jute in Bengal, 447. Diacrisia obliqua confusa, on Lantana camara in India, 289.

Diacrisia rhodophila rhodophilodes, bionomics of, in Formosa, 93, 94. Diacrisia virginica, laboratory rearing of, 557.

Diadromus collaris (see Thyracella). Dialeges pauper, not infesting barked logs in Burma, 78.

Dialeurodes chittendeni, measures against, on rhododendron in U.S.A., 606.

Dialeurodes citri, in U.S.A., 221, 555, 597; bionomics and control of, on Citrus, 597; fumigation of ornamental plants against, 555.

Diamond Back Moth (see Plutella maculipennis).

Diaphania hyalinata, measures against, on cucurbits in Alabama,

405: laboratory rearing of, 557; insecticide tested on, 242.

Diaphania nitidalis (on cucurbits), in Peru, 539; measures against, in U.S.A., 405, 647.

Diaprepes, parasites and control of, on Citrus in St. Lucia, 262.

Diaprepes abbreviatus, on sugar-cane in Barbados, 6, 7; Bufo marinus destroying, 7; termite damage attributed to, 6.

Diapus, not infesting barked logs in Burma, 78.

Diarthronomyia chrysanthemi, distribution of, on chrysanthemum in Br. Isles, 63, 381; legislation against, in N. Ireland, 64; measures against, 63, 64.

Diaspis bromeliae, on pineapple in Fiji, 390.

Diaspis leperii (see Epidiaspis).

Diatomaceous Earth, mixed with stored wheat against Calandra oryzae, 541; as a carrier for dusts, 134, 249, 326, 630, 640, to Eutettix repellent tenellus, 134.

Diatraea, scarce on sugar-cane in Ceylon, 609; on maize and Sorghum in China, 446; parasites introduced against, on sugarcane in Mauritius, 609, 610.

Diatraea auricilia, on sugar-cane in India, 184.

Diatraea canella, on sugar-cane in Br. Guiana, 538; not parasitised by Metagonistylum minense, 538.

Diatraea saccharalis (on sugar-cane), in Brazil, 4, 5, 470; in Colombia, 9; in Br. Guiana, 538; in Peru, 511; in U.S.A., 160, 354, 431, 471, 476, 601; in W. Indies, 3-5, 160, 263, 293, 470, 471, 601; susceptibility of varieties sugar-cane to, 601; on maize, 5, 354, 511; on rice and wheat, 511; parasites and biological control of, 3-5, 9, 160, 263, 354, 431, 470, 471, 476, 538, 601; ecological and climatic factors affecting prevalence and parasites of, 160; dusting with cryolite against, 476.

Diatraea sticticraspis, parasites and biological control of, on sugarcane in India, 183, 184, 268.

Diatraea venosata (on sugar-cane), in Formosa and Java, 92; in India, 137, 184, 268; parasites and biological control of, 92, 137, 184, 268.

diatraeae, Lixophaga; Paratheresia (sée Theresia claripalpis).

Diatraeophaga striatalis, introduced into Formosa from Java against Diatraea venosata, 92.

Dibrachys affinis, enabling Melittobia acasta to parasitise Cydia pomonella, 66.

Dibutyl Phthalate, intermediary solvent for derris resins used with oils, 350.

Dichlorethane (see Ethylene Dichloride).

Dichlorodiethyl Ether, against Mycetophilids on cucumber, 145.

Dichlorodiphenyl Oxide, logs and timber treated with, against ambrosia beetles, 427.

Dichloroethyl Ether, injected into apple trees against Saperda candida, 243; methods of soil treatment with, against pests, 122, 123, 198, 261, 299; sprayed with, against Crambus, 408; and oil for treating maize ears against Heliothis armigera, 217, 481; effect of, on plants, 122.

dichomeridis, Meteorus.

Dichomeris eridantis, parasites of, on Dalbergia sissoo in India, 74. Dichomeris oceanis, bionomics of, in Japan, 100.

dichromellus, Chilo. Dichroplus (see Trigonophymus). Dictyophara, on cotton in Philip-

pines, 649. Dictyoploca japonica, new parasite

of, in Japan, 480. dictyoplocae, Apanteles.

dictyospermi, Chrysomphalus.

diemenensis, Eupsenella.

Diervilla, Xyleborus dispar on, in Br. Columbia, 328.

differentialis, Melanoplus.

Digama marmorea, noxious weed defoliated by, in Queensland, 467. Digitalis purpurea, winter food-plant of Macrosiphum solani in

Maine, **188.**

Di-isobutylketone, intermediary solvent for derris resins used with oils, 350.

dilophonotae, Telenomus.

Dilophus febrilis, bionomics and control of, in turf in Br. Isles, 596, 527.

Dimethyglyoxime, tests of, in baits for locusts, 614.

dimidiatus, Carpophilus. dimmocki, Pikonema.

Dinapate wrighti, observations on, in California, 137.

Dinarmus coimbatorensis, hosts and ecology of, in Madras, 522; parasite of Lophobaris piperis in Netherlands Indies, 33; considered a synonym of D. sauteri, 522.

Dinarmus sauteri, D. coimbatorensis considered a synonym of,

Dindymus versicolor, measures against, on fruit trees, etc., in New South Wales, 290.

Dinitro-cresol, 70; in dormant sprays against orchard pests, 108, 109, 284; in spray against Coccid on pine, 605; in dusts: against locusts and grasshoppers, 20, 326, 327, 538; against other insects, 19, 326. (See Sodium Dinitro-

cresylate.)

Dinitro-ortho-cyclohexyphenol, and oils, in dormant sprays against orchard pests, 128, 134, 191; other uses of, in sprays, 307, 605, 641, 644; in oils for treating maize ears against Heliothis, 641; in dusts against mites, 249, 561; tests of dusts and sprays containing salt of, 560, 561; no danger of residues from dusts of, 647; causing injury to plants, 249, 561, 641.

Dinoderus, parasite of, in stored bamboos in India, 268.

Dinoderus minutus, on bamboo in Fiji, 170; in stored bamboos in Porto Rico, 472.

dinota, Tortrix.

Dioctes (see Angitia).

Diogmites discolor, attacking Lamellicorn larvae in Kentucky, 299.

diophthalmus, Tectocoris.

Dioscorea, Apogonia cribricollis on, in Malaya, 512.

Diospyros kaki (see Persimmon).

diparopsidis, Apanteles.

Diparopsis castanea (on cotton), parasites of, in Nyasaland, 153, 588; in Sudan, 386.

Diphenylene Ketone Oxide, 411. Diplogaster aerivora, parasite of Nodonota puncticollis in Pennsylvania, 192.

Dibloneura cornuta (cleghorni), bred from dead larvae of Heliothis

armigera in Fiji, 390.

Diplotaxis sordida, in forest nurseries in New York, 421, 422.

Diprion frutetorum (see Gilpinia). Diprion nipponicus, parasite of, in Japan, **294.**

Diprion pini, Europe, 294. parasites of,

Diprion polytomum (see Gilpinia). Diprion sertifer (see Neodiprion). Diprion similis, parasite of,

Poland, 294.

diprioni, Exenterus.

diprionis, Delomerista; (see E. diprioni). Exenterus

Diptera (parasitic on insects), records of, in Germany, 70.

dipterocarpi, Trochorrhophalus.

Dipterocarpus tuberculatus, Dasychira cerigoides on, in Burma, 78. Dirhinus, released against fruit-

flies in Fiji, 495, 496.

Dirhinus auratus (parasite of fruitflies), breeding and liberation of, in Ceylon, 594; imported into Mauritius, 610.

Dirhinus banksi, hyperparasite of Artona catoxantha in Java, 79.

Dirhinus giffardi, rearing and utilisation of, against Anastrepha spp. in W. Indies, 471.

discolor, Diogmites; Eutermes (Nasutitermes); Verania (Alesia).

Dishcloth Gourd (see Luffa cylin-

disjuncta, auct., Microphthalma (see M. nigra).

Disophrys sissoo, hosts of, in India, 74, 263; utilisation of, 74.

dispar, Lepidosaphes (Coccomytilus); -Lymantria (Porthetria); Xyleborus.

disparis, Pimpla.

Dissolcus rufiventris, parasite of Eurygaster integriceps in Russia, 582.

introduced Dissosteira carolina, parasite not developing in, in Canada, 309; bionomics of, in U.S.A., 604.

Dissosteira longipennis, in U.S.A., 48. disstria, Malacosoma.

distigma, Chilocorus.

distincta, Galerucella.

distinguendus, Lariophagus.

Ditremena occidentalis, Fundella cistipennis on, in Porto Rico, 345. Diversinervus elegans, parasite of Saissetia oleae in E. Africa, 385; introduced into California, 385.

Diversinervus masakaensis, sp. n., parasite of Saissetia oleae in Uganda, 385; introduced into California, 385.

Diversinervus smithi, sp. n., parasite of Saissetia oleae in S. Africa, 385; introduced into California, 385, 386.

diversus, Archichauliodes; Microdus (Bassus).

divisus. Macrosteles (Cicadula).

Dobson-fly (see Archichauliodes diversus).

Dociostaurus crucigerus subsp. brevicollis, in Russia, 570.

Dociostaurus maroccanus, bionomics and control of, in Turkey and Spain, 289, 459-461, 529.

doclea, Pogonortalis.

Dodecyl Thiocyanate (Rhodanate), 108, 111. (See Thiocyanates.)

Dogstail, Crested (see Cynosurus cristatus).

Dogwood (see Cornus).

dohertyi, Leucoplema (Epiplema).

Doirania leefmansi (egg-parasite of Sexava), distribution of, **380.** dolenda, Pheidole javana.

Dolichocolon rufescens, parasite of Pandesma quenavadi in India,

Dolichos lablab, pests of, in Sudan, 387, 388.

Dolichos pseudopachyrhizus, insecticidal properties of, 483.

domestica, Musca; Thermobia. domesticus, Calotermes; Gryllulus (Gryllus); Opilo.

dominica, Rhizopertha.

Dominican Republic, Coccid pests and their biological control in, 498; parasite of Anastrepha introduced into, from Porto Rico,

Doralis (see Aphis).

dorsalis, Aeolus (see Drasterius mellillus); Chalepus; Dacus (Chaetodacus) ferrugineus; Deltocephalus; Doryphorophaga.

dorsata, Apis.

Dorthesia (see Icerya purchasi).

dorycus, Euagoras.

Doryphora decemlineata (see Leptinotarsa).

Doryphorophaga dorsalis, parasite of Calligrapha in Quebec, 627.

Dosage-Mortality Curves, 433, 434. Douglas Fir (see Pseudotsuga taxifolia).

Dracaena, Pseudococcus adonidum on, in California, 298, 355; injured by methyl bromide fumigation, 555.

drakei, Lachnosterna.

Drasterius mellillus, bionomics of, on tobacco in Kentucky, 363, 482; on sugar-cane, etc., in Florida, 476; measures against, 476, 477, 483.

Drepanosiphum platanoides, potato growing near sycamore in

Eire, **329.**

Drepanothrips reuteri, Leptothrips mali predacious on, on vines and fruit trees in California, 210.

Dreyfusia nüsslini (see Chermes nordmannianae).

Drosicha stebbingi, observations on,

in Punjab, 269. Drosicha townsendi, on cotton in

Philippines, 649.

Drosophila, parasitised by Phaenocarpa leveri in Fiji, 496; technique of rearing and using, for tests of contact sprays, 311.

Drosophila hydei, 311.

Drosophila melanogaster, 221, 311. Drosophila suzukii, bionomics of, on fruit trees and vines in Japan,

Dryinus pyrillae, parasite of Pyrilla in India, 82, 184.

Dryocoetes, on spruce in Canada, 532.

dubius, Ootetrastichus.

dulcivora, Cosmopteryx.

dumbletoni, Meteorus.

Dung Beetles (see Aphodius). duodecimpunctata, Crioceris; Diabrotica.

duplicatus, Necremnus. Du Pont's Repellent, 595.

Dusting, types of equipment for, 47, 52, 91, 162, 249, 250, 271, 408, 504, 513, 569, 630, 638, 654; use of aircraft for, 156, 300, 438, 538; spraying compared with, 42, 135, 197, 202, 296, 297, 463, 466.

Dusts, use of coal-tar distillates as insecticides in, 247; use of oilimpregnated, 161, 197, 343, 504, 561, 569, 647, 654; other conditioning agents for, 250, 353, 504; apparatus for dispersing, with liquid adhesives, 412; effect of electrostatic charges on, 569; equipment for testing, 415, 557; apparatus for determining particle diameter of, 397.

Dutch Elm Disease (see Ophiostoma ulmi).

dysderci, Paraphasiana.

Dysdercus columbicus, associated with boll rots on cotton in Colombia, 53.

Dysdercus fasciatus, food-plants of, in Sudan, 387.

Dysdercus fulvoniger, associated with boll rots on cotton in Colombia, 53.

Dysdercus mendesi, new parasite of, in Brazil, 469.

Dysdercus mimus, on cotton in Peru, 539.

Dysdercus nigrofasciatus, on kapok and Sterculia in Sudan, 387.

Dysdercus peruvianus, associated with boll rots on cotton in Colombia, 53.

Dysdercus poecilus, on cotton in

Philippines, 649.

Dysdercus ruficollis, in Brazil, 469; on cotton in Peru, 510, 539; parasites of, 269, 510; foodplants and migrations of, 539.

Dysdercus sidae, measures against, on fruit trees, etc., in New South

Wales, 290.

Dysdercus superstitiosus, food-plants of, in Sudan, 387.

E.

Earias (on cotton), hyperparasite of, in Nyasaland, 588; in Philippines, 648.

Earias insulana, on cotton in Sudan, 386, 387.

eatoni, Cylindrocopturus.

eccoptogastri, Rhoptrocerus (Pachy-

Echinocystis fabacea, vector of mosaic of, in squash in California, 501.

echinus, Uventius.

echion, Tmolus.
Echthrodelphax bicolor, bionomics of,

in Japan, 101. Echthromorpha conopleura, parasite of Spodoptera mauritia in Guam,

331.

Echthromorpha notulatoria, parasite of Hyblaea puera in Burma, 77.

Eciton nigrescens, destroying Solen-

opsis spp. in Florida, 476. Economic Entomology, book on,

Economic Entomology, book on, 189.

Ecpantheria icasia, on vanilla in Porto Rico, 469.

Ectropis deodarae, effect of grazing on outbreaks of, in India, 74. ectypus, Pheletes (Limonius).

Ecuador, danger of introduction of Platyedra gossypiella into Peru from, 539. egena, Pentilia.

Egg Albumen, as a dispersing agent for sprays, 113; in baits for *Rhagoletis pomonella*, 563.

Egg-plant (Solanum melongena), Gnorimoschema operculella on, in Bermuda, 138; Epitrix cucumeris on, in Connecticut, 162; pests of, in Hawaii, 34, 35; pests of, in India, 265, 266, 516, 517, 523, 542; Acanthocoris sordidus on, in Japan, 171; insects and virus disease in, 140, 542.

Egius platycephalus, establishment of, against Asterolecanium spp. in Porto Rico, 471.

Egypt, Coccids damaging roots of plants in, 30; Platyedra gossypiella on cotton in, 338; parasite introduced into U.S.A. from, against P. gossypiella, 430.

Elachertus, parasite of Lithocolletis

in Porto Rico, 473.

Elachertus evetriae, parasite of Cydia latiferreana in Oregon, 406.

Elaeagnus, Aonidiella aurantii on, in Japan, 102.

Elaeis guineënsis (see Oil Palm). Elaphrium simaruba, Zethus rufinodus damaging fence posts of, in Porto Rico, 473.

Elasmopalpus lignosellus, on maize and Euchlaena in Porto Rico,

469

Elasmus albicoxa, parasite of Parornix prunivorella in Kansas, 190. Elasmus albomaculatus, parasite of lac predator in India, 523.

Elasmus johnstoni, hyperparasite of Sylepta derogata in India, 524.

Elasmus nephantidis, parasite of Nephantis serinopa in India, 620.

eleaphilus, Coccophagus.

electo, Colias.

Electricity, for sterilising cereal products, 605.

Electrostatic Charges, effect of, on insecticidal dusts, **569**.

elegans, Bruchus; Cheiloneurus; Diversinervus; Microterys.

Elephant Yam (see Amorphophallus campanulatus).

Elettaria cardamonum var. major, effect of Poecilips cardamomi on, in India, 74.

Eleusine, locusts damaging, in Uganda, 228, 229.

Eleusine coracana, planted to attract Laphygma exigua in India, 20.

Elgetol (containing sodium dinitrocresylate), uses of, in sprays, 128, 133, 330.

elisus, Lygus.

Ellice Islands, Papuana on Colocasia esculentum in, 171; Bufo marinus imported into, 496.

elliotti, Aulocara.

Elliptone, 135.

ello, Erinnyis.

Ellopia fiscellaria, on conifers in Quebec, 36.

Elm, Galerucella luteola on, in Rumania, 12; pests of, in U.S.A., 40, 116, 119, 120, 163, 212, 276, 364, 365, 416, 427, 502, 550, 551, 563, 595, 606, 627, 628, 635, 644, 645; grasshoppers feeding on, 603; relation of Scolytids to Ophiostoma ulmi in, 116, 119, 212, 364, 365, 502, 550, 563, 606, 644, 645, 646; other possible vectors of O. ulmi in, 40, 644; other fungi associated with Scolytus multistriatus on, 116; injection of chemicals into, against S. multistriatus, 212; treated with selenium against Tetranychus telarius, 276.

Elm Caterpillar, Spiny (see Nymphalis antiopa).

Elm Leaf Beetle (see Galerucella luteola).

Elm Scale, Brown (see Lecanium corni).

Elodia flavipalpis, parasite of Cydia molesta in Japan and Korea, 372.

Elongate Mite (see Tyrophagus

noxius). elongatus, Trigonophymus (Dichroplus).

elutella, Ephestia.

Elytroteinus subtruncatus, marinus destroying, on lemon in Fiji, 389.

emarginatus, Ips. emigratella, Amorbia.

Emilia sonchifolia, vector and identity of virus disease of, in Hawaii, 139, 140.

emissaria, Scopula.

Emmaloceva depressella, parasites of, on sugar-cane in India, 183.

Empoasca, in Argentina, 608; in Fiji, 147; in Peru, 510, 511; on cotton, 147, 510; list of species of, attacking cotton, 516; susceptibility of varieties of cotton to, 148; observations on, in connection with virus diseases of solanaceous plants, 511, 608; measures against, 148, 510.

abrupta, measures Empoasca against, on potato in New Mexico, 638, 639.

Empoasca binotata, sp. n., foodplants of, in India, 265.

Empoasca devastans (in India), bionomics and control of, on cotton, etc., 516, 517; transmitting virus disease of egg-plant, 542.

Empoasca fabae, on fruit trees in Ontario, 305, 306, 548; in U.S.A., 39, 163, 248, 256, 325, 439, 476, 503; on beans, 248, 439, 476; not transmitting cowpea mosaic, 503; on potato, 39, 163, 325; on rose, 256; types of injury to plants by, 256, 305, 548; measures and experiments against, 39, 163, 248, 325, 439-441; Phyllocoptes fockeui spread by, **306.**

Empoasca fabalis, susceptibility of varieties of lima beans to, in Porto Rico, 470.

Empoasca flavescens, measures against, on cotton in Philippines, 648.

Empoasca kerri, sp. n., with var. motti, n., food-plants of, in India, 266.

Empoasca lybica, food-plants of, in Sudan, 387.

Empoasca maligna, type of injury to rose by, in U.S.A., 256.

Empoasca minor, sp. n., on grapevine and cotton in India, 265.

Empoasca parathea, sp. n., foodplants of, in India, 266.

Empoasca punjabensis, sp. n., foodplants of, in India, 266.

Empoasca solana, food-plants of, in Hawaii, 35, 650, 651; experiments with virus disease of papaya and, 35.

Empoasca solanifolia, sp. n., potato in India, 265.

Empoasca tabaci, sp. n., on tobacco in India, 266.

Empusa acaricida, sp. n., infesting Halotydeus destructor in W. Australia, 587.

Empusa grylli, infesting locusts and grasshoppers, 229, 491, 570, 589. Enarmonia caryana, on pecan in U.S.A., **159.**

Encarsia flavoscutellum, introduced into Formosa against Oregma lanigera, 92.

Encarsia formosa (parasite of Trialeurodes vaporariorum), in Britain, 374; introduced into Canada, 310; utilisation of, in Victoria, 32; reared on Aleurodes proletella, 374.

Encarsia partenopea, parasite of Siphoninus phillyreae in Britain, 374; reared on Aleurodes proletella, 374.

Encoptolophus sordidus costalis, experimentally parasitised by Muscina stabulans in Texas, 477.

encyrtcides, Systasis.

Encyrtus fuliginosus, sp. n., parasite of Saissetia oleae in Africa, 385; introduced into California, 385, 386.

Endive, insects and virus diseases in, 140, 646.

Endoconidiophora fimbriata, Cylas fermicarius associated with, on sweet potato in Loochoo Islands, 97,

endogena, Polyodaspis. endonis, Catolaccus.

Endospermum, Xyleborus rameus

on, in Fiji, 169. Endothia parasitica, possible vector of, on chestnut in Japan, 97.

Engytatus geniculatus, bionomics of, in Hawaii, 34, 650, 651.

Engytatus notatus, not transmitting virus disease of tobacco and tomato in Argentina, 608.

Engytatus tenuis, measures against, on tobacco in Sumatra, 319.

Enicospilus, parasite of Plecoptera reflexa in India, 74.

Enicospilus purgatus, parasite of Cirphis unipuncta in U.S.A., 300.

enigma, Oedaleonotus. ensigera, Agalliana.

Entedon pempheridis, hosts and ecology of, in Madras, 522.

Entodecta (see Metallus).

Entomology, books on, 189, 275, 625.

Entomophthora sphaerosperma, infesting Plutella maculipennis in S. Africa, 19.

Entyposis impressa, parasites of, in Zanzibar, 83.

Eperua falcata (Wallaba Wood), termites imported into Barbados in, 6; resistance of, to termites, 7, 138.

Ephestia, Trichogrammatids reared on eggs of, 264, 265, 610.

Ephestia cautella, on coconut and other palms in Brazil, 9; in

stored cacao, 616, 617; in copra, 9, 377; in stored grain, 97; in stored tobacco, 19; parasitised by *Microbracon hebetor*, 377, 616; measures against, 19, 377.

Ephestia elutella, in stored cacao, 616, 617; in stored tobacco, 362, 365, 429; field sprays affecting infestation by, 365; tests of atomised sprays on, 434; fumigation against, 18, 362; parasite of, 616; water vapour affecting survival of, at reduced pressure, 302.

Ephestia figulilella, cold storage of dried fruit against, in U.S.A., 426. Ephestia kuehniella, associated with honey bees in California, 418; influence of cold on pupation of,

Ephialtes, factor affecting parasitism of Cydia pomonella by, 219; application of name, 220.

Ephialtes comstocki, parasite of Rhyacionia buoliana in Br. Columbia, 415.

Ephialtes conquisitor and evetriae (see Pimpla).

Ephialtes grapholithae, parasite of Cydia pomonella in U.S.A., 193.

Ephialtes manifestator, type of genus, 220.

Ephialtes nucicola, parasite of Cydia latiferreana in U.S.A., 405.

Ephialtes nuda, obesa and pedalis (see Pimpla).

ephilida, Lachnosterna (Phyllophaga).

ephippella, Argyresthia.

Epicampocera succincta, parasite of Pieris rapae in Britain, 314.

Epicauta hirticornis, destroying locust eggs in Formosa, 94.

Epicauta megalocephala, destroying grasshopper eggs in Siberia, 485.

Epicanta pennsylvanica (in U.S.A.), bionomics of, destroying grass-hopper eggs, 566; measures against, on asters, 361.

Epicauta waterhousei, destroying locust eggs in Formosa, 94.

Epicometis hirta, parasite of, in Russia, 572.

Epidiaspis leperii, food-plant affecting control of, by sprays in Italy, 155.

Epilachna spp., parasites of, in Mexico and Brazil, 114, 115, 431. Epilachna borealis, 114.

Epilachna corrupta (see E. varivestis).

Epilachna defecta, 114.

Epilachna mexicana, 114, 115.

Epilachna obscurella, 114.

Epilachna varivestis (Mexican Bean Beetle), in U.S.A., 39, 114, 208, 247, 431, 503, 601; effect of temperature on, 601; not transmitting cowpea mosaic, 503; attempted introduction of parasites against, 114, 431; tests and use of insecticides against, 39, 208, 247, 601.

Epilachna vigintioctomaculata, bionomics and control of, on potato

in Korea, 103.

Epilachna vigintioctopunctata, foodplants of, in Fiji, 390, 495; distribution of, in Victoria, 31; measures against, 390.

Epilachna virgata, 114. epilachnae, Paradexodes.

Epilobium angustifolium, effect of, on oviposition by Silpha opaca,

Epiplema dohertyi (see Leucoplema).

epiplemae, Euplectrus.

Epipyrops, predacious on Pyrilla perpusilla in Punjab, 82.

Epitragus aurulentus, on cotton and maize in Colombia, 54.

Epitrimerus gigantorhynchus, associated with Phyllocoptes fockeui on plum in Ontario, 306.

Epitrix, not transmitting virus disease of tobacco and tomato in Argentina, 608; measures against, on potato in New Mexico, 638, 639.

Epitrix cucumeris (in U.S.A.), on potato, 162, 221, 247, 278, 503, 504; on other plants, 162, 221; bionomics of, 221, 278, 503; measures against, 162, 247, 278, 503, 504.

Epitrix parvula (Tobacco Fleabeetle), in U.S.A., 200, 201, 404, 430; bionomics of, 200, 201; measures against, 404, 430.

Epurea luteola, in decayed fruits in Fiji, **390.** eques, Torocampus.

Erannis defoliaria, parasite of, in Britain, 80.

evecta, Phoroceva.

eremita, Bostrychopsis.

Erencyrtus dewitzi, parasite Laccifer lacca in India, 263.

Eretmocerus serius, value of, parasitising Aleurocanthus woglumi in Bahamas, 293.

eveunetiphila, Melittobiopsis.

Ereunetis flavistriata (see Decadarchis).

ericae, Nysius.

Ericerus pela, bionomics and utilisation of, in Szechuan, 153. evichsoni, Pristiphora.

eridania. Laphygma

(Prodenia, Xylomyges).

eridantis, Dichomeris.

Erinnyis ello, parasites of, on cassava in Colombia, 9.

Eriocampoides (see Caliroa).

Eriococcus, possible parasites of, in Argentina, 344.

Eriococcus coriaceus (on Eucalyptus), natural enemies of, in Australia, 290, 545, 546; attempted introduction of predators into New Zealand against, 545.

Eriophyes, measures against, on Juniperus in Japan, 99, 100.

Eriophyes avellanae, on Corylus in California, 223.

Eriophyes gossypii, on cotton in S. America, 54, 511.

Eriophyes ribis, tests of sprays against, on black currant in Britain, 281.

Eriophyes ryderi, thrips probably predacious on, on maple California, 210.

Eriophyes sheldoni, on Citrus in Queensland, 465.

Eriophyes triradiatus, type of injury to species of willow by, in Britain, 146.

Eriophyes vaccinii, sp. n., on blueberry in N. Carolina, 259.

Eriophyes vitis, thrips probably predacious on, on grape-vine in California, 210.

Eriopis connexa, predacious on Toxoptera graminum in Argentina, 36.

Eriosoma lanigerum (on apple), new Encyrtid parasite of, in Argentina, 445; in Britain, 287; in Germany, 68, 231; in New South Wales, 493; in New Zealand, 333; in Peru, 511, 539; in U.S.A., 164, 299, 300, 360, 361; canker associated with, 112; effects of cold on, 68, 333; Aphelinus mali parasitising, and its utilisation, 333, 361, 493, 511; sprays against, 231, 287, 360; soil fumigants against, 299, 300.

Eritrea, Pseudococcus citri on wild olive in, **155.**

Erixestus winnemana, parasite of Calligrapha in Quebec, 627.

Ernobius mollis, damaging building timbers in Sweden, 324.

erosa, Calandra.

Eryngium planum, propagation of, to attract Scoliids in Russia, 572.

Erynnia mitida (parasite of Galerucella luteola), bionomics and attempted establishment of, in California, 416, 417.

Erythrina spp., insects on, in Porto

Rico, 52, 469.

erythrocephala, Calliphora.

Erythroneura (in U.S.A.), type of injury to rose by, 256; measures against, on vines, 117, 558.

erythrothoracis, Pristomerus chinen-

sis (vulnerator).

Ether, effects of, on toxicity of fumigants to *Tribolium confusum*, 327; as a solvent, 29, 352.

327; as a solvent, 29, 352. Ethyl Acetate, effect of ether on toxicity of, to *Tribolium confusum*, 327; in formula for treating timber against *Dermestes*, 393.

Ethyl Alcohol (see Alcohol).

Ethylene Dichloride, emulsion of, against Aegeria exitiosa, 160; fumigation of stored cowpeas with, against Bruchus chinensis, 59; and carbon tetrachloride, against pests of stored grain, 278, 429; as a solvent, 586.

Ethylene Oxide, 434; and carbon dioxide, vacuum fumigation with, against pests damaging books,

536.

Etiella zinckenella, measures against, on beans, etc., in Porto Rico, 345.

Euagoras dorycus, predacious on Amblypelta cocophaga in Solomon Is., 325.

Euaresta aequalis, introduced into Australia against Xanthium pungens, 143.

Eubadizon extensor, parasite of Cydia molesta in Japan, 372.

Eubadizon pleuralis, utilisation of, against Cydia molesta in Connecticut, 87.

Eublemma, predacious on Laccifer lacca in Siam, 94.

Eublemma amabilis, predacious on Laccifer lacca in India, 263, 264, 265, 523; parasites and biological control of, 264, 265.

Eublemma costimacula, predacious on Pseudococcus perniciosus in

Nyasaland, 588.

Euborellia stali, bionomics of, on ground-nut, etc., in India, 518.

Eucalymnatus tessellatus (on coconut), Coccinellids introduced into Seychelles against, 324.

eucalypti, Phylacteophaga.

Eucalyptus, Gonipterus scutellatus on, in S. Africa, 18, 30; pests of, in Australia, 30, 237, 290, 335, 545; Phymatioderus bizonatus on, in Chile, 512; Ericococcus coriaceus on, in New Zealand, 545; pests of, in Nyasaland, 30, 61; G. scutellatus on, in S. Rhodesia, 30; Icerya seychelarum on, in Seychelles, 82.

Eucalyptus crebra, Maskellia globosa causing witches' broom on, in

Queensland, 467.

Eucalyptus maculata, determination of starch content of timber of, in relation to Lyctus in Queensland, 150.

Eucalyptus saligna, new weevil on, in Zululand, 341.

Eucalyptus viminalis, Coleoptera on, in Argentina, 512.

Eucalyptus Snout Beetle (see Gonipterus scutellatus).

Eucelatoria australis, parasite of Lepidoptera on cotton in Peru, **510**. Eucera, parasitised by Myopa in

Europe, 474.

eucera, Eucoila (Rhoptromeris).

Euchlaena mexicana, Elasmopalpus lignosellus on, in Porto Rico, 469; susceptible to Aplanobacter stewarti in U.S.A., 245.

Euchlaena perennis, susceptible to Aplanobacter stewarti in U.S.A., 245.

Eucoila, parasite of Phytomyza spicata in Fiji, 170.

Eucoila eucera, hosts of, in Russia, 618.

Eucomys littoralis, sp. n., parasite of Pulvinaria convexa in Argentina, 344.

Eucosma schistaceana (on sugarcane in Formosa), high temperatures causing sterility in, 172; failure to establish parasites against, 92.

Euderus, parasite of Lophobaris piperis in Netherlands Indies, 33.

Euderus lividus (parasite of Agromyza obtusa), bionomics of, in India, 267.

Euderus pempheriphila, bionomics of, in Madras, 522.

Euetheola rugiceps, factors affecting damage to maize and sugar-cane by, in U.S.A., 2, 3, 601.

Eufallia unicostata, feeding on fungi in houses in Porto Rico, 361.

Eugenia, Dacus spp. on, in Malaya, 90; thrips intercepted on, in New York, 86.

Eugenia aromatica (see Clove).

Eugenia jambolana, new mite on, in Punjab, 271.

Eugenia jambos, Coccid on, in Formosa, 622.

Eugenia malaccensis, insects associated with galls on, in Fiji, 390.

Eugenol, in baits for Popillia, 126, 364, 596.

Eulachnus agilis, measures against, on pines in U.S.A., 607.

Eulachnus rileyi, E. agilis confused with, on pine in U.S.A., 607.

Eulecanium (see Lecanium).

Eulophus femoralis, parasite of Cosmopteryx dulcivora in Java, 524.

salaminia, damaging Eumaenas Citrus fruit in Ceylon, 594.

Eupatorium odoratum, Bemisia tabaci and virus disease in, in Sumatra, 318.

Eupelmella pedatoria, parasite of Euderus pempheriphila in Madras, 522.

Eupelmella vesicularis, parasite of Mayetiola destructor in Russia, 619; search for hosts and egg distribution of, 490.

Eupelmus allyni, hosts of, in U.S.A.,

Eupelmus atropurpureus, parasite of Mayetiola destructor in Russia, 582, 618.

Eupelmus catoxanthae, sp. n., parasite of Artona catoxantha in Java and Malaya, 79.

Eupelmus curculionis, parasite of Lophobaris piperis in Netherlands Indies, 33.

Eupelmus microzonus, parasite of Mayetiola destructor in Russia, 582, 618.

Eupelmus tachardiae, parasite of Laccifer lacca in India, 263.

Eupelmus urozonus; hosts and ecology of, in Madras, 522.

Euphorbia hirta, Bemisia tabaci transmitting virus disease to tobacco from, in India, 183.

euphorbiae, Aphthona.

Euphoria longana, cultivation of lac on, in Siam, 94.

euphytus, Rhizotrogus.

Euplectrini, revision of, 513.

Euplectromorpha artonae, sp. n., parasite of Artona catoxantha in Java, 79.

Euplectromorpha kiambuensis, sp. n., parasite of Tortrix dinota in Kenya, 514.

Euplectromorpha malayensis, sp. n., hosts of, in Malaya and Java, 514.

Euplectromorpha obscurata, sp. n., hosts of, in Africa, 514.

Euplectromorpha salomonis, sp. n., parasite of Zygaenid in Solomon Is., 514.

Euplectromorpha viridiceps, sp. n., parasite of Artona catoxantha in Java and Malaya, 79.

Euplectrus, utilisation of, against Plecoptera reflexa in India, 74; parasite of Niphadolepis alianta in Nyasaland, 154; habits of, 514.

Euplectrus cacoeciae, sp. n., parasite of Tortrix rosana in Bulgaria, 514. Euplectrus coimbatorensis, sp. n.,

host of, in India, 514.

Euplectrus epiplemae, sp. n., hosts of, in Africa, 514.

Euplectrus hargreavesi, sp. n., host of, in Uganda, 514.

Euplectrus indicus, sp. n., hosts of, in India, 514.

Euplectrus laphygmae, sp. n., hosts of, in Africa, 514.

Euplectrus liparidis, sp. n., parasite of Lymantria dispar in Algeria, 514.

Euplectrus parvulus, sp. n., hosts of, in India and Burma, 514.

Euplectrus petiolatus, sp. n., parasite of Lymantriid in India, 514.

Euplectrus phthorimaeae, sp. n., parasite of Gnorimoschema operculella in Cyprus, 514; in Palestine, **514.**

Euplectrus platyhypenae, parasite of Lepidoptera in Cuba and Hawaii, 538, 650.

Euplectrus singularis, sp. n., hosts of, in Uganda, 514.

euploeae, Brachymeria.

Euprepocnemis plorans, on vegetables in Spain, 463. euproctidis, Trichogramma.

Euproctis chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Euproctis terminalis, bionomics and control of, in forests in S. Africa,

Eupsenella diemenensis, parasite of Tortrix postvittana in Tasmania, 151.

Eupteromalus, hyperparasite of Hypera variabilis in California,

Eupteromalus micropterus, parasite of Mayetiola destructor in Russia, 582, 618, 619; bionomics of, 619.

Euribia jacaena, search for hosts by parasites of, 490.

europaea, Microphthalma.

Europe, study of insect outbreaks in, 11.

European Earwig (see Forficula auricularia).

Eurygaster, type of injury to cereals by, in Russia, 574, 575.

Eurygaster austriaca, bionomics of,

on cereals in Russia, 575.

Eurygaster integriceps (on cereals in Russia), bionomics of, 574, 582; parasites and biological control of, 575, 582; experiments with fungus and bacteria against, 583; other measures against, 575.

Eurygaster maura, bionomics of, on

cereals in Russia, 574.

Euryischia comperei, hyperparasite of Eriococcus coriaceus in Australia, 546.

eurytheme, Colias (Eurymus).

Eurytoma, parasite of Nephantis serinopa in Ceylon, 620, 621.

Eurytoma albotibialis, parasite of Artona catoxantha in Java and Malaya, 79.

Eurytoma appendigaster, parasite of Apanteles liparidis in Japan, 172. Eurytoma curta, search for hosts and egg distribution of, 490.

Eurytoma fellis, measures against, on Citrus in New South Wales, 144.

Eurytoma promecothecae (parasite of Promecotheca papuana), bionomics of, in New Guinea Territory, 378.

Eurytoma robusta, search for hosts and egg distribution of, 490.

Eurytoma tylodermatis, parasite of Chalcodermus aeneus in U.S.A., 405.

Euscepes postfasciatus (batatae), on sweet potato in Bermuda and Brazil, 139, 511.

Euschistus tristigmus, on tomatos in New York, 121.

Euschistus variolarius, bionomics of, on tomatos in New York, 121; tests of insecticides against, 121. Eutachina (see Tachina).

Eutermes acajutlae, damaging living trees in Porto Rico, 473.

Eutermes costalis (in West Indies), damage to sugar-cane by, 6; attacking trees and timber, 6, 7, 473; timbers resistant to, 7; measures against, 6, 7.

Eutermes discolor, damaging trees

in Porto Rico, 473.

exitiosus, Eutermes measures against, damaging timber and cables in Australia, 142.

Eutermes octopilis, imported into Barbados in wallaba wood from

Br. Guiana, 6.

Eutetrapha tridentata (on elm in New York), bionomics of, 627; possible relation of, to Ophiostoma ulmi, 644.

Eutettix, not transmitting virus disease of tobacco and tomato in Argentina, 608.

Eutettix phycitis, transmitting virus disease of solanaceous plants in

Madras, 542.

Eutettix tenellus (in U.S.A.), transmitting curly-top of beet and tomato, 134, 631; experiments with curly-top virus and, 66. 501; studies of migrations of, 533, 631; bird destroying, 563; measures against, 134, 631, 632.

Euxanthellus, parasite of Saissetia oleae in S. Africa, 385; introduced into California, 385.

stigmatias, Euxesta measures against, on maize in Porto Rico, 469, 470.

Euxoa segetum (see Agrotis).

Euxoa tritici, bionomics of, Germany, 449, 450. evanescens, Trichogramma.

Evergestis rimosalis, laboratory rearing of, 557.

Evetria (see Rhyacionia).

Elachertus; evetriae, Pimpla (Ephialtes).

ewingi, Persectania (see P. composita).

excessana, Tortrix.

exclamationis, Agrotis.

exempta, Laphygma.

Exenterus, utilisation and value of, against Gilpinia polytoma Canada, 39, 293, 294, 531, 532.

Exenterus abruptorius (parasite of Diprionine sawflies), in Europe, 86, 294; in Japan and Korea, 294; imported into Canada and U.S.A., 294, 637.

Exenterus adspersus (parasite of Diprionine sawflies), in Europe and Far East, 294; imported into Canada and U.S.A., 294, 637; synonymy of, 294; probable erroneous records of, 39, 294.

Exenterus affinis, parasite of Neodiprion in U.S.A. and Canada, 293; characters of, 294.

Exenterus canadensis, parasite of Neodiprion in U.S.A., 294.

Exenterus claripennis (parasite of Gilpinia polytoma), in Czecho-slovakia and Sweden, 294; utilisation and value of, in Canada, 532; introduced into Maine, 637.

Exenterus diprioni, parasite of Neodiprion in U.S.A. and Canada, 293; characters of, 294.

Exenterus diprionis, emendation for E. diprioni (q.v.), 293.

Exenterus flavissimus, sp. n., parasite of Neodiprion sertifer in New Jersey, 294.

Exenterus hullensis, in Canada and U.S.A., 294.

Exenterus laricinus (see E. adspersus).

Exenterus lophyri, parasite of Zadiprion grandis in Nebraska, 294.

Exenterus marginatorius, parasite of Diprionine sawflies in Europe, 86, 294; establishment of, against Gilpinia polytoma in Canada, 294; imported into Maine, 637; recorded as E. adspersus, 637.

Exenterus oleaceus (see E. adspersus).

Exenterus oriolus, parasite of Neodiprion sertifer in Hungary, 294.

Exenterus pini, sp. n., in U.S.A., 294.

Exenterus tricolor, parasite of Gilpinia polytoma in Czechoslovakia, 294; imported into Canada, 294.

Exenterus tsugae, sp. n., parasite of Neodiprion tsugae in Oregon, 294, 552.

Exenterus vellicatus, sp. n., parasite of Gilpinia polytoma in Europe, 294; imported into Canada, 294. Exeristes (see Pimpla).

exigua, Laphygma (Spodoptera); Susumia.

exiguae, Hyposoter.

exitiosa, Aegeria (Conopia, Synan-thedon).

exitiosus, Eutermes.

Exochomus flavipes, predacious on Chironomids in Madagascar, 83; introduction and utilisation of, against Coccids in Seychelles, 82, 324.

Exochomus orbiculus, introduced into Porto Rico from Brazil against Asterolecanium spp., 471.

Exochomus ventralis, introduced into Seychelles from Tanganyika against Coccids, **324**.

Exoprosopa fasciata, attacking Lamellicorn larvae in Kentucky, 299.

Exora apicipennis, on Aleurites in Nyasaland, 62.
Exorista (see Zenillia).
experta, Feltia.
expulsus, Apanteles.
extensor, Eubadizon.
extinctalis, Nodaria.
exuberans, Sarcophaga.
ezea, Achaea.

F.

fabae, Empoasca. fabalis, Empoasca. faber, Achaea. faceta, Antestia. facialis, Catascopus. fagata, Operophtera.

Falco sparvarius, destroying noxious insects in Utah, 217.

Fall Armyworm (see Laphygma frugiperda).

fallax, Piezosternum; Xyleborus.

farinae, Tyroglyphus. fascialis, Hymenia

fascialis, Hymenia (see H. recurvalis). fasciata Frobrosopa: Kolla:

fasciata, Exoprosopa; Kolla; Pochazia.

fasciatus, Dysdercus; Hercothrips. fasciculatum, Trogodendron.

fasciculatus, Araecerus.

Fascista cercerisella, bionomics and control of, on Cercis canadensis in Kansas, 632, 635.

Fatty Acids, discussion of insecticicidal properties of, 592.

fausta, Rhagoletis, fax, Thysanus. febrilis, Dilophus.

fellis, Eurytoma.

Feltia experta, on flax in Peru, 511. Feltia subterranea, bait for, on cabbage and carrot in Bermuda, 138.

femoralis, Eulophus; Pedinus. femorata, Chrysobothris; Tiphia. femur-rubrum, Melanoplus.

fenestralis, Angitia. fenestratus, Scymnus.

Fenusa, bionomics of, on Viola in Massachusetts, 418.

Fenusa pusilla (pumila), on birch in New Brunswick, 348.

Fenusini, key to European genera of, 544.

Ferric Oxide (see Jeweller's Rouge). Ferrisiana (Ferrisia) virgata, on Lantana camara in India, 289; in Philippines, 317, 648; measures against, on cotton, 648; not transmitting mosaic of Musa

textilis, 317.
Ferrous Sulphate, as a corrective for spray injury to foliage, 281, 406.

ferrugineus, Dacus (Chaetodacus, Strumeta); Rhynchophorus.

ferus, Nabis.

festina, Stictocephala. festiva, Hyperaspis.

Festuca, Cecidomyiid affecting seed production in, in Britain, 315.

Festuca novae-zealandiae, pests of, in New Zealand, 334.

festucae, Contarinia.

fici, Hylesinus; Tetranychus.

Ficus spp., pests of, in Brazil, 391, 392; Hylesinus fici on, in New South Wales, 609. (See Fig.) ficus, Chrysomphalus; Pachylia.

fidoniata, Semiothisa.

Field Plots, co-ordination of experiments in, 131.

Fig, Ceratitis capitata on, in S. Africa, 18; C. capitata on, in Algerian Sahara, 616; pests of, in Brazil, 391, 392; Coleoptera on, in New South Wales, 335, 609; Icerya purchasi on, in Peru, Tetranychus fici on, in Punjab, 271; pests of, in Turkey, 288.

Fig (Fruit), juice of, in bait for Ceratitis capitata, 462.

figulilella, Ephestia.

Fiji, miscellaneous pests in, 147, 148, 169-171, 279, 389-391, 495, 496, 592; natural enemies and biological control of insects in, 170, 389, 391, 495, 496; biological control of noxious plants in, 496; pests intercepted in quarantine in, 148; beneficial insects imported into other islands from, 378, 495, 496; relands from, 378, 495, 496; restrictions on importation of bananas into U.S.A. from, 495.

Filbert (see Corylus).

Filbert Big-bud Mite (see Eriophyes avellanae).

finitima, Septis.

Finland, Diprionine sawflies in, 294, 474: Exenterus adspersus in, 294. Fiorinia proboscidaria, on Citrus in Fiji, 389; food-plants of, in Formosa, 622; possibly a variety of F. theae, 622.

Fiorinia theae, F. proboscidaria possibly a variety of, 622.

Fir, Balsam (see Abies balsamea).

Fire Ant (see Solenopsis).

Fire Ant, Small (see Wasmannia auropunctata).

Fire-blight, transmission of, by honey-bees, 112.

Firebrat (see Thermobia domestica). Fireworm, Cranberry (see Rhopobota naevana).

Firmiana platanifolia (see Sterculia).

fiscellaria, Ellopia.

Fish, Dried, Dermestes lardarius infesting, 393; (meal), in baits, 158, 255.

Fish Oil, uses of, in sprays, 112, 163, 256, 371, 415, 437, 439, 595, 639, 642, 643; dispersing agents for, 112, 113; in baits, 255, 538. Fitodendrin, 155.

Flacourtia indica, Plagiodera metallica on, in Philippines, 649.

flammea, Panolis.

flava, Hoplocampa; Sipha.

flavedana, Platynota.

flavescens, Empoasca; Phalera.

flaviceps, Carneocephala; Cryptognatha (Delphastus).

flaviconchae, Apanteles. flavicornis, Sphegigaster.

flavidus, Coccophagus.

flavifrons, Scolia. flavipalpis, Elodia.

flavipennis, Anomala.

flavipes, Apanteles; Exochomus; Reticulitermes.

flaviscutellaris, Cyrtomorpha.

flavissimus, Exenterus.

flavistriata, Decadarchis (Ereunetis).

flavopicta, Spilochalcis.

flavoscutellum, Encarsia. flavus, Aphelinus; Leptomastix.

Flax, pests of, in Formosa, 93; pests of, in India, 266, 523; Lepidoptera on, in Peru, 511; flea-beetles on, in Russian Union, 181; pests of, in U.S.A., 499, 603; relation of wireworms to, 573.

Flea-beetle, Potato (see Epitrix

cucumeris). Flea-beetle, Tobacco (see Epitrix parvula).

Fleahopper, Cotton (see Psallus seriatus).

fletcheri, Opius.

Floor Seal Varnish, effect of, in fixed nicotine spray, 160.

Florida, Acrobasis caryae on pecan in, 117, 426; pests of beans in, 439, 476, 477; pests of Citrus in, 135, 426, 474, 475, 597; miscellaneous pests in, 475, 476; Calotermes brevis in, 158; Coccinellids imported into, against Aspidiotus destructor, 472; Popillia japonica intercepted in, 118; parasite of Nezara viridula imported into other countries from, 170, 496.

Florida Red Scale (see Chrysom-phalus ficus).

floridanus, Horismenus.

Flour, pests in, and their control, 139, 327, 433, 450, 453, 455, 506, 576, 605; effect of injury to grain by Pentatomids of quality of, 574; in baits for locusts, 612; and cryolite, dusting with, against Heliothis armigera, 631; uses of, in sprays, 10, 113, 408, 595.

Flour Beetle, Confused (see Tri-bolium confusum).

Flour Mite (see Tyroglyphus farinae).

Flour Moth, Mediterranean (see Ephestia kuehniella),

Flying Fox (see Pteropus giganteus).

fockeui, Phyllocoptes.
Forest Fires, pests favoured by,

49, 86. Forests, pests of, in S. Africa, 18, 19, 30, 278, 341; Coleoptera in, in S. America, 512; pests of, in Australia, 30, 142, 237, 290, 335, 545, 609; pests of, in Britain, 180, 323, 382, 383, 628; pests of, in Burma, 75-78; pests of, in Canada, 36, 37-39, 293, 294, 308, 312, 328, 347, 348, 395, 415, 424, 431, 474, 509, 530, 531, 532, 626, 637; methods for surveys of pests of, in Quebec, 311; Lepidopterous pests of, in China, 544; pests of, in Europe, 67, 85, 86, 474; pests of, in Formosa, 103, 104; pests of, in India, 74, 75, 512; pests of, in Netherlands Indies, 105, 270, 513; pests of, in Japan, 99, 171, 172, 173, 446; Eriococcus coriaceus in, in New Zealand, 545; Gonipterus scutellatus in, in Nyasaland and S. Rhodesia, 30; pests of, in Porto Rico, 51, 52, 427; pests of, in Sakhalin, 95, 96; new weevil in, in Uganda, 341; pests of, in

U.S.A., 37, 40, 48, 49, 86, 87, 116, 118, 119, 120, 163, 164, 165, 210, 211, 212, 213, 276, 293, 294, 298, 347, 348, 357, 364, 365, 370, 371, 395, 396, 400, 412, 416, 421, 422, 424, 426, 427, 428, 432, 502, 507, 531, 537, 550, 551, 552, 563, 599, 605, 606, 607, 627, 628, 632, 635, 637, 638, 644, 645, 648; types of insect populations in, 396.

Forficula auricularia, bionomics of, in Ontario, 304; damaging bags protecting peaches in Spain, 464; in U.S.A., 537, 538; measures against, 304, 538.

Formaldehyde (Formalin), in bait for Trypetids, 91; reactions of Oscinella frit to, 573; low toxicity of solution of, to Calandra oryzae, 541.

Formica perpilosa, predacious on Lygus spp. in U.S.A., 442.

formicarius, Cylas.

Formosa, miscellaneous pests in, 45, 93, 94, 103, 104; Lepidoptera on mulberry in, 34, 93; sugar-cane pests in, 92, 94, 104, 172, 622; Coccids in, 622; beneficial insects in, 34, 92, 94, 95, 105; biological control of sugar-cane pests in, 92; pests intercepted in imported products in, 95, 103; attempted introduction of Opius fletcheri into Loochoo Islands from, 99.

formosa, Encarsia.

formosanus, Micromyzus; Sipalus. fornicatior, Xyleborus fornicatus.

fornicatus, Xyleborus.

Fornicia, parasite of Nephantis serinopa in Ceylon, **620**, **621**. Fornicia ceylonica, **621**; parasite of

Natada nararia in Ceylon, 174. forskaleana, Tortrix (Argyrotoxa).

Forsythia, injured by basic copper arsenate, 119.

Fowls, destroying Melolonthids, 24, 509.

Foxglove Aphis (see Macrosiphum solani).

fragaefolii, Capitophorus.

Fragaria spp., Capitophorus spp. experimentally surviving on, in Britain, 28. (See Strawberry.)

fragariae, Aleurodes; Anaphothrips (Neophysopus); Ancylis (see A. comptana); Capitophorus (Pentatrichopus); Tyloderma.

France, Melolonthids in, 24, 83; Phyllocoptes fockeui in, 304;

Phylloxera vitifoliae on vines in, 111; insects and virus diseases of plants in, 23, 24, 25; parasites of Cydia pomonella in, 66; attempted acclimatisation of Lebia grandis against Leptinotarsa decembineata in, 26; new spirochaete in Pieris brassicae in, 224; beneficial insects introduced into other countries from, 143, 416, 430.

Frankliniella, transmitting virus disease of tobacco and tomato in

Brazil, 141.

Frankliniella fusca, on tobacco in Florida, 476.

Frankliniella intonsa, on tomato in France, 25.

Frankliniella moultoni, thrips predacious on, on vines and fruit trees in California, 210.

Frankliniella paucispinosa, transmitting virus disease of tobacco and tomato in Argentina, 141, 608.

Frankliniella schultzei, transmitting virus disease on tomato and tobacco in South Africa, 491; spraying against, 491.

Frankliniella tritici, on rose in

U.S.A., 256.

Frankliniella vaccinii, measures against, on blueberry in Maine, 165.

fraterculus, Anastrepha.

fraterna, Lachnosterna (Phyllophaga).

fraternus, Horismenus.

Fraxinus (see Ash).

Frazinus chinensis, food-plant of Ericerus pela in Szechuan, 153.

Fraxinus lanceolata, not attacked by grasshoppers in U.S.A., 603.

frigida, Lissonota.

frit, Oscinella (Oscinis).

froggatti, Brontispa (see B. longissima); Typhlocyba.

Froghopper, Sugar-cane (see To-maspis saccharina).

frontalis, Dendroctonus.

Frontina archippivora (see Achaeto-neura).

frugiperda, Laphygma.

Fruit (Dried), pests in, and their control, 426, 488; low absorption of methyl bromide by, 444.

Fruit-flies, bibliography on, 280. Fruit-fly, Mango (see Dacus ferru-

gineus dorsalis).

Fruit-fly, Mediterranean (see Ceratitis capitata).

Fruit-fly, Melon (see Dacus cucurbitae).

Fruit-fly, Queensland (see Dacus ferrugineus tryoni).

frustralis, Loxostege.

frustrana, Rhyacionia.

frutetorum, Gilpinia (Diprion).

Fuchsia, Coccid on, in California, 223.

fuliginosum, Gymnosoma.

fuliginosus, Encyrtus.

Fuller's Earth, as an emulsifier for oils, 585.

Fuller's Rose Beetle (see *Pantomorus* godmani).

fullo, Polyphylla.

fullonia (fullonica), Othreis.

fulvicornis, Goniozus (Trissomalus). fulvipes, Hemiteles (see H. nanus).

fulvoniger, Dysdercus.

fulvovestitus, Anomocaulus.

fulvus, Cyrtorhinus.

fumiferana, Harmologa (Cacoecia).

Fumigants, principles governing use of, against pests of stored products, 84; new laboratory method for testing, 554; relating of time, concentration and toxicity in tests of, 433; review of recent work on, 483; types of equipment for applying, 40, 113, 124, 132, 276, 499, 555, 568, 569.

fumipennis, Hercothrips.

Fundella cistipennis, on leguminous plants in Porto Rico, 345.

funebrana, Cydia.

funestus, Lagochirus.

Fungi, association of termites with, 19, 338; infesting prickly-pear, 587.

Fungi, Entomogenous, 18, 19, 42, 70, 83, 90, 93, 102, 116, 152, 165, 196, 211, 229, 303, 349, 379, 497, 500, 515, 535, 570, 587, 588, 589, 635, 637; utilisation of, 20, 88, 368, 582, 583; of Mauritius, 424.

Fungi, Injurious, 2, 15, 35, 108, 159, 274, 279, 281, 283, 317, 381, 392, 485, 507, 558, 593, 597, 648; relation of insects to, 3, 23, 32, 40, 48, 53, 96, 97, 107, 112, 116, 165, 166, 167, 168, 177, 178, 179, 212, 239, 315, 316, 357, 364, 365, 495, 501, 502, 533, 550, 563, 581, 597, 606, 644, 645, 646; possibly spread by mites, 365; book on, 111.

Fungus, Brown Rot (see Sclerotinia fructicola).

fur, Ptinus.

furcifera, Phaneroptera; Sogata.

furfura, Chionaspis. furnissi, Cylindrocopturus. Furniture, pests in, 103, 158, 233, 284, 496, 543. Furs, Anthrenus vorax infesting, in U.S.A., **137.** furtifica, Camptothlipsis. Aonidiella infesting Fusarium, aurantii in Japan, 102; associated with Scolytus multistriatus on elm in W. Virginia, 116. Frankliniella; Busseola; Lachnosterna (Phyllophaga). fuscedinella, Coleophora. fuscicauda, Sarcophaga. fuscicornis, Sapromyza. fuscilabris, Ceratomegilla (Megilla). fuscipenne, Microplectron. fuscipennis, Thrips. fusciventris, Anagyrus. fuscum, Prionobrachium.

G.

Galeruca tanaceti, bionomics and control of, in Germany, 68.

Galerucella distincta, on strawberry

in Korea, 152.

futilis, Lachnosterna.

Galerucella luteola (on elm), in Rumania, 12; in U.S.A., 120, 416, 417; natural enemies and biological control of, 12, 416, 417; sprays against, 120.

Galerucella xanthomelaena (see G.

luteola).

galerucellae, Synaplomyia (Aplomy-

iopsis).

Galèrucida bicolor, bionomics and control of, on elephant yam in India, 319.

Galleria mellonella, in bee-hives in S. Africa, 611; parasite of, 611; bacilli isolated from, 583.

Gambier (see *Uncaria gambir*).

Gampsocoris pulchellus, on cacao in Malaya, 513.

Garden Centipede (see Scutigerella immaculata).

gardneri, Angitia (Dioctes).

Gas Oil, tests of, for treating water in rice-fields against Pyralid, 458; in mixture for burning locusts, 459, 460, 529.

Gasoline, in mixture for burning locusts, 459, 460, 529. (See

Petrol.)

Gasterocercodes brasiliensis, bionomics and control of, on cotton in Brazil, 623, 624. gaugleri, Cantheconidea. gei, Metallus.

INDEX.

gei, auct., Macrosiphum (see M. solanifolii).

Gelatine Grub (see Niphadolepis alianta).

Gelechia cercerisella (see Fascista).

Gelis asozanus, parasite of Apanteles liparidis in Japan, 172.

gemella, Podagrica (Nisotra). geminata, Chrysomela (Chrysolina);

Solenopsis.
gemma, Chrysocharis.
gemmatalis, Anticarsia.
Genicide (see Xanthone).
geniculatus, Engytatus.

gentilis, Melanophila.

Geocoris, predacious on Lygus spp. in U.S.A., 442.

Geocoris moderatus, measures against, damaging turf in S. Africa, 615.

Geomermis indica, parasite of Pempherulus affinis in India, **270, 522.** Geonoma, Himatidium spp. on, in

Brazil, 10.
Georgia (U.S.A.), Chalcodermus aeneus on cowpeas, etc., in, 405, 598; cowpea mosaic in, 502; miscellaneous pests in, 594, 598, 599; orchard pests in, 192, 197, 488; beneficial insects and biological control in, 192, 405, 598; Trichogramma introduced into Louisiana from, 601.

Geotrogus deserticola, parasites of,

in Algeria, 83.

Geraniol, in baits for *Popillia*, 126, 364, 596; properties of samples of, 126.

Geranium, Lepidoptera on, in Br. Isles, 146, 525; Engytatus geniculatus on, in Hawaii, 34; Heliothis armigera on, in New York, 481; reported toxicity to Popillia japonica of, 118.

germanica, Blattella.

germanus, Xyleborus (Xylosandrus). Germany, pests and diseases of crucifers in, 68, 231, 232, 234, 235, 450; miscellaneous pests in, 68, 86, 449, 488; orchard pests in, 68, 69, 70, 71, 231, 233, 339, 617; pests of timber in, 238, 234, 385; new Acarapis on honey bee in, 528; beneficial insects in, 70, 79, 234, 294; doubtful value of introduced and indigenous predators against Leptinotarsa decembineata in, 26, 70.

Geum, Metallus gei on, in Britain,

544.

giardi (giardiana), Zeuxidiplosis. Giardomyia rhododendri, measures against, on rhododendron U.S.A., 606.

gica, Balanoecus (Balanephagus).

giffardi, Dirhinus.

giffardianus, Tetrastichus. gifuensis, Macrocentrus.

gigantorhynchus, Epitrimerus.

gigas, Pantophthalmus.

Gilbert Islands, Papuana on Colocasia esculentum in, 171; Bufo marinus imported into, 496.

Gilpinia frutetorum, proposed biological control of, in N. America,

Gilpinia polytoma (on spruce), in Canada, 36, 37-39, 293, 294, 307, 308, 424, 431, 474, 530, 531, 532, 637; in Europe, 294, 474; in U.S.A., 37, 293, 294, 422, 424, 427, 531, 532, 637, 638; forms, cytology and types of parthenogenesis of, 37, 474; effect of climatic factors on, 38, 431, 531; parasites and biological control of, 36, 38, 39, 293, 294, 307, 308, 424, 427, 432, 531, 532, 637, 638; other natural enemies of, 427, 637; disease of, and its utilisation, 37, 38, 422, 427, 432, 531,

Gipsy Moth (see Lymantria dispar). glabratus, Hylastes (Hylurgops).

Gladiolus, pests of, in U.S.A., 136, 204, 214, 277; insecticides causing injury to, 277.

Gladiolus Thrips (see Taeniothrips

simplex).

glaseri, Neoaplectana.

Gleditsia triacanthos, grasshoppers feeding on, in U.S.A., 603.

Gliricidia, possible food-plant of Niphadolepis alianta in Nyasaland, 59.

globosa, Maskellia; Myiophasia.

glomeratus, Apanteles.

gloveri, Lepidosaphes (Mytilococcus). Glue, in baits for Otiorrhynchus, 252; as an emulsifier for oils,

426: as a dispersing agent for

sprays, 113.

Glycerine, tests of, in locust baits, 612; experiments with, sprays, 129, 215, 296.

Glycerol, in mixtures for maintaining humidity, 40.

Glyceryl Monooleate, as an emulsifier for oils, 350.

glycinae, Phalaenoides.

glycinivorella, Cydia (Grapholitha).

Glyphodes (see Margaronia).

Glypta, 151.

Glypta haesitator, parasite of Cydia nigricana in Britain, 311; introduction of, into Canada, 311.

Glypta rufiscutellaris, value utilisation of, against Cydia molesta in U.S.A., 87, 130.

Glyptomorpha, parasite of Zeuzera coffeae in Malaya, 513.

Gnorimoschema heliopa (on tobacco), in Fiji, 390; bionomics of, in Sumatra, 319.

Gnorimoschema operculella, in Australia, 143, 335; in Bermuda, 138; in California, 222; new parasite of, in Cyprus, 514; in Fiji, 390; in Hawaii, 650; Nyasaland, 154; on potato, 138, 154, 222, 650; on tomato, 138, 143, 335; on other plants, 138, 390; measures against, 138, 143, 222, 335.

Gnorimoschema plaesiosema, tomato in Australia, 31.

godmani, Pantomorus (Asynonychus).

Gold Coast, fruit-piercing moths in, 16, 106, 107; pests of stored cacao in, 616; miscellaneous pests in, 492, 516; beneficial insects in, 492.

Gold Mines, Micromalthus debilis in timber in, in S. Africa, 275.

Golden Codling Moth (see Cydia pomonella var. simpsoni).

Golf Courses (see Turf).

gonager, Pachymerus. gonagra, Leptoglossus.

Gonia, revision of American species of. 280.

antipodum, Eupsenella Goniozus diemenensis possibly recorded as, in Tasmania, 151.

Goniozus fulvicornis, parasite of Argyroploce illepida in India, 137; systematic position of, 137.

Goniozus indicus, sp. n., parasite of sugar-cane borers in India, 137.

Goniozus montanus, attempted introduction of, against Hyblaea puera in Burma, 77.

Gonipterus scutellatus (on Eucalyptus), distribution of, in Africa, 18, 30; in Australia, 30; utilisation of Anaphoidea nitens against,

Goniurus proteus (see Urbanus). Gooseberry, Lecanium persicae on, in New Zealand, 58; Nematus ribesii on, in Russian Union, 585;

spray schedule for, in Kentucky, 2.

Gooseberry Sawfly (see Nematus ribesii).

Gordius robustus, bionomics of, parasitising Anabrus simplex in U.S.A., 187.

Gorgopis libania, measures against, damaging turf in S. Africa, 615.

Gorse (see Ulex europaeus).

Gorse-seed Weevil (see Apion ulicis).
Goryphus, parasite of Artona catoxantha 79; parasites of, 79.

Goryphus nursei (parasite of Earias), bionomics of, in India, 183. gossypiella, Platyedra (Pectino-

phora).

gossypii, Aphis; Corythucha; Eriophyes; Phenacoccus.

gossypiperda, Bemisia (see B. tabaci).

Goulac, 191. (See Sulphite Lye.) Gracilaria syringella, on lilac in Russia, 487; effect of leaf-size on oviposition of, 487. gracilis, Lachnosterna (Phyllophaga);

Polygraphus.

Graeffea cocophaga, on coconut in Fiji, 495.

Grain Beetle, Saw-toothed (see Oryzaephilus surinamensis).

Grain Moth, Angoumois (see Sito-troga cerealella).

Grain Weevil (see Calandra gran-aria).

graminum, Toxoptera.

granaria, Calandra (Sitophilus).

granarium, Macrosiphum (see M. avenae).

grandella, Hypsipyla. grandicollis, Ips.

grandis, Anthonomus; Harmolita; Synonycha; Zadiprion (Neodiprion).

granitata, Semiothisa.

Grape Plume Moth (see Oxyptilus periscelidactylus).

Grape-berry Moth (see Polychrosis viteana).

Grapefruit, Ceratitis capitata on, in Algerian Sahara, 616; Pseudococcus comstocki on, in Brazil, 168; Aonidiella auvantii on, in California, 535; Mellesis citri on, in China, 524; pests of, in Fiji, 170, 389; fruit-piercing moths damaging, in Gold Coast, 16, 107; pests of, in Transcaucasia, 654.

Grapenuts, as laboratory food for Lachnosterna, 213.

Grapes, problem of spray residues on, 117, 425.

Graphiphora c-nigrum (see Amathes). Graphium, associated with Scolytus multistriatus in W. Virginia, 116. Graphium ulmi (see Ophiostoma).

Grapholitha (see Cydia).

grapholithae, Ephialtes (Calliephialtes); Phanerotoma.

Grass, Bent (see Agrostis).

Grass, Bermuda (see Cynodon dacty-lon).

Grass, Blue (see Poa).

Grass, Cocksfoot (see Dactylis glomerata).

Grass, Couch (see Agropyrum repens and Cynodon dactylon).

Grass, Johnson (see Sorghum hale-

pense). Grass, Para (see Panicum barbinode). Grass, Rhodes (see Chloris gayana).

Grass, Rhodes (see Chloris gayana). Grass, Timothy (see Phleum pratense).

Grass Caterpillar, Subterranean (see Oxycanus).

Grass Grub (see Odontria zealan-dica).

Grasses, pests of, in S. Africa, 614, 615; pests of, in Australia, 335, 466, 547; pests of, in Britain, 110, 314, 315, 375; Lachnosterna on, in Canada, 90; Cirphis unipuncta on, in Fiji, 495; damaging, Melolonthids France, 83; Tipulids on, in Germany, 68; termites damaging, in Kenya, 338; pests of, in New Zealand, 89, 333, 334, 541, 542; damaged by Melolontha hippocastani in Siberia, pests of, in U.S.A., 213, 214, 300, 365, 408, 506, 602; pests of, in W. Indies, 217, 373; cereal and sugar-cane pests on, 2, 3, 14, 49, 81, 93, 148, 173, 244, 300, 331, 393, 464, 560, 576, 610, 622; reservoir of mosaic disease of cereals, 14; relations of locusts and grasshoppers to, 94, 204, 205, 255, 354, 394, 485, 512, 570. (See Turf.)

Grasshopper, Differential (see Melanoplus differentialis).

Grasshopper, Lesser Migratory (see Melanoplus mexicanus).

Grasshopper, Small Plague (see Austroicetes cruciata).

Grasshoppers (see Locusts).

Greece, Parlatoria oleae possibly introduced into California from, 629.

Greenheart, British Guiana (see Nectandra vodioei).

Greenhouse Fumigation, 25, 32, 132, 145, 146, 203, 276, 375.

Greenhouse Whitefly (see Trialeurodes vaporariorum).

greeni, Dactylopius; Microbracon; Nilaparvata (see N. lugens).

gregaria, Schistocerca.

Grenada, pests of stored cacao in,

Grevillea, Bostrychid in, in New South Wales, 335.

gripusalis, Azochis.

grisescens, Lydella stabulans.

Ground-nuts, Coccids on roots of, in Egypt, 30; Epilachna vigintioctopunctata on, in Fiji, 390; locust attacking, in Formosa, 94; pests of, in Hawaii, 651; pests of, in India, 518, 523; pests of, in U.S.A., 251, 252, 428, 599; insecticides and injury to, 252, 260, 428; (stored), pests imported into Fiji in, 496.

Ground-nut Oil (see Oils, Vegetable). Gryllica melzeri (see Merocentrum). Gryllodes sigillatus, in greenhouses

in Britain, 489.

Gryllotalpa africana, in S. Africa, 615; in China, 446; bionomics of, in New South Wales, 237; in Spain, 462; measures against, damaging turf, 237, 615; characters of, 463.

Gryllotalpa australis, bionomics and control of, in New South Wales,

Gryllotalpa gryllotalpa, on potatoes in Jugoslavia, 72; in Spain, 462; review of data on, 463.

Gryllotalpa hexadactyla, in Florida,

Gryllotalpa unispina, in China, 446. gryllotalpa, Gryllotalpa.

Gryllulus assimilis, in California, 223; introduced parasite of grasshoppers developing in, in Canada, 309.

Gryllulus domesticus, in U.S.A., 88, 164; damaging clothes, 164.

Gryllulus mitratus, bait for, on vegetables in Japan, 173.

Gryllus, auct. (see Gryllulus).

Guadeloupe, original discovery of Leucoptera coffeella on coffee in,

Guam, miscellaneous insects and biological control in, 330-332; Coccinella arcuata introduced into Hawaii from, 331.

guamensis, Apanteles.

Guava, pests of, in S. Africa, 18, 227; moths damaging fruits of, in Gold Coast, 107; Spilonota holotephras on, in Guam and Fiji, 332, 495; pests of, in India, 183, 266; Anastrepha fraterculus on, in Peru, 539; Pseudococcus nipae on, in Porto Rico, 472; fumigation of harvested fruit of, against pests, 18, 499.

Guiana, British, Brassolis sophorae on coconut in, 538; establishment and value of Metagonistylum minense against Diatraea in, **4, 538;** termites imported into Barbados from, **6;** beneficial insects introduced into Porto Rico from, 470.

Guiana, Dutch, Atta spp. in, 169. Guinea Fowl (see Numidia melea-

guineense, Tetramorium.

gularis, Aphomia.

Gulls, destroying Polyphylla perversa, 509.

Gum Arabic, as a dispersing agent for sprays, 112, 113.

gurneyi, Coccophagus.

Gutta-percha (see Palaquium gutta). guttifera, Teragra.

guttulosa, Austracris.

Gymnogryllus, on soy beans in Ceylon, 594.

Gymnosoma fuliginosum, parasite of Chlorochroa ligata in Texas, 500.

Gynandropsis pentaphylla, Myzus persicae on, in Nyasaland, 154.

Gynerium saccharoides, Rhynchophorus palmarum on, in Colombia, 9. Gypsum, as a carrier for dusts, 21,

647; factors affecting toxicity of dusts used with, 301, 569.

H.

haani, Loxoblemmus.

Habrobracon (see Microbracon).

Habrocytus cerealellae, fumigation against, parasitising Sitotroga in laboratory, 193.

Habrocytus phycidis, parasite Macrocentrus cerasivorana Quebec, 626.

Habrocytus trypetae, search for hosts and egg distribution of, 490.

Habrolepis dalmanni, utilisation of, against Asterolecanium variolosum in Australia, 142.

Habrolepis rouxi, food-plant of Aonidiella aurantii affecting parasitism by, 220.

Hadronotus pedester, parasite of Eurygaster integriceps in Russia, 582.

haemorrhoidalis, Heliothrips.

haesitator, Glypta.

Hailstorms, effect of, on grass-hoppers, 394.

haimbachi, Zelleria.

Hairy Mite (see Lepidoglyphus destructor mixtus).

Halisidota caryae, tests of basic copper arsenate against, in U.S.A., 119.

halli, Lepidosaphes. halophila, Blaps.

Halotydeus destructor, in Australia, 141, 142, 236, 546, 587; factors affecting abundance and survival of, 141, 142; new Empusa infesting, on clover, 587; measures against, on crucifers, 546; 'on oats, 141.

Haltica ampelophaga, Bordeaux mixture against, on grape-vines in Spain, 458.

Haltica cyanea, on rice in Ceylon, 594.

Halticoptera, parasite of Agromyza braziliensis in Brazil, 36.

Halticoptera aenea, parasite of Mayetiola destructor and Oscinella frit in Russia, 618.

Halticus citri, measures against, on asters in Maryland, 361.

Halyomorpha reflexa, on cacao and Solanum in Nigeria, 516.

Hambletonia pseudococcina, utilisation of, against Pseudococcus brevipes in Porto Rico, 472.

hampei, Stephanoderes (Hypothene-mus).

hancocki, Anastatus antestiae.

Hapalia machaeralis, parasites and biological control of, on teak in India and Burma, 75, 76, 77.

Haptoncus luteola (see Epuraea). Haptoncus ocularis, in decayed fruits in Fiji, 390.

haradai, Rhynchites.

hargreavesi, Euplectrus; Systates. Harlequin Bug (see Murgantia histrionica).

Harmolita, effects of cultural practices on damage by species of, on wheat in U.S.A., 254.

Harmolita grandis, 254.

Harmolita tritici, in U.S.A., 254, 366; new wheat hybrids resistant to, 366.

Harmòlita vaginicola, 254.

Harmologa fumiferana, on spruce in Canada, 348, 530; factors affecting outbreaks of, on fir and spruce in U.S.A., 348, 507; race of, on pine, 507; natural enemies of, 507; measures against, 508.

Harmonia arcuata (see Coccinella). Harpobittacus, preventing establishment of Tyria jacobaeae in Aus-

tralia, 143.

Harvester Termite (see *Hodotermes*).

havilandi, Coptotermes.

Hawaii, coconut pests in, 274, 331; miscellaneous pests in, 34, 273, 650; pineapple pests in, 49, 140, 273, 274, 298, 346, 390; sugarcane pests in, 35, 49, 50, 273, 274; new thrips on taro in, 273; Micromalthus debilis in timber in, 275; insects and virus diseases of plants in, 35, 139, 140, 608, 650; beneficial insects and biological control in, 34, 35, 49, 272, 273, 274, 298, 331, 332, 390, 458, 459, 472, 496, 650, 651; pests intercepted in quarantine in, 348; danger of introduction of pests in aircraft from Guam into, 330; legislation against introduction of Trypetids into U.S.A. from, 499; pests from, intercepted in other countries, 148, 349; natural enemies of pests introduced into other countries from, 93, 298, 332, 472.

Hawaiian Beet Webworm (see Hy-

menia recurvalis).

hawaiiensis, Melittobia; Pristomerus.

Hawks, destroying noxious insects in Utah, 217.

Hawkweed (see Hieracium).

Hawthorn (see Crataegus).

Hazel (see Corylus).

Heat, uses of, against pests, 53, 59, 146, 216, 338, 381, 385, 448, 454, 506, 576, 593, 605; variations in resistance of certain insects to, 506; effect of, on germinating power of grain, 454.

hebes, Desmidophorus.

hebetor, Microbracon (Bracon).

hederae, Aspidiotus.

Hødychium coronarium, Aphids not transmitting mosaic of Musa textilis to, 317.

helichrysi, Brachycaudus (see Anuraphis padi).

heliopa, Gnorimoschema (Phthorimaea).

Heliothis, new parasite of, in Sudan,

Heliothis armigera, 543; in S. Africa, 81; in Australia, 31, 335, **466**; in Azerbaijan, **235**, **236**; in Fiji, 148, 390; in Guam, 331; in Hawaii, 496, 650, 651; in India, 289; in New Guinea Territory, 377; in Nyasaland, 62; in Porto Rico, 469; in Sudan, 386; in Sumatra, 319; in U.S.A., 3, 64, 199, 217, 257, 367, 419, 429, 430, 481, 534, 598, 599, 631, 640, 641; on Aleurites, 62; on cabbage, 199; on cotton, 3, 81, 235, 236, 257, 386, 430, 466; on geranium, 481; on Lantana camara, 289; on leguminous crops, 3, 390, 481, 534, 535, 598, 599; on maize, 3, 148, 164, 217, 331, 367, 377, 390, 419, 469, 481, 535, 640, 641, 650; susceptibility of varieties of maize to, 367; on Panicum barbinode, 148; on sorghum, 3, 466; on tobacco, 319, 390; on tomato, 31, 32, 148, 335, 390, 429, 466, 481, 534, 535, 598, 631, 640, 641, 650; bionomics of, 31, 80, 123, 235, 367, 419, 481; natural enemies and biological control of, 148, 231, 496, 534, 598, 650; Phorids bred from dead larvae of, 390; measures against, 32, 148, 199, 217, 257, 335, 390, 419, 429, 430, 466, 469, 470, 481, 534, 535, 598, 631, 640, 641.

Heliothis assulta, bionomics of, on tobacco in Sumatra, 319.

Heliothis obsoleta (see H. armigera). Heliothis virescens, bionomics and control of, on cotton and chick peas in Peru, 11, 509, 510; on tobacco in U.S.A., 11; distribution of, as a pest of cotton, 11.

haemorrhoidalis, on New York, Heliothrips rhododendrons in New 86: intercepted on Eugenia from Colombia, 86; tests of new dust on, 561; male of, 86.

Hellebore, American (see Veratrum

viride).

Hellula undalis, measures against, on cabbage in S. Carolina, 199; in Russian Union, 478; laboratory rearing of, 557; larva of, 478.

Helopeltis, records and identity of species of, on cacao and cotton in West Africa, 515, 516; on Solanum verbascifolium, 516.

Helopeltis bergrothi, food-plants of, in Nyasaland, 61, 589; identity of Capsids recorded as, on cacao and cotton in W. Africa, 515, 516. Helopeltis labaumei, on cotton in

Br. Togoland, 516.

Helopeltis sanguineus, doubtful record of, on cotton in Nigeria, 516. Helopeltis theobromae, bionomics and control of, on cacao in Malaya, 512, 513.

Helopeltis westwoodi, on cacao and Solanum in Nigeria, 516.

Helorimorpha spp., parasites Antestia in Uganda, 230.

helvolus, Metaphycus. hemerobiella, Coleophora.

Hemerocampa pseudotsugata, forests in Oregon, 49.

Hemichionaspis (see Pinnaspis). Hemimasipoda alabamae (see Win-

themia). Hemiptarsenus, parasite of Phytomyza spicata in Fiji, 170.

hemipterus, Carpophilus.

hemisphaerica, Saissetia (see coffeae).

Hemitarsonemus (see Tarsonemus). Hemiteles, parasite of Dendrolimus in Sakhalin, 95; hyperparasite of Tortrix postvittana in Tasmania,

Hemiteles areator, parasite of Apanteles glomeratus in Britain, 314. Hemiteles chrysopae, Brischke, 34. Hemiteles chrysopae, Uch. (see H.

kichijoi). Hemiteles fulvipes (see H. nanus). Hemiteles kichijoi, n. n., proposed for H. chrysopae, Uch., 34.

Hemiteles kumamotensis, parasite of Apanteles liparidis in Japan, 172. Hemiteles mesochori, parasite of Apanteles congregatus in N. Carolina, 219.

Hemiteles nanus, parasite of Apanteles spp. in Britain, 314.

Hemiteles submarginatus, parasite of Apanteles rubecula in Britain, 314.

Hemiteles tenuibasalis, sp. n., parasite of Cydia glycinivorella in Japan, 34.

Hemlock, Western (see Tsuga heterophylla).

Sawfly, Western (see Hemlock Neodiprion tsugae).

Hemp, locust attacking, in Formosa, 94; Pyrausta nubilalis on, in Russia, 488; experiments on nutritive value of, for P. nubilalis, 44, 488.

hendecasiella, Phanerotoma.

henryi, Phlugiolopsis.

lupulinus, Hepialus measures against, on Delphinium in Britain, 448.

herbarium, Catorama.

Hercothrips fasciatus, on beans in California, 223; eradication of prickly lettuce against, tests of new dust against, 561.

Hercothrips fumipennis, on cotton in Sudan, 387.

Hercothrips sudanensis, on cotton in Sudan, 387.

hesperidum, Coccus (Lecanium).

hesperus, Lygus.

Hessian Fly (see Mayetiola destructor).

heterobia, Rhabdophaga.

Heterobostrychus aequalis, in logs of Anogeissus acuminata in Burma, 78.

Heteronychus tristis. measures against, damaging turf in S. Africa, 615.

Heteropelma datanae, parasite of Datana integerrima in Missouri,

Heterotermes tenuis, attacking timber in Barbados, 6.

hexacantha, Xyloprista.

hexadactyla, Gryllotalpa.

Hexamera, liberation of, against Oncopera in Victoria, 141.

Heydenia unica, parasite of Den-droctonus frontalis in Virginia, 211.

Hibiscus, Aphis gossypii on, in Hawaii, 651; Desmidophorus hebes on, in India, 263; injured by methyl bromide, 555.

Hibiscus cannabinus, cotton pests on, in India, 517, 521; Bruchid in imported seeds of, in Formosa, 95.

Hibiscus esculentus, insects on, in India, 265, 516, 517, 521, 522, 523, 524; cotton pests on, in Sudan, 387; Platyedra gossypiella on, in U.S.A., 430; prohibition against importation of, into U.S.A. from Porto Rico, against P. gossypiella, 445.

Hibiscus ficulneus, Pempherulus affinis on, in Madras, 521, 522.

Hibiscus vosa-sinensis, Zeuzera coffeae reared on, in Java, 271.

Hibiscus tiliaceus, Xylothrips religiosus on, in Samoa, 170.

Hibiscus vitifolius, insects on, in India, 517, 521, 522.

Hickory, pests of, in U.S.A., 116, 166, 511.

Hickory Bark-beetle (see Scolytus quadrispinosus).

Hickory Shuckworm (see Enarmonia caryana).

hiemalis, Platygaster.

hieracii, Myzus.

Hieracium, Nodonota puncticollis associated with, in Pennsylvania, 192.

hilare, Acrosternum.

Himatidium, systematic position of,

Himatidium bahianum, on palms in Brazil, 10.

Himatidium bondari, on palms in Brazil, 10.

Himatidium neivai, sp. n., measures against, on coconut in Brazil, 10, 468; susceptibility of varieties of coconut to, 468.

hippocastani, Melolontha.

Hippodamia convergens, predacious on noxious insects in U.S.A., 251, 534.

Hippotion celevio, measures against, on Colocasia and sweet potato in Fiji, 170; Bufo marinus controlling, on sweet potato in New Guinea Territory, **377.** *Hirsutella subulata*, infesting *Cydia*

pomonella in U.S.A., 535.

hirsutus, Phenacoccus.

hirta, Epicometis; Scolia.

hirticornis, Epicauta.

hirticula, Lachnosterna (Phyllophaga).

hispaniolae, Termes. histrionica, Murgantia; Zeugenia. Hive Beetle, Small (see Aethina tumida).

Hodotermes (Harvester Termites), in S. Africa, 19, 338; measures against, 199.

Hodotermes mossambicus, damaging grasses in Kenya, 338.

Holcocera pulverea, predacious on Laccifer lacca in India, 263, 264, 265, 523; parasites and biological control of, 264, 265, 523.

Holcus lanatus, Contarinia affecting seed production in, in Britain, 315.

Holland, parasites of Plutella maculipennis in, 65; Psocid in spinach seed imported into California from, 416.

Holly (see Ilex).

Hollyhock (Althaea rosea), pests of, in India, 271, 516, 517, 521, 523.

holotephras, Spilonota. Holotrichia (see Lachnosterna).

Homona coffearia (Tea Tortrix), parasites and biological control of, in Ceylon, 173, 174.

Homona phanaea, on cotton in Philippines, 649.

homonae, Macrocentrus.

Honduras, British, Schistocerca paranensis in, 498: Cordyceps infesting insects in, 152.

Honesty (see Lunaria).

Honey, as a larval attractant in sprays against Cydia pomonella, 129; in bait for Dacus cucurbitae, 99; used for feeding insects, 148, 266, 344, 394, 645. Honey Locust Tree (see *Gleditsia*

triacanthos). Honeysuckle, Argyroploce urticana on, in Britain, 283.

Hop-damson Aphis (see Phorodon humuli).

Hoplocampa, parasitised by Polyodaspis ruficornis, 71.

Hoplocampa flava (Plum Sawfly), in Britain, 108, 286; in Palestine, 622; measures against, 286, 622.

Hoplocampa testudinea (Apple Sawfly), measures against, in Britain, 108, 285.

Hoplophorion pertusa (Hoplophora porosa), measures against cacao, etc., in Brazil, 392.

Hoplorrhinus testaceus, sp. n., on Cocos coronata in Brazil, 468.

Hoplorrhinus unicolor, sp. n., on Cocos coronata in Brazil, 468.

Hops, pests of, in Britain, 282; Tetranychus telarius on, in U.S.A.,

Horismenus floridanus, parasite of Apanteles congregatus in N. Carolina, 219.

Horismenus fraternus, parasite of Parornix prunivorella in Kansas,

Horistonotus uhleri, bionomics and control of, in U.S.A., 253, 602.

Hormius, 176.

Horse-radish, wireworms on, in New Jersey, 122.

Horse Nettle (see Solanum carolin-

horticola, Anomala (Phyllopertha). House-fly (see Musca domestica). howardi, Pleurotropis; Scelio. . howitti, Aphodius.

huidobrensis, Agromyza (Liriomyza). hullensis, Exenterus.

humboldti, Castnia atymnius.

humeralis, Carpophilus. Humidity, relations of locusts and grasshoppers to, 141, 157, 341-343, 484-486, 520, 583; relations of other insects to, 18, 26, 34, 35, 56, 63, 73, 102, 104, 118, 123, 139, 155, 156, 160, 172, 180, 181, 185, 186, 234, 245, 254, 304, 315, 329, 332, 361, 420, 427, 436, 452, 454, 455, 478, 487, 515, 547, 551, 566, 567, 571, 574, 575, 577, 579, 580, 581, 582, 591, 595, 620, 621, 638, 650; mixtures for maintaining, 40.

humilis, Iridomyrmex; Opius.

humuli, Phorodon.

Hungary, new Acarapis on honey bee in, 528; Contarinia medicaginis on lucerne in, 12; beneficial insects in, 12, 71, 294.

hunteri, Sarcophaga.

hyalina, Cinara (see C. pinicola). hyalinata, Diaphania.

hyalinus, Perilampus.

Hyalopterus arundinis (pruni), on peach in Jugoslavia, 72.

Hyalospila ptychis, bionomics and control of, on coconut in Brazil, 9. Hyblaea puera, parasites and biological control of, on teak in

India and Burma, **75, 76, 77.** Hybosorus illigeri, measures against, damaging turf in S. Africa, 615. hydei, Drosophila.

Hydnocarpus, Xyleborus testaceus on in Fiji, 169; pests of, in Uganda,

Hydnocarpus wightiana, Zeuzera coffeae on, in Malaya, 513.

Hydrangea, not attacked by Brotolomia meticulosa, 525.

Hydrochloric Acid, tests of, in baits

for locusts, 612.

Hydrocyanic Acid, 433, 434; tent fumigation with, against Coccids, 102, 280, 336, 337, 350, 398, 399, 410, 437, 463, 553, 564; factors influencing effectiveness of, 336, 399, 410, 553; evaluation of results of using, 463; studies on results of using, 463; studies on resistance to, in races of Aonidiella aurantii, 339, 398, 410, 535; unsatisfactory against Pseudococcus adonidum and Icerya purchasi, 280; as a greenhouse fumigant, 32, 121, 145; against pests in stored products, etc., 19, 47, 362, 429, 610; comparison

of methods of fumigation against Calandra granaria with, 457; effect of sublethal doses of, on Tribolium confusum, 455; against Habrocytus cerealella in parasite laboratories, 193. (See Calcium, Potassium and Sodium Cyanides.)

Hydrogen Sulphide, 433.

hydrophiloides, Dactylosternum. Hylastes, Ophiostoma ulmi asso-

ciated with, in New York, 644.

Hylastes ater, effect of felling on incidence of, on conifers in Britain, 382, 383.

Hylastes glabratus, bionomics of, on

spruce in Sakhalin, 96.

Hylastes obscurus, on clover in Montana, 537.

Hylastes rufipes, treatment of elm logs against, in U.S.A., 427.

Hylastinus (see Hylastes).

Hylemyia antiqua (Onion Fly), measures against, in Britain, 374; in U.S.A., 131; plot method for testing timing of

sprays against, 131.

Hylemyia brassicae (on crucifers), measures against, in Britain, 109, 110, 375, 383—385; in Germany, 232, 235; in New Jersey, 123; in Russian Union, 110, 478; bionomics of, 232, 478; toxicity of dichloroethyl ether to, 123.

Hylemyia cilicrura, outbreak of, on beans in Br. Columbia, 330; in Transcaucasia, 159; bionomics and control of, damaging vegetables and tree seedlings in U.S.A., 158, 348.

Hylemyia jacobaeae, establishment of, on ragwort in New Zealand,

334.

Hylemyia nidicola, attacking grasshoppers in Canada, 308.

Hylesinus fici, measures against, on Ficus spp. in New South Wales, 609.

Hylobius abietis, effect of felling on incidence of, on conifers in Britain, 323, 382, 383, 593; on pine in Latvia, 86; measures against, 86, 323, 593.

Hylobius piceus, on spruce ir Quebec, 37.

Hylotrupes, characters and species of, 497.

Hylotrupes amethystinus, 497.

Hylotrupes bajulus (in building timbers), in Argentina, 497; in France, 497; in Germany, 234,

385; in Latvia, 86; in Sweden, 324; measures against, 385.

Hylotrupes bifasciatus, 497.

Hylurgopinus (see Hylastes). Hylurgops (see Hylastes).

Hymenaea courbaril, susceptibility of timber of, to Calotermes approximatus in Bermuda, 138.

Hymenia recurvalis (fascialis), bionomics and control of, in U.S.A., 46, 152; laboratory rearing of, 557.

Hyoscyamus, factors affecting transmission of virus diseases of, by Aphids, 27, 179.

Hypena abyssinialis (ignotalis), on Lantana camara in India, 289.

Hypera, Coccinellid predacious on, on lucerne in Uzbekistan, 486.

Hypera brunneipennis, distribution of, in U.S.A., 403, 628, 629; question of country of origin of, 403, 629; precautions against spread of, 629.

Hypera postica (see H. variabilis). Hypera rumicis, new Necremnus parasitic on, in Oregon, **569**.

Hypera variabilis, question of distribution of, in Mediterranean Basin, 403; on lucerne in U.S.A., 222, 401, 402, 409, 410, 569, 632; bionomics and apparent climatic limitations of, 401, 402; parasites and biological control of, 401, 409, 569; status of name, 401 (note).

Hyperaspis billoti, predacious on Coccids in Trinidad, 471; introduced into Porto Rico, 471.

Hyperaspis festiva var. apicalis, predacious on Pseudococcus nipae in Porto Rico, 52.

hyperici, Agrilus.

Hypericum perforatum, attempted introduction of insects against, in Australia, 143.

Hyperstylus lineatus, sp. n., on apple in Punjab, 341.

Hyphantria cunea, tests of basic copper arsenate against, 119.

Hypolixus truncatulus, parasites of, on Amarantus in Madras, 522; Spathius vulnificus reared on, 269. Hypomeces squamosus, on cacao in

Malaya, **513.**

Hypomolyx piceus (see Hylobius).

Hyponomeuta padellus, tests of tar distillates against, on plum in Germany, 231.

Hypophloeus parallelus, on Pinus resinosa in Connecticut, 164.

Hypopteromalus apantelophagus, hosts of, in Sakhalin and Japan, 95, 172.

Hypopteromalus tabacum, bionomics of, in N. Carolina, 219.

Hyposidra talaca, on teak in Burma,

Hyposoter exiguae, parasite of Lepidoptera in Hawaii, 35, 650.

Hypothenemus (see Stephanoderes). Hypsipyla grandella, on Cedrela and mahogany in Porto Rico, 427.

Hypsipyla robusta, factors affecting infestation of mahogany by, in India, **75**.

Hyptis atrorubens, Larra americana associated with, in Porto Rico, 553.

I.

icasia, Ecpantheria.

Icerya, new parasite of, on mandarin in Argentina, 344.

Icerya aegyptiaca, Coccinellid predacious on, on Schleichera trijuga in India, 263.

Icerya purchasi, in S. Africa, 320; in S. America, **511**, **512**; in S. in California, Australia, 280; in Cyprus, 544; 280: Dominican Republic, 498; Nyasaland, 62; on Citrus, 280, 320, 498, 511, 544; on other plants, 62, 498, 511, 512; natural enemies and biological control of, 62, 280, 320, 498, 511, 512, 544;

other measures against, 280.

Icerya seychellarum, on Citrus in Fiji, 389; food-plants and biological control of, in Seychelles,

iceryae, Cryptochetum. iceryoides, Phenococcus.

Ichneumon instigator (see Pimpla). Ichneumon irritator, parasite of Cirphis unipuncta in Japan, 98.

Ichneumon manifestator (see Ephial-

ichthyurae, Telenomus.

idaei, Aphis.

Idaho, Nematode parasite of Anabrus simplex in, 187; new mite on clover in, 530; Crioceris asparagi Eutettix tenellus and in, 48; diseases in beet and tomato in, 134, 533; pea Bruchid in, 403; new weevil on pine in, Psylla pyricola in, 48, 371 425; Spilochalcis leptis in, 190.

Idiocerus, on mango in Ceylon, 594.

igneoides, Spilochalcis.

ignotalis, Hypena (see H. abyssinia-

Ilex (Holly), pests of, in U.S.A., 595, 606.

Ilex aguifolium (English Holly), Phytomyza ilicis on, in Columbia and Br. Isles, 417, 489, 490.

ilicicola, Phytomyza. ilicis, Phytomyza.

illepida, Argyroploce (Cryptophlebia). Illiberis pruni, on pear and flowering apple in China, 446.

illigeri, Hybosorus.

Illinoia pisi (see Macrosiphum ono-

brychis).

Illinois, cereal pests in, 48, 245; Conotrachelus nenuphar on peach in, 196; directions for spraying fruit in, 224; hyperpararasite of grasshoppers in, 401.

imaginis, Thrips.

immaculata, Cyclocephala; Scutigevella.

imperata, Boarmia (Ascotis) (see B. selenaria).

impressa, Entyposis. impressithorax, Bruchus. improportionalis, Amphicordus. inaequalis, Coelophora. Inareolata (see Angitia). incarnatus, Tryphetus. incertus, Zatropis. incisus, Dacus ferrugineus. inconsequens, Taeniothrips. inconspicua, Bemisia. incultana, Polychrosis.

Incurvaria rubiella (on Rubus spp.), in Britain, 283; bionomics and control of, in Canada, 307.

Indarbela, on forest trees in Burma, 76, 78; parasitised by Nemeritis tectonae, 76.

Indarbela quadrinotata, 76.

India, cotton pests in, 183, 265, 268, 270, 515, 523; forest pests in, 74, 75, 77, 382, 514; pests of fruit trees in, 75, 183, 260, 270, 272, 523; cultivation and pests of lac and its host-trees in, 263, 264, 523; miscellaneous pests in, 74, 184, 268, 270, 523; sugarcane pests in, 137, 183, 184, 268; tobacco pests in, 20, 183, 266, Galerucida bicolor on elephant yam in, 319; insects and virus diseases in, 75, 183, 382: (Delhi), Aphids of, 266; Dacus ferrugineus and its varieties in, 183, 524; new Jassids in, 265,

266: locusts in, 269, 323, 341-343, 519, 520, 523; new Tenebrionid infesting beehives in, 343; Thysanoptera of, 480; classification and new species of weevils in, 79, 265; beneficial insects and biological control of insect pests in, 74, 75, 77, 137, 175, 183, 184, 263, 264, 265, 266, 267, 268, 270, 480, 514, 523, 524, 620; Lantana insects in, 74, 289; suggested introduction of Teleonemia scrupulosa into, quarantine restrictions in, 290, Cryptorrhynchus mangiferae intercepted in Fiji from, 148. (See also under the various Provinces.) Moth (see interpunctella).

Indian Meal

Indiana, Cydia spp. on apples in, 48, 193, 206, 351, 535; fungus infesting C. pomonella in, 535; cowpea mosaic in, 502.

indianus, Olenecamptus. indica, Apis; Geomermis.

indicum, Calosoma maderae; Syntomosphyrum.

indicus, Euplectrus; Goniozus: Jassus; Paratetranychus.

Indigofera cordifolia, preferred by Schistocerca gregaria, 519.

indistincta, Apate.

Indo-China, Coccinellid introduced Formosa from, against Oregma lanigera, 93.

inermis, Stictocephala. inferens, Sesamia.

infixaria, Boarmia (Serraca).

inflata, Pimpla.

inflatus, Corymbites (Ludius).

infuscatellus, Chilo. injuriosus, Tetrastichus.

innuba, Anomala.

Inocarpus, Xyleborus aplanatide-clivis on, in Fiji, 170.

Inocarpus edulis, Argyroploce ille-

pida on, in Fiji, 496.

Insecticides, tests of organic compounds as potential, 407; methods of testing, 195, 311, 356, 415, 557, 591; rearing of insects for tests of, 199, 261, 311, 407, 591; relating of time, concentration and toxicity in tests of, 433; factors affecting penetration of, into insect cuticle, 239-241; review of recent work on, 483 patents relating to, in U.S.A., 592.

Insects, study of outbreaks of, in Europe, 11; research on preven-

tion of outbreaks of, 355; importance of racial segregation in, 548; problems of spread of, by commerce, 543; aerial distribution of, 262; book on embryology of, 328; microbiology of, 285 review of recent work on control of, 40: factors affecting permeability of cuticle of, 239-241; book on entomophagous, 431; micromortar for triturating, 349. insidiosus, Orius (Triphleps); Psyl-

ledontus. insignis, Axionicus; Macropsis.

insolitus, Phenacoccus. instigator, Pimpla (Ichneumon).

insulana, Earias.

insulare, Melittomma. insularis. Blissus.

integerrima, Datana.

integra, Lineodes. integriceps, Eurygaster.

intermedia, Tiphia.

intermedium, Trichogramma. intermedius, Rhinotermes.

intermixta, Plusia (Phytometra). internodia, Ripersia.

interpunctella, Plodia.

intonsa, Frankliniella.

inversa, Lachnosterna (Phyllophaga). Iowa, grasshoppers in, 326, 327; potato pests in, 325; miscellaneous pests in, 326, 406; Spilochal-

cis leptis in, 190.

Ipomoea spp., Engytatus geniculatus on, in Hawaii, 650; Oncocephala tuberculata on, in India, 270; Cylas formicarius on, in Louisiana, 602; Bemisia tabaci on, in Sudan, 387. (See Sweet Potato.)

Ips, staining associated with, in Pinus ponderosa in California, 357; on spruce in Canada, 532; effect on, of logging killed trees, 357.

Ips bidentatus, effect of felling on incidence of, on conifers in Britain, 382.

Ips calligraphus, on pines in Connecticut, 164.

Ips emarginatus, staining associated with, on Pinus ponderosa California, 357.

Ips grandicollis, on pines in Connecticut, 164.

Ips oregoni, on pines in U.S.A., 49. Ips pini, on pines in U.S.A., 164. 396; nature of outbreaks of, 396.

Ips ponderosae, on Pinus ponderosa in U.S.A., 49.

Iraq, summary of plant quarantines in, 499.

Iridomyrmex humilis, predacious on termites in Louisiana, 499; outbreak and control of, in Melbourne, 332.

Iridomyrmex myrmecodiae, attacking beneficial ants in Solomon Is., 325.

Iris, Macronoctua onusta on, in New York, 644.

Iroko (see *Chlorophora excelsa*). Iron Pyrites, test of residue from, against *Calandra oryzae*, **541**.

Ironbark, Narrow-leaved (see Eucalyptus crebra).

irregularis, Chrysopa.

irritator, Ichneumon (Melanichneumon).

Isaniris ater, bionomics of, in Nyasaland, 62.

Isaria fumosorosea (see Spicaria). Ischiodon (see Xanthogramma).

Ischnaspis longirostris, on mahogany in Porto Rico, 52; Coccinellids introduced into Seychelles against, on coconut, 324.

Iso-eugenol, as an attractant for Dacus spp., 90.

Isodromus, parasite of Pseudococcus comstocki in Palestine, 339.

Italian Millet (see Setaria italica). italicus, Calliptamus.

Italy, Epidiaspis leperii in, 155; natural enemies of Cleonus mendicus in, 316; new Acarapis on honey bee in, 528; Parlatoria oleae possibly introduced into

California from, 629.

Itonida citrulli, observations on, on watermelon in Arizona, 596.

Itoplectis, status and synonymy of, 220. (See Pimpla.)

Iva axillaris, new weevil on, in Saskatchewan, 530.

Ivi Nut (see *Inocarpus edulis*). Ivy, mites on, in California, **223, 586.**

J.

jacaena, Euribia.
jacksoni, Pulvinaria.
jacobaeae, Hylemyia; Tyria.
Jak Fruit (see Artocarpus integrifolia).

Jak Fruit-borer (see Margaronia

caesalis).

Jamaica, miscellaneous pests in, 293, 445; biological control of Cosmopolites sordidus in, 445. jamaicensis, Lachnosterna.

janata, Achaea.

Japan, pests of cereals and grasses in, 93, 97, 98, 99, 100, 101, 152, 173; Nephotettix and virus disease of rice in, 272; pests and fungus disease of chestnut in, 97, 99; pests of Citrus in, 98, 102, 172; forest pests in, 40, 99, 171, 172, 173, 294, 445, 480; Margaronia spp. on mulberry in, 34, 101, 102; orchard pests in, 45, 96, 99, 372, 622; Cydia glycinivorella on soy beans in, 33, 34; pests of stored products in, 97, 102, 172, 173, 508; vegetable pests in, 100, 171, 173; natural enemies of Bombyx mori in 88, 98, 172; experimental utilisation of fungus against Sturmia sericariae in, 88; new Braconid parasite of Dictyoploca japonica in, 480; beneficial insects in, 33, 34, 40, 98, 101, 102, 172, 294, 372, 425, 445, 622; failure to establish Scutellista cyanea in, 172; beneficial insects imported into other countries from, 87, 295, 331, 373, 425.

Japanese Beetle (see Popillia japonica).

Japanese Black Pine (see Pinus thunbergi).

Japanese Mandated Islands, Hispid on coconut in, 331.

Japanese Table Pine (see Pinus densiflora umbraculifera).

japonica, Dictyoploca; Popillia; Theretra.

japonicus, Coccophagus; Spilocryptus.

Jarring, to dislodge weevils from fruit trees, 41, 42, 197; to destroy Saperda in seedling apples, 243.

Jassids, study of feeding habits of, 548.

Jassus indicus, suspected vector of spike disease of sandal in India, 75. jatropha, Cryphalus.

javana, Pheidole.

javanica, Tessaratoma.

javanus, Plaesius.

javensis, Marietta.

Jeffrey Pine (see Pinus jeffreyi).

jesuita, Bostrychopsis.

Jeweller's Rouge, in dust formula against termites, 610.

Jeyes' Fluid, and orthodichlorobenzene, in emulsion against Tipulid and Bibionid larvae, **524**, **527**. jezoensis, Trogus lutorius.
Jimson Weed (see Datura stramonium).
johannes, Parasa.
Johnson Grass (see Sorghum halepense).
johnsoni, Ooencyrtus.
johnstoni, Elasmus.
juglandicola, Chromaphis.
juglandis, Habrobracon (see Microbracon hebetor).
Juglans (see Walnut).
Juglans nigra (Black Walnut), pests

of, in U.S.A., 119, 166.
Jugoslavia, miscellaneous pests in,

71, 72.
julidicolus, Caloglyphus.

Julus moreleti, bait for, on potato in Bermuda, 138.

jungi, Austroicetes (see A. cruciata).
Juniper, Chinese (see Juniperus chinensis).

juniperina, Contarinia.

Juniperus bermudiana, termite in timber of, in Bermuda, 138.

Juniperus chinensis, Contarinia juniperina not attacking varieties of, in Missouri, **432.**

Juniperus procumbens, Eriophyes on, in Japan, 99.

Juniperus scopulorum, Contarinia juniperina on, in U.S.A., 432.

Juniperus virginiana, pests of, in U.S.A., 348, 482, 607, 632.

junodi, Acanthopsyche.

Jute (see Corchorus capsularis and C. olitorius).

K.

K-58, 350.

Kainit, soil treatment with, against cutworms, 450.

kalmi, Lygaeus. Kalo, **633.**

Kalotermes (see Calotermes).

kamiyae, Ceraphron (Calliceras).
Kansas, miscellaneous pests in, 202, 367, 505; orchard pests in, 190, 632, 633-635; pests of shade trees in, 433, 632, 635; Ancylis comptana on strawberry in, 633; insects in stored cereals in, 506;

beneficial insects in, 190, 191, 505.

Kaolin, as carrier for dusts, 92, 290, 335.

Kapok, Lamiid on, in Brazil, 292; Zeuzera coffeae on, in Malaya, 513; Dysdercus spp. on, in Sudan, 387.

Karaya Gum, as an adhesive for nicotine sulphate, 413; in baits for Otiorrhynchus, 252.

Kearfottia, associated with Coccids on bamboo in Porto Rico, 472.

kellyi, Sarcophaga.

Kentucky, Contarinia juniperina in, 433; pests of fruit crops in, 2, 362, 554; Lamellicorn beetles and their natural enemies in, 298, 299; pests of tobacco in, 363, 482; reservoir of tobacco streak in, 166.

Kenya Colony, termites in, 338, 590; parasites of noxious insects in, 80, 177, 385, 514; introduction of parasites of Saissetia oleae into

California from, 385. kenyaensis, Microterys.

Kerosene, uses of, in treating timber against Coleoptera, 147, 393, 448; tests of, for treating water in rice-fields against Pyralid, 458; atomised sprays of insecticides in, 56, 434; as a solvent, 78, 110, 448.

Kerosene Emulsion (see Oil Emul-

sions).

kerri, Empoasca. kiambuensis, Euplectromorpha. kichijoi, Hemiteles (Microtorus).

kirkpatricki, Microbracon. klugi, Campsomeris; Tritneptis.

Knapweed (see Centaurea).

Kolla fasciata, favoured by lawn

sprinkling in Porto Rico, 217.
Korea, Calandra oryzae in stored rice in, 92; Cydia spp. in, 33, 372; miscellaneous pests in, 103, 152, 171; beneficial insects in, 33, 103, 294, 372; beneficial insects imported into U.S.A. from, 87,

295, 373, 431.

Kudzu (see Pueraria hirsuta). kuehniella, Ephestia. kumamotensis, Hemiteles.

kurosawai, Apanteles.

Kurrajong (see Brachychiton populneus).

Kutane, 117.

kuwanae, Chilocorus.

L.

labaumei, Helopeltis.
labdacus, Spathius.
Labia minor, in Montana, 537.
Labrorychus, parasite of Tortrix
cerasivorana in Quebec, 626
laburni, Aphis.

Laccifer lacca (Lac Insect), cultivation and natural enemies of, in Siam, India and Ceylon, 94, 263, 264, 265, 523, 594.

Lachnosterna, in Canada, 89, 90; in U.S.A., 213, 258, 298, 400, 421, 432, 643, 644; in forest nurseries, 400, 421, 432; on strawberry and raspberry, 258, 259, 643; bionomics of, 89, 298; natural enemies of, 90, 299; measures against, 258, 259, 400, 644.

Lachnosterna anxia, in Canada, 89; in New York, 259, 421, 432, 643; bionomics of, 421, 432, 643.

Lachnosterna crenulata, in New York, 421.

Lachnosterna crinita, bionomics of, in Texas, 213, 214.

Lachnosterna drakei, in Canada, 89. Lachnosterna ephilida, in Kentucky, 299.

Lachnosterna fraterna, in New York, 421.

Lachnosterna fusca, in Canada, 89; in New York, 259, 421, 432, 643; bionomics of, 421, 432, 643.

Lachnosterna futilis, in Canada, 89. Lachnosterna gracilis, in New York, 421.

Lachnosterna hirticula, in Kentucky, 298.

Lachnosterna inversa, in Kentucky, 299.

Lachnosterna jamaicensis, on sugarcane in Jamaica, 293.

Lachnosterna lanceolata, bionomics of, in U.S.A., 213, 214.

Lachnosterna morosa, repellent sprays against, in Korea, 171.

Lachnosterna rugosa, in Canada, 89; in New York, 259, 643.

Lachnosterna smithi (on sugar-cane), in Barbados, 6, 7; importation of parasites into Mauritius against, 84, 609, 610; Bufo marinus destroying, 7, 609; damage by Euternes costalis attributed to, 6.

Lachnosterna tristis, bionomics of, in New York, 421, 432.

Lachnus sabinae (see Cinara).

Lactic Acid, in baits for Cydia pomonella, 414.

lactis, Carpoglyphus.

Lactuca scariola, eradication of, against Hercothrips fasciatus in California, 223.

Ladoria desarmata, introduced into Porto Rico against Asterolecanium spp., 471. Laelius voracis, sp. n., bionomics of, parasitising Anthrenus vorax in U.S.A., 137.

Laemophloeus, in stored wheat and cacao, 380, 616.

Laemophloeus minutus, fumigation of sacked grain against, in U.S.A., 209.

laetivirens, Myllocerus.

Laevicephalus striatus, relation of, to mosaic diseases of cereals in Russian Union, 13, 15, 617, 618; bionomics of, 617, 618.

laevicollis, Leucocera. laevigatum, Lygidolon. laevigatus, Orius (Triphleps).

laeviventris, Palmodes.

Laevodeguelin, 135.

Lagenaria siceraria, new weevil on, in Philippines, 649.

Lagerstroemia, Ceratitis capitata on, in S. Africa, 18.

Lagochirus funestus, question of desirability of introduction of, into S. Africa against *Opuntia*, 18.

lagopyga, Rhinoscapha.

Lagria villosa, on strawberry in S. Africa, 19.

lahillei, Chrysomphalus.

Lamellicorns, key to larvae of British species of, 480.

Lamium amplexicaule, Penthaleus major on, in N. S. Wales, 236.

Lampronia rubiella (see Incurvaria). lanceolata, Lachnosterna (Phyllophaga).

lancigera, Cletomorpha; Oregma (Ceratovacuna).

lanigerum, Eriosoma (Schizoneura).

Lantana, insects on, in India, 74,
289; suggested introduction of insects into India against, 289;

Zethus rufinodus on, on Mona Island, 473; establishment of Teleonemia scrupulosa on, in Queensland, 467; distribution of T. scrupulosa against, in Pacific Islands, 496.

lantanae, Agromyza (Ophiomyia); Teleonemia (see T. scrupulosa).

Laphygma eridania, measures against, on potato in Bermuda, 138; parasite of, in Cuba, 538; tests of insecticides on, 39, 242, 356, 414, 481; laboratory rearing of, 557.

Laphygma exempta, in Africa, 154, 514, 589; in Hawaii, 49, 332; on grasses and sugar-cane, 49; on maize, 589; measures against,

on Pennisetum, 154; parasites of, 49, 332, 514, 589.

Laphygma exigua, on vegetables in Hawaii, 650, 651; in India, 20, 447; new parasite of, in Nyasaland, 514; on cotton in Queensland, 466; on jute seedlings, 447; on tobacco, 20, 514; measures against, 20.

Laphygma frugiperda, in S. America, 54, 511; in Porto Rico, 469; in U.S.A., 164, 476, 482; on cotton, 54; on flax, 511; on maize, 54, 164, 469, 476, 482; measures against, 54, 469, 470, 476.

laphygmae, Euplectrus.

Larch (Larix), effect of felling of, on forest insects in Britain, 382; pests of, in Canada, 36, 37, 532; bark-beetles of, in Japan and Sakhalin, 480; other insects on, in Japan, 171, 446; pests of, in U.S.A., 348, 422.

Larch Sawfly (see Pristiphora erich-

soni).

lardarius, Dermestes.

laricicola, Cinara.

laricinus, Exenterus (see E. adspersus).

laricis, Anoplonyx; Coelonotus (Aphidius).

Larinus, parasitised by Rondania cucullata in Italy, 316.

Lariophagus distinguendus, parasite of Lasioderma serricorne in Fiji, 391.

Larix (see Larch).

Larra americana, parasite of Scapteriscus vicinus in Brazil, 553; factors affecting transport and establishment of, in Porto Rico, 553.

Lasioderma serricorne, on cotton in Peru, 511; measures against, in stored tobacco, 362, 390, 493, 610; in other stored products, 391, 493, 616; bionomics of, 493; water vapour affecting survival of, at reduced pressure, 302; parasites of, 391, 616.

Laspeyresia (see Cydia).

Laspeyresia caryana (see Enarmonia).

laspeyresiae, Chaetolixophaga.

lata, Phytolyma.

lataniae, Aspidiotus.

lateralis, Astycus.

laticollis, Bryocoropsis.

latiferreana, Cydia (Melissopus).

latipennis, Adapsilia.

latipes, Mocis (Pelamia).

latistriga, Parasa.

latus, Tarsonemus (Hemitarsonemus). Latvia, forest and timber pests in, 85, 86; parasites of Neodiprion sertifer in, 86.

Launea asplenifolia, Bemisia tabaci transmitting virus disease to tobacco from, in India, 183.

Lauryl Rhodanate, 108. (See Thiocyanates.)

Lawns (see Turf).

Lead, Coleoptera boring in, 335, 845; problems of residues of, on sprayed fruit, 117, 351, 411,

412, 634.

Lead Arsenate, against cereal pests, 148, 464, 534; against Citrus pests, 98, 389; against Sexava on coconut, 91, 92; against cotton pests, 10, 11, 430, 466, 624; against pests of crucifers, 468, 547; against pests of forest and shade trees, 119, 120, 395, 415, 439, 595, 607, 608, 632, 644; against pests of leguminous crops, 39, 252, 534, 600; against pests of nut trees, 116, 133, 134, 167, 426, 561; against orchard pests, 1, 2, 22, 23, 42, 45, 54, 55, 67, 72, 127, 128, 129, 162, 164, 171, 190, 192, 196, 197, 198, 206, 238, 239, 243, 279, 281, 285, 330, 351, 371, 406, 412, 416, 465, 550, 559, 632, 633, 634, 640, 641, 643; against pests of small fruits, 362, 633; against pests of ornamental plants, 146, 448, 526, 595, 606; against pests of solanaceous plants, 20, 21, 31, 32, 39, 103, 143, 146, 163, 292, 318, 335, 365, 390, 492, 534; against vine pests, against pests of 117, 333; other plants, 106, 447, 513, 649; against pests in soil and turf, 42, 90, 259, 262, 408, 527, 542, 548, 614, 615, 643; tests of toxicity of, to insects, 54, 55, 91, 92, 119, 337, 356, 481, 550, 561, 600; effect of particle size on toxicity of, 119, 550; methods of applying, from aircraft, 257, 412; in bait-sprays, 54, 55, 337, 492; in baits, 173, 362, 534, 612; dusting with, 10, 31, 32, 42, 91, 92, 143, 148, 252, 318, 335, 430, 466, 468, 542, 547, 548, 561; carriers for, 92, 252, 318, 335, 542; oil-impregnated dusts of, 196, 412; plants dipped in, 31, 547; formulae for, in sprays, 10, 20, 31, 32, 39, 42, 72, 90, 116, 119, 120, 133, 134.

148, 163, 167, 192, 196, 238, 257, 279, 292, 335, 365, 371, 406, 408, 415, 416, 439, 513, 547, 548, 561, 595, 606, 607, 624, 632, 633, 634, 640, 643, 644; adhesives and spreaders for, 10, 21, 32, 117, 134, 162, 163, 238, 281, 371, 406, 408, 415, 439, 595, 607, 633, 634, 643; factors affecting deposits and retention of sprays of, 206, 281, 371, 513; and aluminium sul-lime, 10, 11, 196, 406, 412, 634; and fime-sulphur, 162, 281, 285, 634, 644; reactions between mixtures of lime-sulphur and, 281; and manganese sulphate, 162; and molasses, plants swabbed with, 148, 466; nicotine, 335, 415, 606, 634; and oils, 67, 127, 129, 133, 134, 279, 281, 381, 412, 606, 632, 633, 634; inverted spray mixtures of oils and, 163, 371, 641; and rotenone, 640; Selocide incompatible with, 559; and sulphur, 10, 634; and zinc sulphate, 134, 196, 633; reactions of, with salts in soil and spray waters, 42; and injury to plants, 33, 43, 54, 190, 196, 197, 238, 281, 351, 426, 466, 599, 633; problems of residues of, on sprayed fruit, etc., 32, 117, 128, 196, 330, 351, 412, 633, 634; Ephestia killed by residues of, in stored tobacco, 365; toxicity of, to sheep and cattle, 44; colloidal forms of, 119, 607; triplumbic form of, 416, 561, 600, 612; other insecticides compared with, 22, 39, 54, 55, 67, 91, 92, 119, 120, 196, 197, 238, 318, 330, 430, 561, 595, 599, 624, 633, 634, 640.

Leaf-cutting Ants (see Atta).

Leafhopper, Aster (see Macrosteles divisus).

Leafhopper, Beet (see Eutettix tenellus).

tenellus).
Leafhopper, Clover (see Aceratagallia

sanguinolenta).
Leafhopper, Indian Sugar-cane (see
Pyrilla perpusilla).

Leafhopper, Plum (see Macropsis

trimaculata).
Leafhopper, Potato (see Empoasca fabae).

Leafhopper, Taro (see Megamelus proserpina).

Leafhoppers, trap for collecting, on lettuce, 260.

Leather, Dermestids damaging, **453**. Lebia grandis (attacking Leptinotarsa decemlineata), limited possibility of acclimatisation of, in Europe, **26**.

Lecaniobius utilis, establishment of, against Saissetia oleae in Peru, 511.

Lecanium corni, on plum in Germany, 231; on fruit and shade trees in U.S.A., 210, 635; bionomics of, 635, 636; thrips predacious on, 210; sprays against, 231, 636.

Lecanium deltae, new parasite of, in Argentina, 344.

Lecanium hesperidum (see Coccus). Lecanium persicae, bionomics and forms of, in New Zealand, 58.

Lecanium persicae spinosum, subsp. n., on Wistaria in New Zealand, 58. lecontei, Agonoderus; Neodiprion. lectularius Gimer.

lectularius, Cimex. leefmansi, Doirania.

Leefmansia bicolor (parasite of Sexava), in Amboina, 380; variety of, in New Guinea Territory, 380; bionomics and utilisation of, 380.

lefroyi, Microbracon. leguminicola; Dasyneura.

Lema cyanella sapporensis, bionomics of, on wheat in Japan and Sakhalin, 93.

Lema melanopa, bionomics and control of, in Spain, 463.

Lema nigrovittata, on Datura and egg-plant in Hawaii, 35.

Lemon, pests of, in California, 295, 296, 297, 410, 561; weevil on, in Fiji, 389; not damaged by fruit-piercing moths in Gold Coast, 107; pests of, in New South Wales, 88, 335; pests of, in Transcaucasia, 653; factors affecting injury to, by insecticide treatments, 296, 410.

Lemon Oil, 612.

Lemon-grass Oil, 94.

Lenzites sepiaria, on spruce killed by bark-beetles in Sakhalin, **96.**

leperii, Epidiaspis (Diaspis). lepida, Ceromasia; Parasa.

Lepidiota caudata, in pastures in Queensland, 466.

Lepidoderma (see Dermolepida).

Lepidoglyphus destructor mixtus, measures against, in stored grain in Russian Union, **576**. Lepidosaphes, measures against, on Citrus in Transcaucasia, 653, 654.

Lepidosaphes alba, on cassava in Dominican Republic, 498.

Lepidosaphes beckii (on Citrus), in Fiji, 389; in U.S.A., 553, 597; in W. Indies, 261, 471, 498; Hyperaspis billoti predacious on, 471; factors affecting infestation by, 262; sprays against, 597; experiments fumigation on against, 553.

Lepidosaphes camelliae, on Camel-

lia in California, 223.

Lepidosaphes dispar, on cassava in Gold Coast, 492.

Lepidosaphes gloveri, not attacked by Comperiella bifasciata Spain, 457.

Lepidosaphes halli, eradication campaign against, in California, 222. Lepidosaphes ulmi, tests of oil

sprays against, in Ohio, 248. Lepinotus, list of species of, 416.

Lepinotus patruelis, in California, 416; infesting spinach seed from Holland, 416.

Lepisma saccharina, tests of spray on, in Australia, 56.

leprosus, Phymateus.

Leptinotarsa decemlineata (Potato Beetle), 112; in Canada, 22, 330, 368; in France, 24; in Jersey, 63; doubtful value of introduced and indigenous predators against, in Europe, 26, 70; in U.S.A., 39, 325, 638; fungus infesting, 368; measures against, 325, 330, 639; tests of insecticides on, 206, 207, 242; laboratory rearing of, 557.

leptis, Spilochalcis.

Leptoglossus gonagra, food-plants

of, in Argentina, 512.

Leptomastidea abnormis (parasite of Pseudococcus spp.), rearing and utilisation of, in Canada and U.S.A., 310; introduced into Dominican Republic, 498; Palestine, 339.

Leptomastidea brethesi, sp. n., parasite of Pseudococcus in Argentina,

344.

Leptomastix (parasite of Pseudococcus comstocki), in Japan, 425; introduction of, into U.S.A., 425.

Leptomastix dactylopii (parasite of Pseudococcus spp.), in S. America, 310, 355; introduction, rearing and utilisation of, in Canada and U.S.A., 310, 355; developing with difficulty in P. adonidum, 355.

Leptomastix flavus, parasite Pseudococcus comstocki in Palestine, 339.

Leptosphaeria coniothyrium, Cecidomyiid associated with, on raspberry in Britain, 283.

Leptothrips mali, bionomics of, in California, 210, 211.

Lestodiplosis, predacious on other Cecidomyiids in Britain, 313, 315.

Lestrodryinus (see Dryinus).

Lethane, 56.

Lethane Spreader, reducing scorching of cotton by cryolite dusts, 358.

lethifera, Blaps.

Lettuce, pests of, in Britain, 145;

Nezara on, in Fiji, 279; cutworms on, in Germany, 450; worms on, in Germany, Empoasca solana on, in Hawaii, 651; Penthaleus major on, in New South Wales, 237; pests of, in U.S.A., 537, 646; insects and virus disease of, 140, 646; trap for collecting leafhoppers on, 260; nutritional experiments with Pyrausta nubilalis and, 43.

Lettuce, Prickly (see Lactuca scari-

ola).

Leucania acontistis, damaging pastures in New Zealand, 334.

Leucania toroneura, damaging pastures in New Zealand, 334.

Leucania unipuncta (see Cirphis).

leucarthros, Necremnus.

Leucoceva laevicollis, on Malpighia punicifolia in Porto Rico, 473. leucoloma, Pantomorus (Naupactus). leuconotus, Anthores.

leucophthalmus, Philaenus.

Leucoplema dohertyi, new parasite of, on coffee in Belgian Congo, 514.

Leucoptera caffeina (on coffee), parasites and biological control of, in Tanganyika, 175-177.

Leucoptera coffeella (on coffee), original discovery of, in Guadeloupe, 35; parasites and biological control of, in Tanganyika, 175–177.

Leucoptera daricella, L. caffeina recorded as, in Tanganyika, 175.

leucoptera, Phytomyptera. leucopterus, Blissus.

Leucothrips theobromae, on cotton in Peru, 510.

leucotreta, Argyroploce. leveri, Phaenocarpa.

libania, Gorgopis. lienardi, Achaea. ligata, Chlorochroa.

Light, reactions of insects to, 156, 314, 527, 553; (artificial), insects attracted to, 167, 541, 602, 627.

Light-traps, for Delphacids and Jassids, 101, 517; for Lamellicorn beetles, 90, 298, 421, 432, 509, 614; for moths, 357, 534, 641; analyses of captures of insects in, 152, 224, 280.

lignosellus, Elasmopalpus.

ligustici, Otiorrhŷnchus (Brachyrrhinus).

Ligustrum (see Privet).

Ligustrum lucidum, food-plant of Ericerus pela in Szechuan, 153.

Ligyrus tumulosus, measures against, on lilies in Bermuda, 138.

Lilac, Prociphilus oriens on, in Japan, 173; Gracilaria syringella on, in Russia, 487; copper arsenate causing injury to, 119. lilacinus, Pseudococcus.

Lilies (*Lilium*), pests and virus diseases of, in Oregon and Ber- muda, 45, 138.

lilii, Myzus.

Lima Beans, Lepidoptera on, in Hawaii, 35; Argyroploce illepida on, in Java, 182; susceptibility of varieties of, to pests in Porto Rico, 345, 470; pests of, in U.S.A., 252, 481, 534, 535; Aphid transmitting cowpea mosaic to, 503.

limacina, Caliroa (Eriocampoides). Limber Pine (see Pinus flexilis).

Lime (Citrus), fungi associated with Pseudococcus comstocki on, in Brazil, 168; not damaged by fruit-piercing moths in Gold Coast, 107; Zeuzera coffeae on, in Malaya, 513.

Lime, spraying with, against Empoasca on potato, 325, 326, 440; addition of, to arsenical sprays, 10, 11, 148, 196, 325, 326, 406, 412, 479, 503, 634; addition of, to other sprays (see also Bordeaux mixture), 296, 412; as a carrier for dusts, 32, 53, 54, 92, 121, 136, 148, 162, 236, 249, 250, 252, 301, 359, 479, 504, 513, 596, 598, 603, 639, 640, 647; as a repellent, 391, 392, 539; stored cowpeas mixed with, against Bruchid, 59; formulae containing, 10, 54, 92, 136, 148, 162, 236,

250, 252, 325, 406, 412, 479, 503, 504, 513, 605, 634, 639, 640, 647. Lime-sulphur, against Coccids, 21, 223, 640, 642; preventing reproduction in Aspidiotus perniciosus, 409; unsatisfactory against Calandra oryzae, 540; tests of toxicity of, to Empoasca fabae, 441; against mites, 22, 100, 108, 109, 135, 237, 249, 281, 285, 286, 321, 330, 465, 641, 648; ineffective against Oxyptilus periscelidactylus, 133; against Psyllids, 21, 232, 278, 638, 639; against Scirtothrips citri, 296, 297; against Scolytus multistriatus and Ophiostoma ulmi, 644; effect of, against eggs of Stictocephala inermis, 216; as a repellent, 98, 171, 312, 472; in mixed dust, 639; formulae for, 21, 22, 23, 133, 278, 281, 284, 286, 321, 640 ; formulae containing, 21, 23, 109, 135, 162, 191, 223, 237, 249, 281, 296, 634, 639, 644; and calcium arsenate, 171, 281; and derris, 285; and lead arsenate, 162, 281, 285, 634, 644; and nicotine, 100, 108, 109, 127, 238; and oil emulsions, 23, 191, 223, 285, 642, 648; and sodium arsenite, 312; and sulphur, 21, 135, 249, 296, 297, 638; and tar, 472; zinc arsenite, 278; addition of other agents to, 162, 249, 281, 285, 286, 296, 638; and injury to plants, 21, 191, 281, 441, 598.

Limnaea, measures against, on watercress in Britain, 111.

limniace, Danaus (Danais).

Limnophilus lunatus, measures against, on watercress in Britain, 111.

Limonius ectypus (see Pheletes). Lina scripta (see Melasoma lineatopunctata).

linearis, Lyctus.

lineata, Sarcophaga (Blaesoxipha).

lineatella, Anarsia. lineaticollis, Antestia.

lineatopunctata, Melasoma,

lineatum, Trypodendron (Xyloterus). lineatus, Agriotes; Aleuas; Hyperstylus.

Lineodes integra, on Capsicum in Peru, 539.

lineola, Tectocoris (see T. dioph-thalmus).

lineolatus, Phytorus.

Linseed Oil (see Oils, Vegetable).

Linum usitatissimum (see Flax).

Liocola lugubris, Scoliids reared on, in Russia, 572.

Liothrips urichi, distribution of, against Clidemia hirta in Pacific Islands, 495, 496.

liparidis, Apanteles; Euplectrus. Liquidambar formosana, Lymantria dispar on, in China, 545.

Liquidambar styraciflua, Monolepta varicornis on, in Louisiana, 600. Liriomyza (see Agromyza).

Lissonota, parasite of Cydia lati-ferreana in U.S.A., 405.

Lissonota frigida, parasite of Crambus vulvivagellus in Kansas, 505. Listroderes costirostris, auct. (see L. obliquus).

Listroderes obliquus (on vegetables), measures in Hawaii, 35; Victoria, 31, 546, against, in 547.

Lithocarpus, Dichomeris oceanis on,

in Japan, 100.

Lithocolletis, on Tephrosia spp. in Porto Rico, 473; parasites of, 473.

Lithocolletis populifoliella, on poplar in Russia, 487; effect of leafsize on oviposition of, 487.

littoralis, Eucomys. litura, Prodenia.

lividus, Euderus.

Lixophaga diatraeae, failure to introduce, into Barbados against Diatraea saccharalis, 3.

Lixus truncatulus (see Hypolixus). Lizards, destroying noxious insects, 251, 380.

Lobotrachelus, parasite of, on Hibiscus in Madras, 522.

Locust, Bombay (see Patanga succincta).

Locust, Brown (see Locustana pardalina)

Locust, Bush (see Phymateus lepvosus).

Migratory (see Locust, Locusta migratoria).

Locust Tree, Black (see Robinia pseudacacia).

Locust-tree Leaf Miner (see Chalepus

dorsalis).

Locusta migratoria, distribution of, in Europe, 322, 323; in India, 322, 519; in Russian Union, 487, 569-571, 583; breeding centres, phases and migrations of, 322, 323, 487, 570, 571; other bionomics of, 570, 571; experiments with Beauveria bassiana against, 583.

Locusta migratoria migratorioides, in E. Africa, 177, 229, 491, 528, 589; in S. Africa, 322; in Belgian Congo, 388; in Gold Coast, 492; in N. Rhodesia, 491; in Sudan, 388; crops attacked by, 229, 491; migrations of, 229, 322, 491, 492; Empusa grylli infesting, 491; mites parasitic on, 177, 229, 322; Stomatorrhina lunata parasitising eggs of, 229; eye as phase character in, 341.

Locusta migratoria rossica, in Central Russia, 570; distribution of, 323. locustae, attacking Locustacarus Locusta migratoria migratorioides in E. Africa, 177.

Locustana pardalina (in S. Africa), phases of, 19; tests of baits and dusts against, 20, 613.

Locusts and Grasshoppers, in Br. E. Africa, 145, 175, 177, 228, 229, 388, 491, 528, 589; in Portuguese E. Africa, 156; in S. Africa, 19, 20, 143, 144, 156, 320, 322, 435, 612, 613; in Argentina, 308, 322, 497, 511, 512, 528, 538; in Australia, 141, 144, 169, 466; in Canada, 112, 143, 241, 280, 308, 309, 393, 394, 428, 648; in Belgian Congo, 228, 388; in Formosa, 94; in Gold Coast, 492; in Hawaii, 49, 650; in Br. Honduras, 498; in India, 269, 271, 323, 341-343, 519, 520, 523; in Peru, 539; in N. Rhodesia, 491; in Russian Union, 485, 487, 569, 571, 583; in Spain, 459, 460, 461, 529; in Sudan, 388; in Sumatra, 318; in Turkey, 289; in U.S.A., 48, 136, 143, 204, 205, 217, 222, 223, 224, 254, 255, 326, 327, 354, 394, 401, 428, 477, 536, 596, 603, 604, 638; identification of eggs of mid-western species of, 224; phases of, 19, 322, 323, 341, 459, 519, 570, 571; breeding centres and ecology of, 322, 484 486, 487, 497, 520, 570, 571; migrations of, 154, 228, 229, 322, 323, 428, 466, 491, 492, 497, 520, 523, 536, 543, 570, 571, 589, 604; effects of temperature, humidity and rainfall on, 141, 205, 341-343, 394, 484-486, 487, 497, 520, 523, 570, 583, 589, 603, 604; effect of hailstorms on, 394; changes in nymphal development of, 518, 604; physiology of diapause eggs of, 328; periodicity of, 12, 538; cultural

practices in relation to, 112, 254, 255, 497; as a factor in soil erosion, 204, 603; surveys and estimation of populations of, 222. 529, 537, 569, 570, 604; losses due to, 393, 537; algae on, 269; birds destroying, 217, 497; insect enemies of, 49, 94, 169, 229, 280, 401, 435, 477, 485, 486, 491, 566; mites parasitic on, 177, 229, 322; *Mermis* in, 486; fungi infesting, 220, 229, 485, 491, 497, 570, 583, 589; reactions of, to castor plants, 327; measures against (see also under Baits), 20, 145, 156, 169, 241, 289, 326, 435, 459, 460, 498, 529, 538, 539, 589; comparison and economic aspects of different methods of controlling, 223, 393, 459, 460, 529, 537; organisation of control of, 143, 144; international cooperation in control of, 62, 308, 322.

Loganberry, Incurvaria rubiella on, in Britain, 283; Polyphylla perversa on, in Br. Columbia, 508.

lolii, Contarinia.

Lolium, susceptibility of species of, to grass caterpillars in New Zealand, 89.

Lolium perenne, 89; Contarinia lolii on, in Britain, 315.

Lonchaea, Dacus confused with, in New South Wales, 237.

Lonchaea aurea, attacking damaged tomatos in Victoria, 32.

Lonchocarpus, uses of, in sprays, 109, 145; effect of diluents on toxicity of dusts of, 301; effect of variety and treatment on insecticidal constituents of, 470; rotenoids in, 135. (See Cubé and Timbo.)

longicaudata, Ctenolepisma.

longiceps, Orgilus.

longiclava, Anastatus antestiae.

longiclavis, Pagiophloeus.

longicorne, Stromatium.

longicornis, Paratrechina.

longimanus, Acrocinus. longipennis, Dissosteira; Longitar-

sus.

longipes, Anoplolepis. longirostris, Ischnaspis.

longisetaceus, Cheiloneurus.

longispinus, Pseudococcus (see P. adonidum).

longissima, Brontispa.

Longitarsus longipennis, bionomics and control of, on sweet potato, etc., in Crimea, 479.

Longitarsus parvulus, ecological factors affecting, on flax in Russia, 181, 182.

ongitarsus pellucidus, bionomics and control of, on sweet potato, Longitarsus etc., in Crimea, 479.

Long-tailed Mealybug (see Pseudococcus adonidum).

longulus, Cylindrocopturus.

lonicerae, Aleurodes.

Loochoo Islands, Cylas formicarius associated with fungus on sweet potato in, 97; Dacus cucurbitae in, 99; Myzus varians in, 45; Cerambycid damaging structural timber in, 103; attempted introduction of Opius fletcheri into,

Lophobaris piperis, bionomics and control of, in Netherlands Indies, 32, 33, 328.

Lophobaris servatipes, not attacking healthy pepper vines in Netherlands Indies, 33.

lophyri, Exenterus.

Lophyroplectrus luteator, liberation of, against Gilpinia polytoma in Maine, 637.

Loro, **249.**

Lorol (Sulphonated), as a spreader

for sprays, 145.

Louisiana, cotton pests in, 257, 358, 359, 599, 600, 602, 603; Cylas formicarius in, 413, 601; miscellaneous pests in, 600-603; Pantomerus spp. and regulations against them in, 494, 568; sugarcane pests in, 160, 354, 471, 601; beneficial insects and biological control in, 160, 354, 471, 600, 601, 602; Iridomyrmex humilis destroying termites in, 499.

lounsburyi, Aspidiotiphagus; Meta-

phycus.

Loxoblemmus haani, on vegetables in Japan, 173.

Loxostege frustralis, on fodder bushes in S. Africa, 19.

Loxostege sticticalis (on beet), in Russian Union, 479, 654; in U.S.A., 136, 161, 537; climatic · factors affecting distribution of, 161; measures against, 136, 654; Titanio normalis confused with,

lozinskii, Microbracon (Habrobracon) simonovi.

lucasseni, Podothrips.

Lucerne, 536; relation of grass-hoppers to, in Argentina, 497, 511: resistant to Lachnosterna in Canada, 90; Contarinia medicaginis on, in Hungary, 12; Contarinia pests of, in India, 265, 266, 523; Pantomorus spp. on, in New South Wales, 494, 608, 609; pests of, in Russian Union, 486, 576; Prodenia litura on, in Spain, 463; pests of, in Sudan, 387, 388; pests of, in U.S.A., 198, 202, 222, 223, 252, 354, 357, 376, 401, 402, 409, 429, 441, 567, 600, 601, 603, 604, 628, 640, 643; Lygus spp. affecting seed production of, 441, 567; reaction of Macrosiphum onobrychis to varieties of, 202; orthodichlorobenzene causing injury to, 199; (baled hay), Nodaria extinctalis on, in S. Africa, 19; (stored seed), Bruchid in, in Peru, 139; (meal), in baits for armyworms, 300.

Lucerne Gall-midge (see Contarinia medicaginis).

lucicola, Anomala (Pachystethus). lucidulus, Pseudocentrinus.

Lucilia sericata, tests of organic compounds on, 407.

luctuosa, Psyllobora. luctuosus, Cryptoideus.

Ludius inflatus (see Corymbites).

Luffa, new weevil on, in Philippines,

Luffa cylindrica, Engytatus geniculatus on, in Hawaii, 650.

lugens, Nilaparvata.

lugubris, Brachymeria; Liocola.

Lunaria, not attacked by Pieris rapae in Victoria, 546.

lunata, (Chilomenes); Cydonia Stomatorrhina.

lunatus, Limnophilus.

Luperodes brunneus (see Monolepta varicornis).

Luperodes quaternus, on Dolichos lablab in Sudan, 388.

lupulinus, Hepialus.

luridula, Anomis.

luridus, Troilus.

luteator, Lophyroplectrus.

luteipes, Azya. luteola, Epuraea (Haptoncus); Galerucella.

luteum, Trichogramma.

lutorius, Trogus.

lybica, Empoasca.

Lycaena boetica (see Cosmolyce). lycopersici, Phyllocoptes.

Lyctids, digestive enzymes of, 340. Lyctus, in Australia, 150, 447, 466; in Britain, 147, 447, 448; in U.S.A., 447; chemical treatments

of timber against, 146, 447, 448; determination of starch content of Eucalyptus in relation to, 150; depletion of starch from timber trees against, 466.

Lyctus brunneus (in timber), in Britain, 447, 448; in Queensland,

Lyctus linearis, in timber in Britain, 448.

Lyctus parallellipipedus, in timber in U.S.A., 447.

Lyctus planicollis, in timber in U.S.A., 447.

Lydella stabulans grisescens, considered correct name for Ceromasia senilis (q.v.), 331.

Lydinolydella metallica (parasite of Epilachna spp.), introduced into U.S.A. from Brazil, 431.

Lygaeus kalmii, tests of ovicides on, 352.

Lygidea mendax, on apple in New York, 2.

Lygidolon laevigatum, experiments against, on wattle in S. Africa, 19, 278.

Lygocerus, parasite of Pseudococcus comstocki in Virginia, 196.

Lygus, factors affecting damage to cotton by, in Uganda, 230; in U.S.A., 429, 441, 442, 557; affecting seed production of beet and lucerne, 429, 441, 567; study of damage to lucerne by, 441, 567; type of injury to peaches by, 441, 557; natural enemies of, 442; measures against, 429, 442, 557.

Lygus elisus, on lucerne and peach in U.S.A., 441, 557, 567; other food-plants of, 441.

Lygus hesperus, bionomics of, on lucerne, etc., in U.S.A., 441, 442, 567.

Lygus hesperus var. viridiscutatus, on peach in U.S.A., 557.

Lygus pratensis, on strawberry in Britain, 28; in U.S.A., 503; not transmitting cowpea mosaic, 503.

Lygus pratensis oblineatus, foodplants of, in U.S.A., 359, 361, 441; measures against, 359, 361. Lygus simonyi, on cotton in Uganda,

230.

Lymantria dispar, in Algeria, 514; in forests in China, 544, 545; variety of, on apricot in China, 446; in Japan, 171, 172; in U.S.A., 161, 164, 212, 412, 428; bionomics of, 171, 545; parasites of, 172, 514; carriage of eggs and larvae of, by wind, 164, 213; problem and importance of controlling spread of, 212; use of scent of females for attracting males of, 428; method of dusting woodland against, 412: sprays protecting apple against, 164.

Lymantria monacha, different importance of, in Latvia Germany, 85, 86.

Lysiphlebus, parasite of Myzus persicae in S. Rhodesia, 184.

Lysiphlebus testaceipes (parasite of Aphids), in Colombia, 54; in Hawaii, 651; in Oklahoma, 251.

M.

macaensis, Polistes.

Macalla plicatalis, on teak in Burma, 77.

machaeralis, Apanteles; Hapalia. Macrocentrus, parasite of Chilo zonellus in Ceylon, 610.

Macrocentrus ancylivorus, utilisation of, against Cydia molesta in Australia, 141; utilisation and value of, against C. molesta in U.S.A., 87, 129, 130, 192, 193, 295, 425, 549; parasite of C. latiferreana in Oregon, 406; Ancylis comptana the normal and laboratory host of, 192, 193, 549; apparent development of races of, 549.

Macrocentrus calacte, parasite of Argyroploce illepida in Fiji, 496. Macrocentrus cerasivorana, parasite of Tortrix cerasivorana in Quebec,

626.

Macrocentrus gifuensis, parasite of Pyrausta nubilalis in Massachusetts, 471; attempted introduction of, into Porto Rico against Diatraea saccharalis, 471.

Macrocentrus homonae, establishment of, against Homona coffearia

in Ceylon, 174.

Macrocentrus thoracicus, parasite of Cydia molesta in Japan and Korea, 372.

Macrocheles, attacking Bombyx mori

in Japan, 172.

Macrodactylus subspinosus, on vines in U.S.A., 139; intercepted on roses in Bermuda, 139.

Macronoctua onusta (Iris Borer), tests of ovicides against, in New York, 644.

macrophallus, Sturmia.

Macrophylla, habits and control of, damaging golf courses in Africa, 614.

Macrophylla nigricollis, 614.

Macropiper methysticum (see Piper). Macropsis insignis, on Prunus in Ontario, 307, 548.

Macropsis trimaculata, on Prunus spp. in Ontario, 306, 548; transmitting virus diseases of peach, 306, 548; bionomics of, 306; mode of feeding of, 307, 548.

Macrosiphoniella (see Macrosiphum). Macrosiphum avenae, bionomics of, on cereals, etc., in Japan and India, 100, 266.

Macrosiphum gei, auct. (see M. solanifolii).

Macrosiphum granarium (see M. avenae).

Macrosiphum onobrychis (pisi) (Pea Aphis), in Russia, 576; in U.S.A., 42, 136, 164, 202, 249, 353, 376, 404, 415, 503, 504, 631, 639, 640; development affecting recovery of peas from attacks of, 42; reaction of, to resistant varieties of lucerne, 202; relation of lucerne to attacks of, on peas, 376, 576, 639, 640; on clover, 164; susceptibility of species and varieties of vetch to, 404; transmitting mosaic diseases of narcisus and cowpeas, 415, 503; not feeding on potato, 165; natural enemies of, 164, 165; measures against, 136, 202, 249, 353, 504, 631.

Macrosiphum rubiellum (in Britain), possible vector of raspberry mosaic in, 287; on strawberry, 28. Macrosiphum rubifolium, possible

vector of raspberry mosaic in Britain, 287.

Macrosiphum sanborni, rearing and use of, for testing insecticide treatments, 29, 207, 591.

Macrosiphum solani (on potato), in Br. Isles, 28, 329; in Canada, 303; in Maine, 188; relation of, to virus diseases of potato, 188; winter food-plant of, 188; strawberry, 28.

Macrosiphum solanifolii, in Br. Isles, 28, 145, 283, 329; Canada, 303; in Hawaii, 35. 650; in Maine, 188; in New South Wales, 57; on beans, 145; on potato, 188, 303, 329, 650; on strawberry, 28, 283; rose the winter food-plant of, 188; transmitting virus disease of potato, 188; experiments with virus diseases of other plants and, **35, 57, 503**; feeding methods of, in relation to transmission of viruses, 179; measures against, 145, 650.

Macrosiphum solidaginis, on Carthamus tinctorius in India, 266.

Macrosiphum sonchi, on tomato in Hawaii, **651.**

Macrosteles divisus, relation of varieties of, to aster yellows in vegetables in U.S.A., 301, 302, 646; multiplication of asteryellows virus in, 502; 'overwintering of, 646; measures against, 647.

Macrosteles sexnotata, not transmitting mosaic diseases of cereals in Russian Union, 13, 16.

Macrotermes spp., habits and control of, damaging grasses in Kenya, 338.

maculata, Ceratomegilla (Megilla); Nephrotoma (Pales).

maculatus, Bruchus (Callosobruchus); Neoplectrus.

maculipennis, Plutella. maculiventris, Podisus.

Madagascar, Melittomma insulare on coconut in, 341; Exochomus flavipes predacious on Chironomids in, 83; Scolia oryctophaga introduced into Samoa from, 148. maderae, Calosoma.

madiunensis, Aulacaspis (Chionas-

Madras, insects and virus diseases of egg-plant, etc.; in, 542; foodplants and parasites of Pempherulus affinis in, 521-522; miscellaneous pests in, 517, 519.

Maerua, Achaea lienardi on, in S. Africa, 19.

Magdalis, Ophiostoma ulmi cultured from, in New York, 644.

Magicicada septendecim, persistance of damage to fruit trees, etc., by, in W. Virginia, **562.** agnesite, mixed with

Magnesite, with wheat against Calandra, 541.

Magnesium Arsenate, ineffective against Heliothis armigera, 535; and sand, applied to strawberry roots against Lachnosterna, 259.

Magnesium Carbonate, reducing dusts containing toxicity of rotenone, 301; ineffective against Dermestids, 454.

Mahogany (Swietenia), Xylothrips religiosus on, in Fiji, 170; pests of, in India, 75; pests of, in Porto Rico, 52, 427; failure to rear Zeuzera coffeae in, 271; resistance of timber of, to Eutermes costalis, 7. maidis, Aphis; Peregrinus.

Maine, forest pests in, 37, 86, 164, 347, 424, 427, 531, 637, 638; miscellaneous pests in, 88, 164, 594; Septis spp. on wheat and wild rice in, 560; Aphids and virus diseases of potato in, 165, 188; beneficial insects and biological control in, 86, 424, 637, 638.

Maize, Lepidoptera on, in S. Africa, 19, 615; pests of, in Canada, 90, 303, 508; pests of, in China, 446; pests of, in Colombia, 54; Ripersia internodia on, in Egypt, 30; Pyrausta nubilalis on, in Central Europe, 71, 72; pests of, in Fiji, 148, 170, 390, 496; pests of, in Guam, 331; Heliothis armigera on, in Hawaii, 650; pests of, in India, 81, 266, 523; P. nubilalis on, in Japan, 99; Heliothis armigera on, in New Guinea Territory, 377; pests of, in Nyasaland, 154, 589; pests of, in Peru, 511, 539; Nomadacris septemfasciata damaging, in N. Rhodesia, 491; Calandra oryzae on, in S. Rhodesia, 184; Prodenia litura on, in Spain, 463; Sesamia cretica on, in Sudan, 387; locusts damaging, in Uganda, 228, 229; pests of, in U.S.A., 2, 3, 43, 48, 88, 162, 164, 187, 214, 217, 244, 245, 250, 253, 275, 276, 277, 300, 354, 367, 419, 428, 476, 481, 482, 505, 508, 529, 534, 535, 560, 599, 600, 602, 603, 640, 641; pests of, in W. Indies, 5, 346, 469, 470; insects and virus diseases of, 13, 469; host range and insect vectors of bacterial wilt of, 244; varieties and time of planting of, in relation to pests, 44, 187, 217, 245, 250, 253, 303, 367, 481, 503, 529, 599; methods of protecting ears of, against pests, 217, 469, 470, 535, 640, 641; nutritional experiments with Pyrausta nubillalis and, 43, 44; treated with selenium against mites Aphids, 275, 276; cryolite causing injury to, 260.

Maize (Stored), pests in, and their control, 72, 73, 184, 210, 292, 429, 453, 455, 493; olfactory

reaction of Rhizopertha dominica to, 451; (cobs), in bait for Otiorrhynchus ligustici, 252, 253; (meal), in baits for locusts and Heliothis, 20, 429, 534, 598, 612, 631.

Maize Weevil (see Calandra oryzae). majalis, Amphimallus (Amphimallon).

major, Penthaleus; Polistes.

Malacosoma americana, bionomics and control of, in Canada and U.S.A., 395; tests of insecticides on, 119, 242.

Malacosoma disstria, bionomics and control of, in Canada and U.S.A.,

37, 395.

Malacosoma neustria, on fruit trees,

etc., in China, 446.

Malay Apple (see Eugenia malac-

censis).

Malaya, cacao pests in, 512; Citripestis sagittiferella on Citrus in, 91; coconut pests in, 79, 148, 620; Dacus spp. in, 90; Calotermes domesticus damaging sepetir wood in, 543; beneficial insects in, 79, 148, 513, 514, 620; deferment of introduction of Metagonistylum minense into, 538; beneficial insects introduced into other countries from, 148, 466.

malayensis, Euplectromorpha.

mali, Aphelinus; Leptothrips;

Psylla.

Malic Acid, as a larval attractant for sprays against Cydia pomonella, 129.

maligna, Empoasca.

malivorella, Coleophora. Leucocera Malpighia punicifolia, laevicollis on, in Porto Rico, 475. Malvastrum coromandelianum, Pempherulus affinis on, in India, 270, 521, 522.

Malvaviscus drummondi, Platyedra gossypiella on, in U.S.A., 430.

measures Mamestra brassicae, against, on crucifers in Britain, 145. Mammea americana (Mamey), Pseudococcus nipae on, in Porto Rico, 472.

Manchuria, Myzus persicae on peach in, 100; parasites of Cydia glycinivorella on soy beans in, 33, 98; Scolytids in, 592.

Mandarin Orange (see Orange,

Mandarin).

Manganese Sulphate, as a safener for sprays of lead arsenate and lime-sulphur, 162.

Mangifera indica (see Mango). mangiferae, Cryptorrhynchus; Dasyneura.

mangiferus, Paratetranychus.

manglissiensis, Scolytus.

Mango, pests of, in Ceylon, 594; Trypetid on, in Formosa, 94; moths damaging fruits of, in Gold Coast, 107; pests of, in Hawaii, 331, 332; pests of, in India, 269, 270, 271; pests of, in Peru, 511, 539; Cryptorrhynchus mangiferae intercepted in seeds of, in Fiji, 148.

Mango-Fruit-fly (see Dacus ferru-

gineus dorsalis).

Mangrove (see Bruguiera gymnorrhiza).

manifestator, Ephialtes (Ichneumon). Manila Hemp (see Musa textilis).

Manilabaris, gen. nov., species of, in Philippines and Java, 649. Manilabaris cucurbitae, sp. n.,

habits of, on cucurbits in Philippines, 649.

Manitoba, grasshoppers and their natural enemies in, 308, 309, 394, 648; other field-crop pests in, 648.

Maori Mite (see Phyllocoptruta oleivorus).

Maple (Acer), Tortrix forskaleana on, in Britain, 628; Xyleborus dispar on, in Br. Columbia, 328; pests of, in U.S.A., 210, 551, 635. marcida, Phaenacantha.

margaritaria, Metrocampa (see Campaea margaritata).

margaritata, Campaea.

Margaronia caesalis, on Artocarpus communis in Ceylon, 594.

Margaronia marinata, study of, on gambier, etc., in Sumatra, 105, 106; bacteria infecting, 106.

Margaronia pryeri, bionomics of, on mulberry in Japan, 102.

Margaronia punctiferalis, on Sarcocephalus cordatus in Burma, 78.

Margaronia pyloalis (on mulberry), in Formosa, 34; in India, 75; in Japan, 34, 101, 102; parasites of, **34, 75, 101.**

marginatorius, Exenterus. marginifrons, Tapeigaster.

mariana, Brontispa.

Marietta, parasite of Chrysom-phalus ficus in Brazil, **625**; hyperparasite of Ceroplastes of Chrysomof Ceroplastes rubens in Japan, 172.

Marietta javensis, parasite of Laccifer lacca in India, 264.

marinata, Margaronia. maritimus, Pseudococcus.

Markhamia platycalyx, new weevil on, in Uganda, 341.

marmorea, Digama. marnoana, Colias.

maroccanus, Dociostaurus.

Maroga unipunctata (see Cryptophasa).

marshalli, Bruchus (Callosobruchus). martinezi, Apion; Platystolus.

Maruca testulalis, on Xylia dolabriformis in Burma, 78; on beans and lima beans in Hawaii, 35, 650; food-plants of, in Porto Rico, 345; quarantine against introduction of, into U.S.A., 444; measures against, 345, 650.

Maryland, pests of china asters in, 361; Conotrachelus nenuphar in, 41; pests of leguminous plants in, 202, 208, 354; miscellaneous pests in, 438, 505, 629; new mite on strawberry in, 530; biological and other measures against Popillia japonica in, 125, 426,

504.

masakaensis, Diversinervus. maskelli, Morganella.

Maskellia globosa, causing witches' broom on Eucalyptus crebra in Queensland, 467.

masrensis, Trionymus.

Massachusetts, forest pests in, 87, 412, 424, 552; Oxyptilus periscelidactylus on vines in, 133; Fenusa on violets in, 418; beneficial insects and biological control in, 87, 424, 552; ants predacious on Reticulitermes flavipes in, 499; Macrocentrus gifuensis introduced into Porto Rico from, 471.

Mastrus neodiprioni, parasite of sawflies in U.S.A., 637, 638.

matsumurai, Crematogaster brunnea. Matthiola, introduction of selenium into, against mites, 207.

maura, Eurygaster

mauritanicus, Tenebroides.

mauritia, Spodoptera.

Mauritius, miscellaneous pests in, 610; Pseudococcus spp. on pineapple in, 514; sugar-cane pests in, 83, 609, 610; termites in, 6, 610; Coccids described by de Charmoy from, 424; genous fungi of, 424; entomobiological control in, 84, 609, 610; Scolia oryctophaga introduced Samoa from, 148.

Mayetiola destructor (Hessian Fly), in Russia, 576, 582, 618, 619; in U.S.A., 205, 215, 264, 366, 400, 505; parasites of, 400, 582, 618, 619; resistance of varieties and hybrids of wheat to, 215, 366, 505; cultural practices affecting damage by, 205, 254; measures against, 576.

Moth, Indian (see Plodia Meal interpunctella).

Mealworm (see Tenebrio molitor). Mealybug, Comstock's (see Pseudo-

coccus comstocki). Mealybug, Long-tailed (see Pseudococcus adonidum).

Mealybug, Pineapple (see Pseudococcus brevipes).

medicaginis, Aphis; Contarinia. Medicago denticulata, Bemisia tabaci

on, in India, 523. Medicago sativa (see Lucerne).

Mediterranean Flour Moth (see Ephestia kuehniella).

Mediterranean Fruit-fly (see Ceratitis capitata)

megacephala, Pheidole.

Megacoelum modestum, food-plants of, in Australia, 149.

Megaderus stigma, boring in lead sheathing in Brazil, 345.

megalocephala, Epicauta. megameli, Ootetrastichus.

Megamelus proserpina, natural enemies of, on taro in Fiji and Philippines, 170, 280.

Megarhogas, parasite of Plecoptera reflexa in India, 74.

Megaselia scalaris, bred from dead larvae of Heliothis armigera in Fiji, **390.**

Megatrioza vitiensis, parasite of, on Eugenia malaccensis in Fiji, 390.

Megilla (see Ceratomegilla).

Meigenia mutabilis, introduced into U.S.A. against Crioceris asparagi, 430.

Melanagromyza (see Agromyza). Melanichneumon (see Ichneumon). Melanocallis caryaefoliae, on pecan

in U.S.A., 159. melanocephala, Diabrotica. melanocephalum, Tapinoma. melanochra, Stathmopoda. melanogaster, Drosophila.

melanopa, Lema. Melanophila californica, bionomics of, on pines in California, 357. 551; effect on, of logging killed

trees, 357.

Melanophila gentilis, bionomics of, on pines in California, 551.

Melanoplus (in U.S.A.), effect of idle land on species of, 255; as a factor in soil erosion, 205, 603; reactions of, to castor plants, destroyed by sparrow hawks, 217.

Melanoplus bivittatus, introduced and native parasites of, in Canada, 309; in U.S.A., 48, 223, 255, 326, 327, 536, 603, 604; bionomics of, 48, 255, 603, 604; tests of contact dusts on, 326.

Melanoplus devastator, surveys for,

in California, 222.

Melanoplus differentialis, in U.S.A., 3, 48, 223, 255, 327, 328, 401, 603, 604; bionomics of, 3, 48, 255, 603, 604; physiology of diapause egg of, 328; parasite of, 401.

Melanoplus femur-rubrum, introduced parasite reared on, in Canada, 309; in U.S.A., 223,

603, 604; bionomics of, 604.

Melanoplus mexicanus, in Canada, 241, 309, 393, 394, 428; in U.S.A., 48, 205, 223, 255, 327, 394, 428, 536, 604; climatic and ecological factors affecting, 48, 255, 393, 394; migrations of, 428, 536; other bionomics of, 604; introduced parasite reared on, 309; measures against, 241, 428. Melanoplus packardi, effect of hail-

storms on, in Canada, 394. Melanotus, damaging potato in New

York, 645.

communis, Melanotus measures against, damaging sugar-cane, etc., in Florida, 476, 477.

Melasoma lineatopunctata, measures against, on willow and poplar in Kansas, 632.

Melcha nursei (see Goryphus).

Melia azadirachta, extract of, in spray against Schistocerca, 519.

Melia azedarach, Boarmia selenaria on, in India, 74; Bostrychid in, in New South Wales, 335.

Meligethes aeneus (on rape), mobile traps against, in Germany, 231; pyrethrum sprays against, in Russian Union, 585.

Melilotus, resistant to Lachnosterna

in Canada, 90.

Melilotus alba, probable reservoir of tobacco streak in Kentucky, 165.

Melilotus indica, Hypera brunneipennis on, in Arizona, 629.

Melilotus parviflora, Bemisia tabaci on, in India, 523.

Melissopus (see Cydia).

melitarae, Antrocephalus (Stomatoceras).

Melittia satyriniformis, cultural measures against, on squash in Connecticut, 163.

Melittobia acasta, parasitism of Cydia pomonella by, possibly facilitated by Dibrachys affinis in France, 66.

Melittobia hawaiiensis, hyperparasite of Artona catoxantha in Java and Malaya, 79; Melittobiopsis ereunetiphila recorded as, Hawaii, 274.

Melittobiopsis ereunetiphila, nomics of, in Hawaii, 274.

Melittomma insulare, on coconut in Madagascar, 341; bionomics and control of, in Seychelles, 340, 341.

Mellesis citri, sp. n., on Citrus in China, **524**.

mellillus, Drasterius (Aeolus). mellitor, Microbracon. mellonella, Galleria.

Melochia corchorifera, probably attacked by Pempherulus affinis

in Madras, **522.** *Melolontha hippocastani*, in Germany, 488; in forests in Latvia, 86; measures against, in Russian Union, 571, 572, 577; damaging crops and pastures, 577; nomics of, 86, 488, 571, 577; parasite of, 572.

Melolontha melolontha, damaging pastures in France, 83; time of flight of, in Germany, 488.

Melon, Diaphania spp. on, in Alabama, 405; Creontiades pallidifer on, in India, 523; Aspongopus viduatus on, in Sudan, 388; scorched by derris and sulphur dust, 405.

Melon Fruit-fly (see Dacus cucur-

bitae). melzeri, Merocentrum (Gryllica).

memnonius, Metaphycus. menapia, Neophasia.

mendax, Lygidea. mendesi, Dysdercus.

mendicus, Cleonus (Conorrhynchus).

mendosa, Dasychira. menippus, Rhaconotus.

menozzii, Neoaplectana.

Meraporus crassicornis (parasite of Mayetiola destructor), bionomics of, in Russia, 618, 619.

merceri, Contarinia.

Mercuric Chloride (Mercury Bichloride), against *Hylemyia bras*sicae on crucifers, 109, 383, 384; in spray against thrips, 414.

Mercurous Chloride, ineffective against Dilophus febrilis in turf, 527; uses of, against Hylemyia brassicae on crucifers, 109, 375, 383, 384, 385; seed treatment of onion with, against H. antiqua, 374; ineffective for protecting strawberry roots against Lachnosterna, 259.

sterna, **259.**Mercury Bichloride (see Mercuric Chloride).

meridionalis, Agathiella.

Merisus destructor, parasite of Mayetiola destructor in Russia, 582, 618, 619; bionomics of, 619.

Mermis, parasite of Acridids in Siberia, 486.

Mermis nigrescens, habits of, parasitic on Homona coffearia in Ceylon, 174.

Merocentrum, gen. nov., 292.

Merocentrum (Gryllica) melzeri, on Coccoloba ilheense in Brazil, **292.** Mescinia peruella, on cotton in

Peru, 40, 509.

mesochori, Hemiteles (Acrolyta).

Mesochorus ornatus, parasite of Rhogas in Nyasaland, 588.

Mesochorus tachypus, parasite of Apanteles spp. in Britain, 314.

Mesoleius tenthredinis, parasite of Pristiphoraerichsoni in Quebec, 37. Mesopeltis (parasite of Saissetia oleae), introduced into California from Africa, 385, 386.

Mesostenoideus clavinervis, parasite of Scirpophaga in India, 183.

Metagonistylum minense (parasite of Diatraea saccharalis), in Brazil, 4, 5, 470, 471; establishment and utilisation of, in Br. Guiana, 4, 538; introduction of, into U.S.A., 354, 431, 471; distribution and utilisation of, in W. Indies, 4, 5, 263, 470, 471; deferment of introduction of, into Malaya, 538; not parasitising D. canella, 538; parasite of, 470; strains of, 4, 354, 431, 470, 471; bionomics and experiments in crossing strains of, 4; technique of rearing, 5, 354.

metallica, Lydinolydella; Plagiodera (Coccinella); Pleurotropis. Metallus albipes, on Rubus in

Britain, **544.**

Metallus gei, on Geum in Britain, 544.

Metallus pumilus, on Rubus in Britain, 544.

Metalnikov's Bacilli, experimental utilisation of, **583**, **584**.

Metamasius sericeus, on sugar-cane in Jamaica, 293.

metanastriae, Rhogas (see R. den-drolimi).

Metaphycus (parasitising Saissetia oleae), African species of, and their introduction into California, 385, 386.

Metaphycus helvolus, 385, 386. Metaphycus lounsburyi, 385, 386.

Metaphycus memnonius, sp. n., parasite of Eriococcus coriaceus in S. Australia, 290, 546; recorded as Metaphycus sp., 546.

Metaphycus stanleyi, 385, 386.

Metaplectrus thoseae, gen. et sp. n.,
hosts of, in Ceylon and Sumatra,
514.

Metarrhizium anisopliae (infesting Lamellicorns), in Algeria, 83; in Canada, 90; introduced into Formosa from Hawaii, 93.

Meteorological Conditions, relations of insects to, **152.** (See Humidity, Rainfall and Temperature.)

Meteorus, parasite of Calligrapha in Quebec, 627.

Meteorus autographae, parasite of Colias eurytheme in Lousiana, 600.

Meteorus dichomeridis, parasite of Dichomeris eridantis in India, 74.

Meteorus dumbletoni, parasite of Tortrix postvittana in Tasmania, 151.

Methyl Bromide, fumigation of soil, plants and planting materials with, 199, 203, 208, 413, 426, 499, 554, 555, 568; soil treatments with solutions containing, 203, 208; against pests in fruit and vegetables, 18, 113, 123, 124, 138, 222, 364, 499; against Bruchids and pests of stored grain, 19, 222, 429; determination of toxicity of, to Thermobia domestica, 554; absorption of, by food-stuffs, 444; causing injury to plants and fruit, 203, 364, 413, 499, 554; methods of fumigating with, 113, 123, 124; risks associated with, 444.

Methyl Eugenol, in baits for Trype-

tids, **91, 94.**

Methyl-n-amylketone, intermediary solvent for derris resins when used with oils, **350**.

meticulosa, Brotolomia.

meticulosalis, Terastia.

Metrocampa margaritaria (see Campaea margaritata).

Metroxylon sagu (Sago Palm), coconut pests on, in New Guinea Territory, 378, 379.

Mexican Bean Beetle (see Epilachna

varivestis).

mexicana, Epilachna.

mexicanus, Melanoplus.

Mexico, Myzus porosus in, 46; parasite of Epilachna in, and its attempted introduction into U.S.A., 114, 115.

Meyer Colour Test, 135.

micans, Dendroctonus.

Mice, destroying noxious insects, 38, 223.

Michelia champaca, Urostylis punctigera on, in India, 74.

Michigan, gladiolus thrips in, 277; grasshoppers and soil erosion in, 204; pests of pines in, 563, 605; Scolytus quadrispinosus in, 116; pests of vines in, 117, 558; insect food of trout in, 224. michiganensis, Microphthalma.

Micrapate, boring in lead sheathing

in Brazil, 345.

Microbiology, of insects, 285.

Microbracon, parasite of Lepidoptera in India, 74, 184; parasite of Saperda candida in New York, 243; parasite of Anomis luridula in Peru, 510.

Microbracon albolineatus, parasite of Chilo zonellus in Ceylon, 610; importation of, against sugarcane pests in Mauritius, 610.

Microbracon brevicornis, 184; hosts of, in S. Africa, 17, 611; parasite of Nephantis serinopa in India, 620.

Microbracon greeni (parasite of lac predators), laboratory breeding and utilisation of, in India, 264, 265; not pairing with M. lefroyi, 264.

Microbracon hebetor, laboratory breeding and utilisation of, against lac predators in India, 264, 265; parasite of Ephestia spp., 377, 616; experimental hosts of, 264; doubtful record of, 184; theory of sex determination by chromosomes in, 474.

Microbracon juglandis (see M. hebe-

tov).

Microbracon kirkpatricki (parasite of Platyedra gossypiella), in Nyasaland, 588; in Sudan, 386; introduced into U.S.A. from Egypt, 430.

Microbracon lefroyi, not pairing

with M. greeni, 264.

Microbracon mellitor, parasite of Chalcadermus aeneus in U.S.A., 405.

Microbracon omiodivorum, parasite of Nacoleia accepta in Hawaii, 49.

Microbracon sanninoideae, parasite of Aegeria exitiosa in Ontario, 226.

Microbracon simonovi (with f. lozinskii, n.), parasite of Heliothis armigera in Azerbaijan, 235.

Microcryptus basizonus, parasite of Neodiprion sertifer in Latvia, 86; breeding and liberation of, against Gilpinia polytoma in Maine, 637; fungus infesting, 637; effect of food on adults of, 394.

Microcryptus tsugae, parasite of Neodiprion tsugae in Oregon, **552.** Microctonus basalis, parasite of

Longitarsus spp. in Crimea, 479. Microdus agilis, parasite of Tortrix cerasivorana in Quebec, 626.

Microdus diversus, importation and colonisation of, against Cydia molesta in U.S.A., 87, 129, 161, 295.

Microdus nucicola, sp. n., parasite of Cydia latiferreana in U.S.A., 86, 406.

Microdus pini, sp. n., parasite of Rhyacionia spp. in U.S.A., 86, 87.

Microdus stigmaterus (parasite of Diatraea saccharalis), in Brazil, 470; introduction of, into Porto Rico, 470, 471.

Microgaster plecopterae, parasite of Plecoptera reflexa in India, 74.

Micromalthus debilis (attacking timber), in Hawaii, 275; distribution of, 275.

Micromortar, description of, for triturating insects, 349.

Micromyzus formosanus, on chives and Japanese onion in Hawaii, 651.

Microphanurus semistriatus, parasite of Eurygaster integriceps in Russia, 575, 582.

Microphanurus suranus, parasite of Antestia in Uganda, 230.

Microphanurus vassilievi, parasite of Eurygaster integriceps in Russia, **582**.

Microphthalma, parasite of Geotrogus deserticola in Algeria, 83. Microphthalma capensis (see

sejuncta).

Microphthalma disjuncta, auct. (see M. nigra).

Microphthalma europaea, parasite of Melolonthids in Zanzibar, 83.

Microphthalma michiganensis, parasite of Lachnosterna in Canada,

Microphthalma nigra, parasite of Lachnosterna in Canada, 90.

Microphthalma sejuncta, in Africa, 609; imported into Mauritius against Lachnosterna smithi, 609.

Microplectron fuscipenne, utilisation and value of, against Gilpinia polytoma in Canada and U.S.A., 36, 38, 424, 427, 432, 531, 532, 637; parasitising Pristiphora erichsoni, 532.

Microplitis theretrae, parasite of Theretra japonica in Japan, 40. microptera, Pararcyptera (Arcyp-

tera).

micropterus, Eupteromalus.

Microtermes, damaging vines in S. Africa, 19; attacking tung trees in Nyasaland, 588; measures against, 589.

Microterys elegans, sp. n., parasite of Coccids in Argentina, **344.**

Microterys kenyaensis (parasite of Saissetia oleae), in Kenya, 385; introduced into California, 385.

Microterys saissetiae (parasite Saissetia oleae), in Uganda, 386; introduced into California, 386.

Microterys speciosus, parasite of Ceroplastes rubens in Japan, 172. Microtorus (see Hemiteles).

microzonus, Eupelmus.

Mictis profana, on Citrus in Fiji, 389.

middletoni, Aphis.

migratoria, Locusta.

migratorioides, Locusta migratoria. Migratory Grasshopper, Lesser (see Melanoplus mexicanus).

Migratory Locust (see Locusta migratoria).

miliaris, Asterolecanium.

militaris, Apanteles.

Milk, uses of, in sprays, 113, 160, 215, 371.

Milky Diseases, types of, infesting Popillia japonica and their utilisation in U.S.A., 124, 125, 369, 426, 504.

Millepedes, mite parasitic on, in S. Africa, 322; relation of, to potato scab in New York, 644.

Millet (see Panicum miliaceum). Millet, Italian (see Setaria italica). Millettia pachycarpa, insecticidal properties of, 483.

mimus, Dysdercus.

minense, Metagonistylum.

Mineola vaccinii, measures against, on cranberry in Washington, 640,

Minnesota, Crioceris duodecimpunctata on asparagus in, 48; fungi associated with Trypodendron spp. in aspen and birch in, 165; Harmologa fumiferana on conifers in,

minor, Apanteles; Baeoanusia; Calotermes; Empoasca; Labia; Myelophilus; Oncometopia; Pinnaspis (Hemichionaspis).

Minthea rugicollis, in timber of Anogeissus in Burma, 78.

minutula, Platygaster (see P. hie-

minutum, Trichogramma.

Dinoderus; minutus, Laemophloeus.

Miscanthus, Phaenacantha marcida

on, in Formosa, 104. Mischocyttarus phthisicus, predacious on Terastia meticulosalis in Porto Rico, 469.

Miscogaster, parasite of Mayetiola destructor in Russia, 618.

Mississippi, Cirphis unipuncta on oats in, 356; pests and disease of cucurbits in, 647.

Missouri, Cydia pomonella on apple in, 45, 160, 633; Contarinia juniperina on Juniperus spp. in, 432; Datana integerrina in, 166; beneficial insects in, 167, 166; 190, 433.

Misumenops asperatus, destroying Anasa tristis in Connecticut, 443.

Mites, attacking insects, 172, 177, 211, 215, 229, 322, 354, 376, 379, 488, 575, 620; attacking other mites, 306, 321; natural enemies of, 175, 210, 211, 273, 282, 306, 587; classification and new species of, 271, 444, 530, 565.

mitratus, Gryllulus (Gryllus).

miwai, Ochyromera.

Mixdrin, 232.

mixta, Orgyia.

mixtus, Anthonomopsis; Lepidoglyphus destructor.

Mocis latipes, on soy beans in U.S.A., 599.

moderatus, Geocoris.

modestum, Megacoelum.

moelleri, Solenopsis.

moesta, Proutista.

Molasses, in baits for grasshoppers and locusts, 144, 269, 461, 612; in baits for other insects, 304, 542, 583, 584; effects of, in baitsprays for Trypetids, 54, 55, 337; as a larval attractant for sprays against Cydia pomonella, 129; other uses of, in sprays, 352, 413, 414, 607, 636; and lead arsenate, plants swabbed with against Heliothis, 148, 466.

Mole-cricket (see Gryllotalpa).

Mole-cricket, Porto Rican (see Scapteriscus vicinus).

molesta, Cydia (Grapholitha, Laspeyresia); Solenopsis.

molestae, Angitia (Dioctes, Inareo-

lata); Apanteles. molitor, Tenebrio.

mollis, Ernobius.

mombinpraeoptans, Anastrepha.

Momordica charantia, new weevil on, in Philippines, 649.

monacha, Lymantria (Ocneria). Monanthia monotropidia, on Cerdana alliodora in Porto Rico, 51. Monarda, Psallus seriatus on, in

Texas, 198. Monarthropalpus buxi, measures against, on box in U.S.A., 636.

Monda, on tea in Nyasaland, 60. Moneilema ulkei, introduced into S. Africa against Opuntia, 18.

moneta, Plusia.

Monilia candida, Trypodendron spp. associated with, in trees

Minnesota, 165.

Monilia cinerea, Rhynchites bacchus associated with, on fruit trees in Russia, 581.

moniliferus, Pachyrrhynchus.

Monkeys, destroying Cactoblastis cactorum in S. Africa, 17.

Monkshood (see Aconitum).

Monochamus, measures against, attacking pine logs in Canada,

Monogonogastra agrili, parasite of Saperda candida in New York, 243.

Monolepta varicornis, habits and control of, on maize, cotton, etc., in Louisiana, 600.

Monomorium, predacious on Pempherulus affinis in Madras, 522. monotropidia, Monanthia.

Montana, Anabrus simplex and its natural enemies in, 187, 537; pests of conifers in, 371, 537, 648; grasshoppers in, 428, 536,

miscellaneous pests in, 537 : Spilochalcis leptis in, 190.

montana, Pimpla (Itoplectis).

montanus, Goniozus.

monticolae, Dendroctonus. montrouzieri, Cryptolaemus.

Moonia albimaculata, suspected vector of spike disease of sandal in India, **75.**

Moorish Literature, early references to insect pests in, 157.

moreleti, Julus.

Morganella maskelli, on fig in Brazil, 392.

mori, Tetranychus.

Mormon Cricket (see Anabrus simplex).

Morocco, parasite of Melolonthids in, and its introduction into Mauritius, 83, 84, 609.

morogorensis, Allodontermes.

morosa, Lachnosterna (Holotrichia); Rhyparida.

morstatti, Xyleborus.

Morus (see Mulberry).

mosellana, Sitodiplosis (Thecodiplosis).

Mosquitos, tests of toxicity of insecticides on immature stages of, 414, 433.

mossambicus, Hodotermes.

motti, Empoasca kerri. moultoni, Frankliniella.

Mountain Ash (see Sorbus americana).

Mountain Pine Beetle (see Dendroctonus monticolae).

Mulberry, Lepidoptera on, in Formosa, 34, 93; pests of, in India, 75, 271; Margaronia spp. on, in Japan, 34, 101, 102; Lachnosterna morosa on, in Korea, 171; Lecanium corni on, in Oklahoma, 635; preference of grasshoppers for, 603.

multilineata, Zagrammosoma. multipunctata, Calligrapha.

multistriatus, Scolytus.

Mundulea, rotenoids in, 135.

Mundulea sericea, insecticidal properties of varieties of, 29.

Mungo Bean (see Phaseolus mungo). Murgantia histrionica, food-plants of, in Hawaii, 35, 651; in U.S.A., 35, 560; egg-parasites and biological control of, 560, 651.

murgantiae, Trissolcus.

Musa, Aphis gossypii transmitting virus disease of banana to other species of, in New South Wales, 57. (See Banana.)

Musa textilis, Aphids and virus diseases in, in Australia and

Philippines, 57, 317.

Musca domestica, insecticide tests on, 127, 301, 398, 414, 470, 555, 556.

Muscina assimilis, possible parasite of Polyphylla perversa in Br.

Columbia, 509.

Muscina stabulans, possible parasite of *Polyphylla perversa* in Br. Columbia, **509**; parasitising Acridids in Texas, **477**.

Mushroom Houses, equipment for sulphur fumigation of empty,

276.

Muslin Covers, use of, for protecting stored tobacco from Lasioderma serricorne, 493.

Mussidia nigrivenella, in stored cacao in Gold Coast, 616.

Mussidia pectinicornella, bionomics of, on Parkia speciosa in Java, 182.

Mustard, new Jassid on, in India, Rhopalosiphum pseudobrassicae on, in U.S.A., 596.

mutabilis, Carpophilus; Meigenia. mutilatus, Xyleborus.

Mycetophilids, soil insecticides against, on cucumber in Britain,

Mycodiplosis alternata, bionomics and control of, on Cornus in

U.S.A., 606.

Myelophilus spp., prevention of outbreaks of, due to felling of conifers in Britain, 323, 382, 383,

Myiophasia globosa, parasite of Chalcodermus aeneus in U.S.A., 405.

Mylabris, destroying eggs of Acridids in Siberia, 485.

Mylabris phalerata, destroying locust eggs in Formosa, 94.

Myllocerus laetivirens, food-plants of, in India, 523.

Myllocerus spinicollis, sp. n., on Zizyphus jujuba in Punjab, 341.

Myopa, parasitising honey bee in Ontario, 473; hosts of, in Europe, 474.

myrmecodiae, Iridomyrmex.

Myrmecopria, probable hyperparasite of Paradexodes epilachnae in Ohio, 115.

Mystrops palmarum, sp. n., on Cocos spp. in Brazil, 10.

Mytilococcus (see Lepidosaphes).

Myzine quinquecincta, parasite of Lamellicorns in Kentucky, 299. Myzocallis annulata, utilisation of parasites against, on oak in Australia, 142.

Myzodes (see Myzus).

Myzus, revision of, in N. America, 45.

Myzus cerasi, on cherry in New York, 415; transmitting mosaic of narcissus, 415.

Myzus circumflexus, on papaya and rose in Hawaii, 651; feeding methods of, in relation to transmission of viruses, 179.

Myzus clematifoliae (see M. varians). Myzus hieracii, on lettuce in Britain,

Myzus lilii, sp. n., suspected vector of virus diseases of Lilium candidum in Oregon, 45.

Myzus ornatus, on strawberry in Britain, 28; in California, 45;

intercepted in U.S.A., 45.

Myzus persicae, in S. Africa, 19; in Argentina, 608; in Australia, 144, 465; in Br. Isles, 145, 329, 381, 590, 591; in Canada, 303; in France, 23, 24; in Germany, 235; in Hawaii, 35, 650; in India, 523; in Netherlands Indies, 318, 492; in Manchuria, 100; in Nyasaland, 154; in S. Rhodesia, 84, 184; in Spain, 457; in U.S.A., 45, 188, 638; on peach and plum, 23, 100, 144, 188, 465, 523, 590; on crucifers, 145, 235, 329, 381, 590; possibly transmitting virus disease cauliflower, 235; on potato, 303, 329, 381, 523, 590, 591, 650; relation of, to potato virus diseases, 23, 24, 179, 188, 329, 638; on tobacco, 19, 179, 318, 492, 523; relation of, to virus diseases of tobacco, 84, 154, 184, 608; on other plants, 35, 145, 154, 523; relation of, to virus diseases of other plants, 35, 65, 457, 503, factors affecting transmission of certain viruses by, 27, 65, 66, 179, 180; notice of list of food-plants of, 84; bionomics of, 23, 100, 188; natural enemies of, 184, 303; measures against, 144, 145, 184, 189, 318, 381, 492, 638; tests of insecticides on, 132, 353.

Myzus porosus, on rose and strawberry in U.S.A. and Mexico, 45, 46, 256.

Myzus pseudosolani (see Macrosiphum solani)

Myzus scammelli, sp. n., on cran-

berry in U.S.A., 45.

Myzus sensoriatus, sp. n., on Crataegus in U.S.A., 45.

Myzus varians (tropicalis), foodplants, synonymy and distribution of, 45.

Nabis ferus, predacious on other Rhynchota in U.S.A., 442, 443. Nacoleia accepta, parasites of, on sugar-cane in Hawaii, 49.

Nacoleia annubilata, hyperparasite

of, in Ceylon, 620.

naevana, Rhopobota. nana, Trichogrammatoidea.

nanus, Hemiteles (Astomaspis).

Naphtha, 239, 240.

Naphthalene, and oils, in sprays against Coleoptera in logs, 312, 427; soil treatment with, against pests, 110, 111, 146, 282, 448; pests, 110, 111, for sum formulae containing, for sum 110, 111; other uses treatment, 110, 111; other uses of, as a fumigant, 25, 56, 132, 555; and nicotine, 132; method of volatilising in orthodichlorobenzene, 555, 556; as a repellent against Gnorimoschema operculella, 138.

Naphthalenes, Chlorinated, and · oils, in sprays against Coleoptera in elm logs, 427; as a preservative against termites and Lyctus, 142, 147; types of, 142, 427.

Naphthalene Sulphonates, use of in sprays and dusts, 360, 600 (see Vatsol)

nararia, Natada.

Narcissus, Aphids and mosaic of, in New York, 415.

Narosa conspersa, parasite of, on tea in Ceylon, 174.

Narrow-leaved Ironbark (see Eucalyptus crebra).

Nasturtium (see Tropaeolum).

nasuta, Prorops.

Nasutitermes (see Eutermes).

Natada nararia, parasites of, on tea in Ceylon, 174, 514. natadae, Platyplectrus.

natalensis, Macrotermes (Termes);

Nysius. Naupactus (see Pantomorus). nawai, Pachyneuron; Telenomus.

Neasteropaeus, parasite of Stomatomyia bezziana in Ceylon, 621.

Nebraska, miscellaneous pests in, 277, 278, 433, 603; Exenterus spp. parasitic on sawflies in, 294.

Necremnus, revision of, 569.

Necremnus duplicatus, sp. n., parasite of Ceuthorrhynchus assimilis in Washington, 569.

Necremnus leucarthros, imported into U.S.A. against Hypera vari-

abilis, 569.

Necremnus oregonensis, sp. n., parasite of Hypera rumicis in Oregon, 569.

Nectandra rodioei, resistance of timber of, to termites, 7, 138.

Nectarine, Capsid on, in Australia, Tortrix sarcostega on, in 149; Baluchistan, 518.

neivai, Himatidium.

Neivamyrmex nigrescens (see Eciton).

Nematodes, infesting insects, 116, 173, 174, 187, 191, 211, 270, 316, 369, 370; utilisation of, against Popillia japonica, 370.

Nematospora, Antestia associated with, on coffee in Tanganyika,

Nematus ribisii (Gooseberry Sawfly), tests of pyrethrum against, in Russia, 585.

nemea, Zenillia (Phryxe).

Nemeritis tectonae, parasite of Xyleutes and Indarbela in Burma, 76.

nemoralis, Neurotoma. nemorata, Phyllotoma.

nemorivagus, Nysius.

nenuphar, Conotrachelus.

Neoanisotylus bimaculatus, gen. et sp. n., parasite of Eriosoma lanigerum in Argentina, 445.

Neoaplectana glaseri, bionomics and economic importance of, U.S.A., 369, 370.

Neoaplectana menozzii, bionomics of, infesting Cleonus mendicus in Italy, 316.

Neoclytus acuminatus, on elm in New York, 628.

Neoclytus unicolor, on apple in Peru, 511.

Neocoelostoma erophila, new parasite of, in Argentina, 344.

Neocopidosoma roccidophaga, gen. et sp. n., parasite of Icerya in Argentina, 344.

Neodendrin, 155, 232.

Neodiprion (in forests), parasites and biological control of, in Canada and U.S.A., 293, 294, 637, 638.

Neodiprion abietis, on spruce in Canada, 36, 530.

Neodiprion grandis (see Zadiprion). Neodiprion lecontei, on pine in Quebec, 37.

Neodiprion pinetum, on pine Quebec, 37.

Neodiprion sertifer (on pine), in Canada and U.S.A., 293, 294; in Europe, 86, 294; in Japan, 294; parasites and biological control of, 86, 293, 294, 552.

Neodiprion tsugae, bionomics of, on Tsuga heterophylla in Oregon, 294, 552.

neodiprioni, Mastrus.

Neomaskellia bergi, on sugar-cane in Guam, 331.

Neopales pavida, parasite of Cirphis unipuncta and Bombyx mori in Japan, 98.

Neophasia menapia, effects of defoliation by, on Pinus ponderosa in U.S.A., 422.

Neophysopus (see Anaphothrips).

Neoplectrus, gen. nov., 79.

Neoplectrus areolatus, sp. n., hosts of, in Celebes and Malaya, 514.

Neoplectrus bicarinatus, sp. n., parasite of Artona catoxantha in Java, 79.

Neoplectrus maculatus, sp. n., parasite of Artona catoxantha in Java,

Neotermes (see Calotermes).

nephantidis, Brachymeria; Elas-

mus; Perisierola.

Nephantis serinopa, natural enemies and biological control of, on coconut in Ceylon, 594, 619-622; in India, 620; bionomics of, 619.

Nephotettix apicalis (see N. bipunctatus).

Nephotettix bipunctatus cincticeps, bionomics of, on rice in Japan, 101; retention of dwarf disease virus through several generations by, 272.

Nephrosperma, Melittomma insulare associated with, in Seychelles, 341. Nephrotoma maculata, damaging

cereals and pastures in Germany,

Nesopimpla, considered congeneric with Itoplectis, 220.

Netherlands Indies, new Coccid on Amorphophallus variabilis in, 88: pests of gambier in, 105, 106; Lophobaris on pepper in, 32, 33, 328; miscellaneous pests in, 155, 270, 319, 649; pests of Pithecolobium lobatum and Parkia speciosa in, 182; sugar-cane pests in, 92, 524; pests and diseases of tobacco in, 318, 319, 492; Lasioderma serricorne in stored beneficial tobacco in, 493; insects in, 33, 79, 105, 106, 380, 492, 514, 524, 620; beneficial insects introduced into Formosa from, 92.

Nettle Weed (see Lamium amplexicaule).

Neurotoma nemoralis, tests of contact sprays against, on peach in Germany, 233.

neustria, Malacosoma.

Nevada, Anabrus simplex in, 562; Spilochalcis leptis in, 190.

nevadensis, Zootermopsis.

New Brunswick, potato Aphids in, 302; Cydia nigricana on peas in, 310: forest pests in, 37, 38, 39, 347, 348, 531, 532; Incurvaria rubiella on raspberry in, 307; beneficial insects and biological control in, 38, 39, 303, 532; disease of Gilpinia polytoma in, 531,

New Caledonia, miscellaneous pests in, **391.**

New Guinea, Dutch, parasites of Sexava in, 380.

New Guinea Territory, coconut pests in, 91, 377, 379, 380; miscellaneous pests in, 377; natural enemies and biological control of insects in, 377, 378, 379, 380.

New Hampshire, outbreak, disease and parasites of Gilpinia polytoma on spruce in, 37, 422, 424,

427, 531, 637, 638.

New Jersey, new Aphid on cranberry in, 45; forest pests in, 212, 294, 428, 438, 552; Scolytids and Dutch elm disease in, 119, 212, 365, 550; orchard pests in, 127, 129, 130, 194, 195, 295, 364, 549; pests of ornamental plants in, 132, 203; wireworms in, 122, 123; virus diseases of plants in, 502, 646; diseases and natural enemies of Popillia japonica and their utilisation in, 125, 368, 369, 370: parasites and biological control of other insects in, 129, 130, 193, 294, 295, 549, 552.

New Mexico, cotton pests in, 430; Cydia pomonella in, 414; potato pests in, 638.

New South Wales, pests of Brachychiton populneus in, 467; pests of fruits and fruit-trees in, 54, 88, 141, 144, 149, 151, 237, 290, 298, 335, 336, 337, 493, 494, 609; Aphids and virus disease in banana, etc., in, 57; grassgrasshoppers in, 141, 144, 466; miscellaneous pests in, 170, 236, 237, 335, 436; Aphodius damaging pastures in, 436, 547; Pantomorus spp. in, 494, 608, 609; Pieris rapae in, 290, 436; other vegetable pests in, 31, 142, 236, 290, 335; natural enemies and biological control of insects in, 142, 170, 298, 493, 608; biological control of prickly-pear in, 587; pests imported into Fiji from, 148, 391, 496.

New York, Coleoptera and Ophiostoma ulmi on elm in, 119, 120, 502, 551, 563, 627, 644, 645; other forest pests in, 86, 212, 400, 421, 424, 428, 432, 438, 552 Otiorrhynchus ligustici on lucerne etc., in, 198, 252, 357, 643; Lepidoptera on maize in, 481, 482; orchard pests in, 1, 2, 119, 127, 128, 196, 292, 293, 563, 643, 645; pests of ornamental plants in, 86, 136, 415, 481, 644; vegetable pests in, 121, 221, 481, 643, 644, 645, 646; insects and virus diseases of plants in, 302, 415, 646; Alabama argillacea in, 404; Attagenus piceus in, 644; Lachnosterna spp. in, 258, 400, 421, 432, 643, 644; new records of European Tortricids in, 628; beneficial insects and biological control in, 86, 121, 243, 424, 552.

New Zealand, insects damaging grasses in, 89, 333, 334, 541; miscellaneous pests in, 333, 335, 545; orchard pests in, 58, 150, 151, 333; forms of Lecanium persicae in, 58; Archichauliodes diversus in, 334; beneficial insects and biological control in, 64, 151, 314, 333, 545; biological control of noxious plants in, 334; introduction of beneficial insects into Australia from, 142, 468, 546.

New Zealand Dobson-fly (see Archichauliodes diversus). New Zealand Spinach (see Tetragonia expansa).

Newfoundland, biological control of Trialeurodes vaporariorum in, 310; Gilpinia polytoma apparently absent from, 37.

Nezara viridula, in Australia, 31, 170; food-plants and attempted introduction of parasite of, in Fiji, 170, 279, 493; possible vector of spike disease of sandal in India, 75; on cotton in Philippines, 649; in Tonga Is., 170; bionomics and control of, on vegetables, 31, 279; list of food-plants of, 152.

Nezara viridula var. smaragdula, 279.

Nicandra physaloides, Tarsonemus latus on, in S. Africa, 320.

nicolasi, Campylomma.
Nicotiana glutinosa, Thrips tabaci
transmitting virus disease to,
140.

Nicotiana tabacum (see Tobacco). Nicotine, against Aleurodids, 32, 132, 606; against Aphids, 132, 135, 145, 250, 281, 283, 287, 375, 384, 504, 556, 603, 631, 638; against Jassids, 117, 440, 497, 517, 558, 638; against other Rhynchota, 31, 121, 290, 603; against Cydia pomonella, 1, 22, 45, 67, 127, 128, 129, 160, 206, 239, 351, 411, 633, 634, 639, 645; against other Lepidoptera, 10, 32, 117, 132, 135, 162, 163, 290, 425, 448, 468, 558, 563, 633; against Diptera, 64, 384, 607; against mites, 32, 108, 109, 237, 321, 546, 627 637; against Monolepta varicornis, 600; against sawflies, 108, 233; against thrips, 25, 31, 121, 437; dusting with, 31, 32, 121, 135, 145, 162, 163, 233, 237, 290, 437, 448, 468, 504, 546, 600, 603; lime as carrier for, 121, 603; fumigation with, 132; applied by vaporisation, 250, 504, 631; plants dipped in, 283, 375, 384; aircraft for applying spray mixtures containing, 256; fixed compounds of, 22, 45, 117, 128, 129, 160, 162, 163, 206, 207, 233, 239, 256, 351, 352, 425, 556, 558, 563, 633, 634, 639, 644, 645; preparation of fixed compounds of, 162, 558, 634; tests of toxicity and action of various forms and compounds of, on insects, 206, 207, 353, 556, 644, 645; ovicidal

action of, 281, 352, 556; formulae containing, 25, 31, 32, 109, 121, 128, 145, 233, 237, 290, 321, 437, 497, 504, 546, 558, 563, 607, 684, 639; influence of chlorides on toxicity of, 556; and copper fungicides, 558; and creosote, 31; and derris, 135, 237, 287, 546; and lead arsenate, 634; and lime-sulphur, 108, 109; mixtures of naphthalene and, as a fumigant, 132; and oils, 1, 109, 127, 128, 160, 287, 351, 563, 606, 633, 634; factors affecting distribution of, between water and oils, 245; atomised sprays of oils containing, 250, 504; and, pyrethrum, 31; and rotenone, 250: Selocide compatible with, 560; and soap, 25, 233, 283, 375, 607; addition of other agents to sprays containing, 25, 160, 351, 556, 607; persistence of deposits and toxicity of, 206, 232, 233, 256, 351, 352; problem of residues of, on fruit, 45, 425, 634, 639; other insecticides compared with, 233, 250, 414.

Nicotine Sulphate, against Aphids, 54, 96, 127, 128, 136, 191, 250, 335, 359, 360, 381, 544, 596, 635, 650; against Coccids, 280, 632; against Jassids, 148, 291, 558, 639; against Psyllids, 534, 636, 639; against Trialeurodes vapovariorum, 32; against other Rhynchota, 82, 105, 121, 373, 492, 603; against Cydia pomonella, 45, 128, 160, 238, 239, 465, 634, 639; against other Lepidoptera, 117, 127, 128, 136, 190, 415, 447, 558, 594, 633; against Diptera, 55, 99, 170, 606, 636; injected into tunnels of Gryllotalpa spp., 237; against mites, 32, 100, 191, 237, 249, 546; against thrips, 204, 277, 413; unsatisfactory against Calandra oryzae in sacking, 541: dusting with, 54, 82, 136, 249, 250, 359, 596, 639; carriers for, 54, 136, 249, 250, 359, 596, 639; use of aircraft for applying sprays con-606, 632, 633, 634, 636, 639; and bentonites, 45, 160, 351, 352, 558, 634, 639; and Bordeaux mixture, 127, 148, 191, 381, 558; and derris, 136; and dinitro-ocyclohexylphenol, 249: lead arsenate, 335, 415, 606, 633, 634; and lime-sulphur, 100, 127, 238; and molasses, 636; addition of, to fixed nicotine sprays, 128; and oil emulsions, 32, 55, 127, 128, 160, 190, 191, 237, 238, 280, 291, 465, 534, 606, 632, 633, 634, 635; and soap, 82, 96, 121, 170, 277, 291, 373, 534, 544, 546, 594, 603, 635, 636; other wetters and adhesives for sprays containing, 117, 160, 277, 351, 352, 360, 413, 415, 595, 634, 639; persistance and duration of activity of deposits of, 233, 351, 352.

nidicola, Hylemyia.

niger, Apanteles obliquae; Thysanus.

Nigeria, pests of field and stored cacao in, 515, 516, 616; Helopeltis on cotton in, 515, 516; records of food-plants of insects in, 88.

Nightshade, Black (see Solanum nigrum).

nigra, Microphthalma; Saissetia. Nigra Scale (see Saissetia nigra). nigrescens, Eciton (Neivamyrmex); Mermis.

nigricana, Cydia (Laspeyresia). nigricans, Coccidaphycus. nigricella, Coleophora. nigricollis, Macrophylla.

nigricornis, Anagyrus (see A. fusciventris); Oecanthus; Valanga. nigricoxis, Syntomosphyrum zygaenarum.

nigripennis, Cryptogonus orbiculus.

nigripes, Thelaira. nigritus, Chilocorus; Coccophagus; Typophorus.

nigrivarius, Aphycus. nigrivenella, Mussidia. nigriventris, Anagrus armatus.

nigrofasciatus, Dysdercus. nigronervosa, Pentalonia.

nigropicta, Anomala. nigrovariegatus, Cirrospilus.

nigrovittata, Lema. Nilaparvata lugens (greeni), on rice in Ceylon, 594.

Nilaparvata oryzae, bionomics of, on rice in Japan, 98, 101.

niloticus, Coccidoxenus.

Nipa fruticans, Sexava on, in New Guinea Territory, 380. nipae, Pseudococcus.

Niphadolepis alianta, in Nyasaland, 59, 60, 61, 62, 154, 588; in Tanganyika, 514; on coffee, 59, 514; on tea, 59, 60, 61, 154, 588; on other plants, 59, 61, 62; bionomics of, 60, 588; parasites and disease of, 61, 154, 514, 588; measures against, 59, 61.

nipponicus, Diprion. Nisotra (see Podagrica).

nitens, Anaphoidea; Setora.

nitida, Cotinis; Erynnia; Telsimia.

nitidalis, Diaphania. nitidula, Anachaetopsis.

Nodaria extinctalis, on baled lucerne in S. Africa, 19.

nodiceps, Cryptognatha.

Nodonota puncticollis, bionomics and control of, in Pennsylvania, 192.

Nomadacris septemfasciata, in S. Africa, 20, 156, 612, 613; in Belgian Congo, 228; in Mozambique, 156; in Nyasaland, 154, 528, 589; in N. Rhodesia, 491; in Sudan, 388; in Uganda, 228, 388, 491, 492; bionomics and migrations of, 154, 228, 323, 589; crops attacked by, 228, 491; parasite of, 491; fungus infesting, 229, 589; measures and experiments against, 20, 154, 156, 589, 612, 613.

nordmannianae, Chermes.

normalis, Titanio.

Nornicotine, toxicity of forms of, to Aphis rumicis. 353.

Aphis rumicis, 353.
Norway Spruce (see Picea abies).
Nosema, probably infecting Tineola
biselliella in Switzerland, 376.

Nosema cactoblastis, Cactoblastis cactorum infected with, in S. Africa, 17.

notatus, Engytatus; Pissodes. notulatoria, Echthromorpha.

Nova Scotia, potato Aphids in, 302; Cydia nigricana and its biological control on pea in, 310, 311; forms of Rhagoletis pomonella in, 226.

Novius (see Rodolia). noxius, Tyrophagus. nubila, Sexava. nubilalis, Pyrausta.

nubilipennis, Acerophagus.

nucicola, Ephialtes (Calliephialtes); Microdus (Bassus).

nucum, Curculio (Balaninus). nuda, Pimpla (Ephialtes).

Nudaurelia cytherea, subspecies and strains of, in forests in S. Africa, 18. Nudaurelia cytherea capensis, 18.

Numidia meleagris, of little value against Sexava in New Guinea Territory, 380.

numidicus, Rhizotrogus. numismatica, Toumeyella. nunenmacheri, Scymnillus.

nursei, Goryphus (Melcha); Oncochalcis.

nüsslini, Dreyfusia (see Chermes nordmannianae).

Nyasaland, pests of Aleurites in, 60, 62, 588; cotton pests in, 60, 62, 153, 588; miscellaneous pests in, 30, 154, 589; tea pests in, 59, 60, 61, 154, 588; pests and diseases of tobacco in, 153, 154, 514, 588; locusts in, 154, 528, 589; beneficial insects in, 30, 62, 153, 154, 514, 588, 589.

Nygmia phaeorrhoea, tests of contact insecticides against, in Ger-

many, **232.**

nymphaeae, Rhopalosiphum.

Nymphalis antiopa, tests of lead arsenates against, in U.S.A., 119. Nysius coenosulus, experiments with virus disease of papaya and, in Hawaii, 35.

Nysius ericae, on beet in Arizona, 429.

Nysius natalensis, measures against, damaging turf in S. Africa, **615.** Nysius nemorivagus, food-plants of, in Hawaii, **651.**

Nysius vinitor, bionomics and control of, in Victoria, 31.

O.

Oak (Quercus), Lepidoptera on, in S. Africa, 19, 227; pests of, in Australia, 142; Aphids on, in Britain, 180; Lymantria dispar on, in China, 545; Acrocercops on, in Japan, 99; Tortrix viridana on, in Latvia, 86; Melolontha hippocastani on, in Russia, 572; pests of, in U.S.A., 119, 298, 551, 599; Cydia latiferreana on, in California, 86; (timber), Coleoptera in, 147, 178, 179, 233, 315, 316, 447, 593; termite damaging, 333.

Oak, Cork (see Quercus ilex var. suber).

Oak, Pin (see Quercus palustris).
Oats, Toxoptera graminum on, in
Argentina, 36; Halotydeus destructor on, in W. Australia, 141;
Cephus cinctus on, in Canada,
393; pests of, in India, 81, 266,
523; Criocerid experimentally
feeding on, in Japan, 93; pests
of, in Russian Union, 13, 14, 16,

572, 573, 574, 576, 617; pests of, in U.S.A., 251, 300, 356, 576, 428, 603; susceptibility of varieties of, to pests, 251, 393; insects and mosaic diseases of, 13, 16, 617; attraction of juices of, for Oscinella frit, 573.

Oats (Stored), pests in, and their control, 187, 210, 436, 576; reaction of Rhizopertha dominica to, 451; in bait for Thermobia, 406.

obesa, Pimpla (Ephialtes). oblineatus, Lygus pratensis.

obliqua, Diacrisia.

obliquae, Apanteles.

obliquus, Listroderes.

oblivia, Anomala (Pachystethus).

obovatus, Tenuipalpus.

obscura, Rhabdocnemis; Rodolia(Novius).

obscurata, Brachymeria; Euplectromorpha.

obscurella, Epilachna.

obscuriceps, Syntomosphyrum.

obscurus, Agriotes; Chrysomphalus; Hylastes (Hylastinus)

obsoleta, Heliothis (see H. armigera). obtectus, Bruchus (Acanthoscelides,

Bruchidius). obtusa, Agromyza.

occidentalis, Bittacus; Polybia;

Tortrix (Cacoecia). oceanica, Pheidole. oceanis, Dichomeris.

ocellana, Spilonota (Tmetocera).

ochraceus, Coccophagus.

Ochrogaster contraria, hand-collection of eggs of, on Acacia in Australia, 142.

Ochroma (Balsa), pests of, in Formosa, 103, 104; Xestobium rufovillosum not readily infesting wood of, 316.

ochus, Telenomus.

Ochyromera miwai, bionomics of, on persimmon in Formosa, 93.

Ocimum basilicum, Empoasca lybica on, in Sudan, 387.

Ocneria monacha (see Lymantria). octopilis, Eutermes (Nasutitermes). octosignatus, Brumus.

ocularis, Haptoncus.

Odontria, damaging pastures in New Zealand, 334.

Odontria zealandica, damaging pastures and crops in New Zealand, 333, 541; bionomics and control of, 541, 542.

Oecanthus nigricornis var. quadripunctatus, parasites of, on Daucus carota in U.S.A., 400.

Oecophylla smaragdina, measures against, on tea in Ceylon, 175; other ants preventing control of Amblypelta cocophaga by, Solomon Islands, 325.

Oecophylla smaragdina vivescens, attacking Promecotheca papuana, in New Guinea Territory, 378.

Oedaleonotus enigma, in California, 222.

Oenothera spp., relation of Psallus seriatus to, in Texas, 198.

Ohio, pests of beans and potato in, 39, 115, 248, 439; maize pests in, 43, 88, 253, 529; orchard pests in, 195, 248, 273, 425; Polychrosis viteana in, 425; Xyleborus germanus in, 551; beneficial insects and biological control in, 86, 114, 115, 425.

Oides decempunctata, on vines in

China, 446.

Emulsions, against Acridids, 241, 435; against Aleurodids, 32, 597, 606; against Aphids, 127, 128, 171, 191, 194, 248, 280, 284, 287, 360, 389, 494, 635; against Coccids, 23, 73, 102, 128, 155, 222, 223, 336, 349, 350, 426, 585, 597, 605, 632, 636, 642, 648, 652, 653, 654; against Psyllids, 232, 284, 534, 653; against other Rhynchota, 108, 216, 284, 373, 392, 440, 441, 603; against Calandra spp., 540, 576; against Cydia pomonella, 1, 127, 128, 129, 160, 238, 239, 279, 351, 411, 412, 465, 585, 633, 634; against other Lepidoptera, 108, 127, 128, 133, 134, 163, 190, 284, 307, 563, 585, 632, 633, 642, 644; against Diptera, 55, 131, 606; against mites, 23, 108, 109, 134, 191, 195, 237, 248, 281, 284, 285, 286, 287, 465, 559, 576, 597, 641, 653; against sawflies, 109, 285, 286; ineffective against *Thrips tabaci*, 121; tests of ovicidal action of, 216, 585; residual effect of, 74, 534; types of oils for, 32, 102, 108, 109, 127, 128, 129, 133, 134, 155, 171, 190, 191, 194, 195, 223, 232, 237, 238, 241, 245, 246, 248, 279, 280, 281, 284, 285, 286, 287, 307, 349, 350, 360, 373, 389, 392, 411, 426, 435, 441, 534, 540, 559, 563, 584, 585, 598, 603, 605, 606, 632, 633, 634, 635, 636, 541, 642, 644, 651, 652, 653, 654; emulsifiers and spreaders for, other than soap, 112, 113, 191, 241, 248, 350,

411, 426, 540, 559, 585, 605, 654; formulae for, 23, 102, 108, 127, 128, 133, 134, 191, 194, 216, 223, 232, 238, 241, 279, 280, 285, 307, 360, 435, 534, 540, 584, 585, 598, 603, 605, 606, 633, 636, 641, 642, 644, 652, 653, 654; formulae containing, 23, 32, 67, 108, 109, 127, 128, 133, 134, 190, 191, 195, 215, 223, 237, 238, 241, 248, 280, 281, 284, 285, 286, 291, 307, 349, 350, 435, 441, 534, 540, 559, 563, 605, 606, 632, 633, 634, 641, 642, 644 and Bordeaux mixture, 191, 441, 597, 634, 648; and calcium arsenate, 585; and cresylic acid, 128, 194; and cubé resins, 426; and cryolite, 134, 633; and derris, 240, 285, 286, 287, 349, 350; intermediary solvents for derris resins when used with, 349, 350; and dinitro-ortho-cresol, 108, 109, 128, 284, 605; and dinitroortho-cyclo-hexylphenol, **134, 191, 307, 605, 644**; inverted spray of salt of dinitro-o-cyclohexylphenol and, 561; and lead arsenate, 67, 127, 129, 133, 134, 279, 281, 351, 412, 606, 632, 633, 634; inverted mixtures of lead arsenate and, 163; and limesulphur, 23, 191, 223, 642; and nicotine, 1, 32, 55, 109, 127, 128, 160, 190, 191, 237, 238, 280, 287, 291, 351, 426, 465, 534, 563, 606, 632, 633, 634, 635; factors affecting distribution of nicotine in, 245; and oleic acid, 190; and oxydiphenyl, 576; and pyrethrum, 411, 642; and rotenone, 109, 122, 237, 350, 426, 642; and Selocide, 559; and sugar, 129; and tar distillates, 108, 128, 191, **194, 195, 248, 284, 585**; and tartar emetic, 215; and thiocyanates, 73, 108, 109; and zinc sulphate, 134, 633; electrolytes increasing deposits of, 411; injury to plants, 98, 222, 238, 280, 284, 351, 534, 584, 585, 633, 642, 652, 653; factors affecting action of, on leaves and trees, 652; methods of determining deposits of, on Citrus, 410, 411; effect of sprays of, on removal of spray residues from fruit, 412, 634; review of history of, as insecticides, 651.

Oil Palm (Elaeis guineënsis), coconut pests on, in New Guinea Territory, 378, 379; pests of, in Uganda, 175.

Oils, in baits for grasshoppers, 428; in sprays against Coleoptera in logs, 312, 427; and supplementary materials, treatment of maize ears with, against pests, 217, 469, 470, 481, 535, 640, 641; tests of, for treating water in rice-fields against Pyralid, 458; against permeability of termites, 6; insect cuticle by types of, 240; atomised sprays of insecticides in, 19, 230, 246, 250, 434, 435, 504; effect of type of, on atomised sprays, 398; phytotoxic properties of types of, 246; equipment for applying atomised sprays of, 654; aeroplanes for applying sprays of, 655; uses of, with insecticides applied from aircraft, 256, 412; dusts impregnated with, 161, 197, 353, 561, 584, 654; electrostatic charges of dusts impregnated with, 569; solution of resin in, as a spray adhesive, 419, 420; uses of sulphonated, in sprays, 360, 595, 637. (See Kerosene.)

Oils, Essential, in baits for Trypetids, 90, 94, 183, 594; unsatisfactory in baits for locusts, 612; attraction of, for Oscinella frit, 573.

Oils, Vegetable, in emulsion for treating turf against pests, 373, 374; in mixed sprays, 162, 353, 439, 634; uses of sulphonated, in sprays, 595, 637; and calcium cyanide, against Saperda candida, 243; permeability of insect cuticle by, 240; uses of, for conditioning dusts, 353, 504.

Oklahoma, pests of cereals and sorghum in, 213, 251, 258, 356; Lecanium corni in, 635; natural enemies of Toxoptera graminum in, 251; cowpea mosaic in, 502.

Okra (see Hibiscus esculentus).

Olea (see Olive)

Olea chrysophylla, Pseudococcus citri on, in Eritrea, **155.**

oleaceus, Exenterus (see E. adspersus).

oleae, Dacus; Parlatoria; Saissetia. Oleander, tests of distillates of, in baits for locusts, **612**.

Oleic Acid, uses of, in sprays, 163, 190; and caustic soda (see Sodium Oleate).

oleivorus, Phyllocoptruta (Phyllocoptes).

Olene achatina (see Dasychira).

Olenecamptus indianus, in timber in Burma, 78.

oleracea, Polia; Tipula.

Oligonychus ulmi, auct. (see Paratetranychus pilosus).

Olive, Parlatoria oleae on, in California, 629, 630; Teleonemia australis on, in Nyasaland, 589; Saissetia oleae on, in Peru, 511, 539; Dacus oleae on, in Spain, 462.

Olive Oil (see Oils, Vegetable). Olive Scale (see *Parlatoria oleae*). Omiodes accepta (see *Nacoleia*). omiodivorum, Microbracon.

Omocena convergens, on tea in Nyasaland, 154.

Oncideres, review of data on, in U.S.A., 213.

Oncocephala tuberculata, bionomics of, in India, 270.

Oncochalcis nursei, 74.

Oncometopia minor, on cotton in Peru, 510.

Oncopera, liberation of Hexamera against, in Victoria, 141.

Onion, Hylemyia antiqua on, in Britain, 374; Thrips tabaci on, in France, 25; pests of, in U.S.A., 121, 131; juice, etc., of, not improving spray against T. tabaci, 121.

Onion, Japanese, Micromyzus formosanus on, in Hawaii, 651.

Onion Fly (see *Hylemyia antiqua*). Onion Thrips (see *Thrips tabaci*). onobrychis, Macrosiphum.

Ontario, miscellaneous pests in, 303, 304; orchard pests in, 51, 225, 304, 305, 306, 548; insects and virus diseases of plants in, 306, 548, 646; Myopa parasitising honey bee in, 473; beneficial insects and biological control in, 225, 306, 308, 309, 310; Microplectron fuscipenne imported into U.S.A. from, 424.

onusta, Macronoctua.

Ocencyrtus johnsoni (egg-parasite of Murgantia histrionica), in U.S.A., 560, 651; importation of, into Hawaii, 651.

Ocencyrtus papilionis, parasite of Pyrilla perpusilla in Punjab, 82. Ocencyrtus pinicolus, parasite of

Dendrolimus in Sakhalin, 95. Oospora citri-aurantii, fruit-piercing moths associated with, on grapefruit in Gold Coast, 107.

Ootetrastichus dubius (parasite of Sexava), distribution of, 380.

Ootetrastichus megameli, sp. n., parasite of Megamelus proserpina in Philippines, 280.

opaca, Silpha (Aclypea).

Opatrum sabulosum, observations on, in Russia, 571, 573.

operculella, Gnorimoschema (Phtho-rimaea).

Operophtera brumata, in Austria, 232; in Britain, 80, 108, 281, 284; in Germany, 80; on birch in Latvia, 86; measures and tests of sprays against, on apple, 108, 232, 281, 284; parasitised by Apanteles praepotens, 80.

Operophtera fagata, parasitised by Apanteles praepotens in Germany,

80.

Opharus astur, on cacao in Brazil, 391.

Ophiomyia (see Agromyza).

Ophiostoma ulmi (causing Dutch elm disease in U.S.A.), relation of Scolytus multistriatus to, 116, 212, 364, 502, 563, 606, 644, 645; isolation of, from S. multistriatus stored at different temperatures, 166; relation of other Coleoptera to, 119, 365, 550, 644; list of insects associated with, 40; possibly spread by mites, 365; apple experimentally infected with, 119; measures against, 644.

opifera, Sarcophaga.

Opilo domesticus, predacious on Anobium punctatum in Germany, 234.

Opius, parasite of Rhagoletis in California, 565; parasite of Agromyzid on tea in Nyasaland, 588.

Opius fletcheri, attempted introduction of, into Loochoo Islands against Dacus cucurbitae, 99.

Opius humilis, suggested introduction of, into Spain against Ceratitis capitata, 459.

Opius ilicis (parasite of Phytomyza ilicis), bionomics of, in Britain, 489, 490; establishment of, in Br. Columbia, 417.

Opogona cleonyma, on Rhodes grass in Queensland, 466.

optabilis, Paranagrus.

Opuntia spp., establishment and utilisation of insects against, in S. Africa and Australia, 17, 18, 587; Arthropods attacking, in America, 587; fruit-piercing moths feeding on, in Gold Coast, 107; diseases of, 587; danger

of injury to spineless species of, by Lagochirus funestus, 13; in baits for cutworms, 615.

Opuntia aurantiaca, 17.

Opuntia monacantha, 17.

Opuntia tardiospina, 18.

opuntiae, Dactylopius; Tetranychus. Orange, Argyroploce leucotreta on, in S. Africa, 18; pests of, in Australia, 280, 335, 336; pests of, in Brazil, 167, 168, 392; Mellesis citri on, in China, 524; fruitpiercing moths damaging, in Gold Coast, 17, 107; Phytorus lineolatus on, in Guam, 332; pests of, in Spain, 462, 463; pests of, in Transcaucasia, 654; pests of, in U.S.A., 134, 209, 296, 297, 298, 475, 561, 597; fungi and ants associated with Pseudococcus comstocki on, 167, 168; risk of injury to, by oil sprays, 280; (fruit), cold storage of, against Trypetids, 544; juice of, in bait for Ceratitis capitata, 461, 462, 464.

Orange, Mandarin or Tangerine, Ceratitis capitata on, in Algerian Sahara, 616; Icerya on, in Argentina, 344; fruit-piercing moths damaging, in Gold Coast, 107; pests of, in Transcaucasia, 653; factors affecting action of oil emulsions on, 652.

Orange, Satsuma, pests of, in U.S.A., 597; lime-sulphur causing injury to, 598.

Orange Oil (see Oils, Essential). orbiculus, Cryptogonus; Exochomus. orbignyi, Bostrychopsis uncinata. Orchelimum, parasites of, on Daucus

carota in U.S.A., 400.

ordinarius, Apanteles.

Oregma lanigera, natural enemies and biological control of, on sugar-cane in Formosa, 92, 93.

Oregon, forest pests in, 49, 294, 371, 552; pests of fruit and nut trees in, 222, 261, 359, 415; Myzus lilii and virus diseases of Lilium candidum in, 45; Bruchus brachialis on vetches in, 48; Forficula auricularia in, 538; beneficial insects in, 294, 406, 552, 569.

oregonensis, Necremnus.

oregoni, Ips.

Oreodoxa regia, Brontispa longissima on, in New Guinea Territory, 379. Oreta carnea, bionomics of, on gambier in Sumatra, 105, 106; spraying against, 106.

Organothrips bianchii, gen. et sp. n., habits of, on taro in Hawaii, 273.

Orgilus longiceps, liberation of, against Cydia molesta in Connecticut, 161.

Orgyia mixta, food-plants of, in Nyasaland, **60, 61, 62.**

orichalcea, Plusia.

oriens, Prociphilus.

Oriental Fruit Moth (see Cydia molesta).

orientalis, Anomala; Aonidiella (Aspidiotus); Bruchocida.

oriolus, Exenterus.

Orius, lists of African and predacious species of, 320.

Orius insidiosus, predacious on Phyllocoptes fockeui in Ontario, 306; predacious on Heliothis armigera in Tennessee, 534.

Orius laevigatus, predacious on Phorodon humuli in Britain,

282.

Orius thripoborus, sp. n., predacious on thrips in Transvaal, 320.

orizicola, Sogata.

ornatus, Mesochorus; Myzus; Platyplectrus.

ornigis, Xenosternum.

Ornix prunivorella (see Parornix).
Orobanche, Myzus persicae on, in
India, 523.

Orphulella speciosa, introduced parasite of, in Canada, 309.

Orthocraspeda catenatus, new parasites of, in Celebes, **514.**

Orthocraspeda trima, new parasite of, in Java, 514.

orthocraspedae, Platyplectrus.

Orthodichlorobenzene, as a preservative for timber against *Lyctus*, 147; in sprays against Coleoptera in elm logs, 427; uses of emulsions containing, against pests in grassland, 111, 524, 525, 527; unsatisfactory as a soil fumigant against *Otiorrhynchus ligustici*, 198; causing injury to lucerne, 199; tests of volatilised mixture of naphthalene and, 555, 556.

orthogonia, Agrotis.

Orthol K, 633.

Oryctes rhinoceros (on coconut), search for natural enemies of, and their introduction into Samoa, 148.

oryctophaga, Scolia.

Calandra (Sitophilus); oryzae,.

Chlorops; Nilaparvata.

Oryzaephilus surinamensis (in stored products), in Australia, imported into Fiji, 496; U.S.A., 426; of little importance in cacao, 616; influence of environment and food on, 186; measures against, 380, 426; rearing and use of, for insecticide tests, 29, 591.

oryzella, Agromyza.

Oscinella (Oscinis) frit (on cereals in Russian Union), parasites of, 618; experiments on control of, 573, 574.

Othreis (Fruit-piercing Moths), observations on, in Gold Coast, 16. 106, 107, 492; in New Caledonia, fungi associated with damage by, 107.
Othreis fullonia (damaging Citrus

fruit), in Ceylon, 594; in Fiji, 389; in Gold Coast, 106, 107.
Othreis fullonica (see O. fullonia).
Othreis salaminia (see Eumaenas). Otiorrhynchus ligustici, measures against, on clover and lucerne in New York, 198, 252, 357, 643; soil conditions in relation to, 643. Otiorrhynchus ovatus, measures

against, on cranberry in Washington, 642.

Otiorrhynchus singularis, Campaea margaritata confused with, on apple in Britain, 282.

Otiorrhynchus sulcatus, measures against, on cranberry in Washington, 640, 641.

ovata, Brachymeria.

Otiorrhynchus (Brachyovatus, rrhinus).

Owls, destroying Polyphylla perversa, 509.

Oxo Bordeaux, 602.

Oxya chinensis, on sugar-cane and potato in Hawaii, 49, 650; utilisation and value of Scelio pembertoni against, 49.

Oxycanus (Porina), damaging grasses in New Zealand, 89, 333, 541; bionomics and control of,

Oxycanus cervinatus, bionomics and control of, in New Zealand, 541,

Oxydiphenyl, in formula for spraying granaries against pests, 576.

Oxyptilus periscelidactylus, bionomics and control of, on vines in Massachusetts, 133.

Oyster-shell Scale (see Lepidosaphes ulmi).

P.

Pachyceras eccoptogastri (see Rhoptrocerus)

Pachydissus birmanicus, bionomics of, on Xylia dolabriformis in Burma, 77.

Pachylia ficus, measures against, on Ficus spp. in Brazil, 392.

Pachymerus gonager, bionomics of, in stored tamarind pods in India, 523.

Pachyneuron altiscuta, parasite of Pseudococcus comstocki in Virginia, 196.

Pachyneuron coccorum, parasite of Pseudococcus comstocki in Palestine, 339.

Pachyneuron nawai, parasite Dendrolimus in Sakhalin, 95.

Pachyneuron pentatomivora, parasite of Urostylis punctigera in India, 75.

Pachyosa perplexa, on Citrus and balsa in Formosa, 103.

Pachyrrhynchus moniliferus, intercepted in cacao in Formosa, 103.

Pachystethus (see Anomala). Pachyzancla (see Psara). pacificus, Tetranychus.

packardi, Cydia; Melanoplus. padellus, Hyponomeuta.

padi, Anuraphis. pagana, Tipula.

Pagiophloeus longiclavis, bionomics and control of, in India, 75.

Pagyda traducalis, on Zizyphus jujuba in Siam, 94.

Palaquim gutta, Apogonia cribricollis on, in Malaya, 512.

Pales maculata (see Nephrotoma). Pales pavida (see Neopales).

Palestine, Pseudococcus comstocki and its natural enemies on Citrus in, 73, 74, 339; miscellaneous pests in, 403, 622, 623; Chalcidoid parasite in, 514.

pallida, Trichogramma. pallidifer, Creontiades.

pallidus, Creontiades; Tarsonemus. pallipes, auct., Agonoderus (see A. lecontei).

palmarum, Bitoma; Mystrops: Rhynchophorus.

Palm, Sago (see Metroxylon sagu). Palm Weevil, Black (see Rhynchophorus papuanus).

Palmetto, Bermuda (see Sabal bermudiana).

Palmodes laeviventris, predacious on Anabrus simplex in Montana,

Palms, Coccotrypes dactyliperda on, in Bermuda, 139; pests of, in Brazil, 9, 10, 468; Dinapate wrighti on, in California, 137; Nephantis serinopa on, in Ceylon, 619; Rhynchophorus palmarum on, in Colombia, 9; Aeolarchis sphenotoma on, in Fiji, 496; pests of, in New Guinea Territory, 378, 379, 380; Pseudococcus nipae on, in Porto Rico, 472; Melittomma insulare on, in Seychelles, 341. (See Coconut, Date and Oil Palms.)

paludosa, Tipula.

Pamphilius sylvaticus, rearing of, on plum in Germany, 339.

Pandanus, Aeolarchis sphenotoma on, in Fiji, 496.

Pandemis (see Tortrix).

Pandesma quenavadi, parasites of, on Acacia catechu in India, 263. panicea, Sitodrepa.

Panicum barbinode, Lepidoptera on,

in Fiji, **148, 495.**

crus-galli, Delphacodes Panicum striatellus and mosaic of cereals in, in Siberia, 14.

Panicum dichotomiflorum, Chaetocnema denticulata on, in U.S.A.,

Panicum miliaceum, Pyrausta nubilalis on, in Russia, 488; attraction for Oscinella frit of juices of, 573.

Paniscus testaceus, parasite of Dendrolimus in Sakhalin, 95.

Panolis flammea, unimportant on pine in Latvia, 86.

pantherinus, Xanthippus corallipes. Pantomorus godmani, in New Zea-

land, 545; review of data on, 545.

Pantomorus leucoloma, in Chile,
428; in New South Wales, 494, 609; in U.S.A., 208, 209, 370, plants 428, 494, 568, 632; attacked by, 428, 494; bionomics of, 494, 609; Neoaplectana glaseri parasitic in, 370; measures and legislation against, 208, 428, 568.

Pantomorus peregrinus, measures and legislation against, in U.S.A.,

Pantomorus taeniatulus, bionomics and distribution of, 608.

Pantophthalmus spp., bionomics of, on Artocarpus in Brazil, 392.

Papaipema purpurifascia, ovicides against, on Aquilegia in New York, 644.

Papaya, Rhynchophorus palmarum on, in Colombia, 9; Coccid on, in Dominican Republic, 498; Dacus ferrugineus on, in Malaya, 90; pests and disease of, in Hawaii, 35, 650, 651; (fruit), fumigation against introduction of Trypetids into U.S.A. from Hawaii in, 499.

Paper, constituents of, palatable to Ctenolepisma longicaudata, 56.

Paper Covers, types of, for protecting bananas against Scirtothrips signipennis in Queensland, 436, 465; for protecting peaches from Ceratitis capitata, 464.

Papilio schmeltzi, on Citrus in Fiji,

389.

papilionis, Ooencyrtus.

Pappea, Achaea lienardi on, in S. Africa, 19.

Papuana, distribution of, on Colocasia esculentum in Pacific Is.,

papuana, Promecotheca.

papuanus, Rhynchophorus. Para Grass (see Panicum barbinode). Parabrachycoma ruficauda, gen. et sp. n., parasite of Alabama argillacea in Argentina, 169.

paraclina, Porthmologa.

Paradexodes epilachnae (parasite of Epilachna spp.), in Mexico, 114; bionomics and attempted establishment of, in U.S.A., 114, 115.

Paradichlorobenzene, against borers in trees, 78, 160, 243; against clothes moths and carpet beetles, 363; against Ctenolepisma longicaudata, 56; as a soil fumigant, 110, 145, 278, 279, 280, 614; ineffective for treating tobacco plant-beds against Epitrix parvula, 404; causing injury to tomato, 145; solutions and emulsions containing, 78, 110, 243, 579.

paradoxa, Cedria. Paraffin Wax, uses of, in sprays, 112, 113, 134.

Paraguay, Conotrachelus denieri on cotton in, 445.

Paralitomastix varicornis, parasite of Dichomeris eridantis in India, 74.

parallelipipedus, Lyctus. parallelus, Hypophloeus. Paranagrus optabilis, apparently parasitic on Perkinsiella thompsoni in Guam, 331.

paranensis, Schistocerca.

Paranomalon (see Anomalon).

Paraphasiana dysderci, gen. et sp. n., parasite of Dysdercus spp. in Brazil, **469.**

Paraphorantha peruviana, parasite of Dysdercus ruficollis in Peru,

510.

Pararcyptera microptera, effect of climate on outbreaks of, Siberia, 485: natural enemies of, 485.

Parasa johannes, on Quercus robur in S. Africa, 19.

Parasa latistriga, on tea in Nyasa-

land, 154.

Parasa lapida, new parasite of, in Malaya, 514.

Parasa vivida, bionomics of, in

Nyasaland, 60.

Parasites, search for, in artificial environments, 362; problems of host-finding by, 397, 490; factors preventing establishment of, 219. Parasymmictus clausa (see Trichop-

sidea).

Paratetranychus, on date palm in Sudan, 388; on oaks in U.S.A.,

Paratetranychus bioculatus, relation of shade to infestation of tea by, in Nyasaland, 61.

Paratetranychus citri, measures and experiments against, on Citrus

in U.S.A., 561, 597.

Paratetranychus indicus, on sugarcane and Sorghum spp. in Punjab,

271, 520, 521; bionomics of, 521. Paratetranychus mangiferus, sp. n., food-plants of, in Punjab, 271.

Paratetranychus pilosus (on fruittrees), in Britain, 108, 281, 284, 285, 286; in Br. Columbia, 22; in U.S.A., 2, 134, 191, 195, 248, 559, 561; bionomics of, 108, 281, 285, 286; measures against, 22, 108, 109, 134, 191, 195, 248, 281, 284, 285, 286, 559, 561; technique for testing ovicides against, 195.

Paratetranychus ununguis, 599.

parathea, Empoasca.

Paratheresia diatraeae (see Theresia claripalpis).

Paratrechina longicornis, infesting food in Connecticut, 164.

Paratrioza cockerelli (Potato Psyllid), in Canada, 330; in U.S.A., 21, 278, 537, 596, 638; on tomato, 21, 278; causing Psyllid yellows, 21, 278, 596, 638; bionomics of, 21, 278; measures against, 21, 278, 638, 639.

pardalina, Locustana.

Parechthrodryinus clavicornis, parasite of Laccifer lacca in India, 263. Parendymia pilipes, in mangrove timber in Fiji, 170.

Paria canella, on strawberry in

Montana, 537.

Paris Green, in baits, 138, 283, 300, 354, 355, 357, 542, 612, 613; dusting with, 91, 92, 208, 325, 326; carriers for, 92, 325, 326; aeroplane dusting with, 300; spraying with, 10, 136, 208, 277, 325, 326, 392, 413, 468, 479, 624; uses of, against termites, 6, 7, 387, tests of toxicity of, to 589; insects, 208, 481; danger of, to bees, 136; and Bordeaux mixture, 10, 325, 326; and calcium arsenate, 300; and lime, 325, 326, 479; and sugar, 129, 277, 413; and injury to plants, 91, 208, 277, 326; other insecticides compared with, 91, 92, 277, 325, 326, 357, 413, 642.

Parkia speciosa, pests of, in Java,

182.

Parlatoria oleae, outbreak, bionomics and distribution of, in U.S.A., 222, 629, 630; characters of, 629.

Parornix prunivorella, bionomics and control of, on fruit trees in

Kansas, 190, 191.

Parsley, Lachnosterna crinita damaging, in Texas, 214.

Parsnip, Listroderes obliquus on, in Victoria, 31. partenopea, Encarsia.

Parthenocissus quinquefolia, Scirtothrips citri on, in California, 565. parummaculata, Conchyloctenia.

parvipennis, Dasyscapus.

parvula, Epitrix.

parvulus, Euplectrus; Longitarsus; Pleurotropis.

Pasania (see Lithocarpus).

Passiflora, Leptoglossus gonagra on, in Argentina, 512.

Passiflora edulis, pests of, in Australia, 144, 465.

passiflorae, Dacus (Chaetodacus.

Strumeta). Passion Vine (see Passiflora edulis). Patanga succincta, outbreaks of, in

India, 519. patruelis, Lepinotus.

pauciseta, Sciara.

paucispinosa, Frankliniella.

pauper, Dialeges. pavida, Neopales (Pales).

zeylanica, Pempherulus Pavonia affinis on, in Madras, 521, 522.

pawneepae, Aphis.

Pea, pests of, in Australia, 31, 142, Cydia nigricana on, in 236; Britain, 310; C. nigricana on, in Canada, 22, 310; Bemisia tabaci on, in India, 523; pests of, in Russia, 575, 576; pests of, in U.S.A., 42, 136, 158, 164, 202, 249, 353, 376, 408, 428, 504, 630, 631, 639, 640; stage of development affecting recovery of, from Macrosiphum onobrychis, nutritional experiments with Pyrausta nubilalis and, 43; not harbouring Aphid vectors of potato virus diseases, 165; cryolites and injury to, 260, 428; (stored), Bruchids and their control in, 408, 492, 630.

Pea, Chick (see Cicer arietinum). Pea, Pigeon (see Cajanus cajan). Pea Aphis (see Macrosiphum onobrychis).

Pea Bruchid (see Bruchus pisorum). Pea Moth (see Cydia nigricana).

Peach, Ceratitis capitata on, in Algerian Sahara, 616; pests of, in Australia, 55, 141, 144, 149, 335, 465; Myzus persicae on, in Brittany, 23; Malacosoma neustria on, in China, 446; Neurotoma nemoralis on, in Germany, 233; pests of, in India, 267, 518, 523; Epidiaspis leperii on, in Italy, 155; pests of, in Japan, 372, 622; Hyalopterus arundinis on, in Jugoslavia, 72; Cydia molesta on, in Korea, 372; Myzus persicae on, in Manchuria, 100; pests of, in Ontario, 225, 306, 548; pests of, in Peru, 511, 539: Ceratitis capitata on, in Spain, 464; pests of, in U.S.A., 2, 41, 45, 87, 113, 118, 129, 130, 159, 161, 192, 196, 210, 248, 348, 403, 405, 418, 425, 557, 602; spray schedule for, in Kentucky, 2; insects and virus diseases of, 306, 403, 548; insecticides causing injury to, 55, 197; (fruit), fumigation of, against *Anarsia* lineatella, 113; juice of, in bait for Ceratitis capitata, 461, 462.

Peach Aphis, Green (see Myzus

persicae).

Peach Borer (see Aegeria exitiosa). Peach Twig Borer (see Anarsia lineatella).

Pear, pests of, in S. Africa, 18, 435; pests of, in Australia, 279, 291; injury to, by Tomaspis sanguinea in Britain, 282; pests of, in Br. Columbia, 22, 508; Dacus ferrugineus on, in Ceylon, 594; Illiberis pruni on, in China, 446: Anthonomus piri on, in Germany, 617; pests of, in India, 183, 518; Epidiaspis leperii on, in Italy, 155; pests of, in Japan, 96, 99, 372; Cydia pomonella on, in Jugoslavia, 72; pests of, in Korea, 171, 372; Aspidiotus perniciosus on, in Russian Union, 653; pests of, in Switzerland, 66, 617; pests of, in U.S.A., 48, 113, 190, 192, 211, 222, 223, 243, 261, 359, 409, 425, 533, 534; spur-bull on, 359; Rhagoletis pomonella developing on, 226; bees transmitting fire-blight to, 112; (fruit), fumigation of, against pests, 18, 113, 222.

Pear Psyllid (see Psylla pyricola). Pear Thrips (see Taeniothrips in-

consequens).

Pecan, pests of, in U.S.A., 116, 117, 133, 159, 166, 192, 211, 222, 426, 635; Acrobasis caryae completing development on foliage of, 117; insecticides causing injury to, 222, 426.

Pecan Aphis, Black (see Melanocallis caryaefoliae).

Pecan Nut Casebearer (see Acrobasis caryae).

pectinicornella, Mussidia.

Pectinophora gossypiella (see Platyedra).

pectoralis, Arge.

pedalis, Pimpla (Ephialtes).

pedatoria, Eupelmella.

pedester, Hadronotus.

pedestris, Podisma.

pedicellata, Scutiphora.

Pediculoides ventricosus, attacking Nephantis serinopa in India, 620; attacking Bruchus pisorum in Russia, 575; attacking Tineola biselliella in Switzerland, 376: bionomics of, destroying cultures of Bruchids in U.S.A., 215.

Pedinus femoralis, habits of, on wheat in Russia, 571; studies on resistance of, to cold, 573.

pela, Ericerus. Pelamia latipes (see Mocis). pellucida, Camnula. pellucidus, Longitarsus.

Pelochyta avontes, on cacao in Brazil,

pembertoni, Scelio.

Pempheres affinis (see Pempherulus). pempheridis, Entedon.

pempheriphila, Euderus. Pempherulus, gen. nov., 265.

Pempherulus affinis (Cotton Stem Weevil), in India, 268, 270, 515, 521, 522; measures against, in Philippines, 649; relations of parasites to food-plants of, 270, 521, 522; possibility of utilising Spathius vulnificus against, 268; effects of temperature and humid-

ity on, 515; type of genus, 265. Pempherulus pterospermi, system-

atic position of, 265.

Pemphigus bursarius, irrigation against, on lettuce in Montana, 537.

Penicillium, susceptibility of pineapple varieties to, in Hawaii, associated with Scolytus multistriatus on elm in W. Virginia, 116.

Penicillium digitatum, fruit-piercing moths associated with, on grapefruit in Gold Coast, 107.

pennipes, Trichopoda.

Pennisetum, Laphygma exempta on, in Nyasaland, 154.

Pennisetum macrostacium, Sexava experimentally feeding on, 380.

Pennisetum typhoideum, new Jassid on, in India, 266; preferred by Schistocerca gregaria, 519.

Pennsylvania, cereal pests in, 48, 400; orchard pests in, 129, 191, 192, 371; Lymantria dispar in, 213, 428; vegetable pests in, Plagiodera versicolora on 247; willow in, 438; beneficial insects and biological control in 87, 129, 192, 400; utilisation of milky diseases against Popillia japonica in, 124.

pennsylvanica, Epicauta.

Pentachlorphenol, as a preservative for timber against Lyctus, 448.

Pentachrysis (see Chrysis).

suavis, resistance Pentacme of barked logs of, to borers in Burma, 78.

pentagona, Aulacaspis.

Pentalonia nigronervosa, relation of, to virus diseases of Musa spp. in Australia and Philippines, 57, 149, 317.

pentatomivora, Pachyneuron.

Pentatrichopus, considered distinct from Capitophorus (q.v.), 27.

Penthaleus major, food-plants and control of, in Australia, 236, 546.

Pentilia castanea (predacious on Coccids), distribution and utilisation of, in W. Indies, 471, 498; into Florida introduced Hawaii. 472.

Pentilia egena (predacious on Coccids), in Brazil, 471, 625; distribution and utilisation of, in W.

Indies, **263, 471.**peponis, Plusia.
Pepper (see Capsicum and Piper nigrum).

Pepper Tree (see Schinus molle). Pepper Weevil, Small (see Lophobaris piperis).

Peppermint, reaction of Rhizopertha dominica to, 452.

perdix, Rhinaria.

Peregrinus maidis, transmitting stripe disease of maize in Porto Rico. 469.

peregrinus, Pantomorus; Tetracnemus.

perforans, Xyleborus.

Perilampus, parasite of Ascogaster carpocapsae in U.S.A., 193.

Perilampus hyalinus, hyperparasite of grasshoppers in Canada, 309. Perilitus basalis (see Microctonus). Perilla Oil (see Oils, Vegetable).

Periodical Cicada (see Magicicada septendecim).

Periplaneta americana, insecticide tests on, 352, 353, 556.

Periplaneta australasiae, in greenhouses in Britain, 489.

periscelidactylus, Oxyptilus.

Perisierola, parasite of Tortrix sarcostega in Baluchistan, 518; bionomics of, parasitising Nephantis serinopa in Ceylon, 620, 621; possibly parasitic on Kearfottia in Porto Rico, 472.

Perisierola angulata, importation and colonisation of, against Cydia

molesta in U.S.A., 295.

Perisierola nephantidis, parasite of Nephantis serinopa in India, 620. Perkinsiella thompsoni, parasite of, on sugar-cane in Guam, 331.

perniciosus, Aspidiotus (Quadraspidiotus); Pseudococcus.

perobtusa, Batrachedra.

Peromyscus, destroying egg-pods of Anabrus simplex in California, 223.

Peronea comariana, on strawberry in Britain, 28.

Peronia rostrata, Dacus confused with, in New South Wales, 237. perpilosa, Formica.

perplexa, Pachyosa. perpusilla, Pyrilla.

Persectania composita (ewingi), damaging pastures in New Zealand, 334.

persicae, Lecanium; Myzus (Aphis, Myzodes).

Persimmon, Ochyromera miwai on, in Formosa, 93; Drosophila suzukii on, in Japan, 622.

persona, Xyleutes.

personata, Phaonia (see Peronia rostrata).

Peru, cotton pests in, 11, 40, 404, 509-511, 539; miscellaneous pests in, 139, 511, 539; beneficial insects and biological control in, 510, 511; danger of introduction of Platyedra gossypiella into, 539.

peruella, Mescinia.

peruviana, Acaulona; Paraphorantha.

peruvianus, Dysdercus. perversa, Polyphylla. petiolatus, Euplectrus.

Petrol, for destroying termite nests, **6.** (See Gasoline.)

Petrova albicapitana (see Rhyacionia).

Petunia, Thrips tabaci transmitting virus disease to, 140.

Pexopsis pyrrhaspis, parasite of Lamellicorns in Zanzibar and Natal, 83, 609; imported into Mauritius against Lachnosterna smithi, 609.

Pezomachus (see Gelis).

Phacelia, Scoliids attracted to, in Russia. 572.

Phaenacantha marcida, bionomics and control of, on sugar-cane in Formosa, 104, 105.

Phaenocarpa leveri, parasite of Drosophila in Fiji, 496.

Phaenopria, parasite of Drosophila suzukii in Japan, 622.

Phaeogenes ater, parasite of Aegeria exitiosa in Ontario, 226.

phaeorrhoea, Nygmia.

Phalaenoides glycinae, measures against, on vines in New Zealand, 333.

Phalera flavescens, on apple and cherry in Japan, 96.

phalerata, Campsomeris; Mylabris. phanaea, Homona.

Phaneroptera brevis, on tobacco in Sumatra, 318.

Phaneroptera furcifera, on cotton in Philippines, 649.

Phanerotoma grapholithae (parasite of Cydia molesta), in Japan and Korea, **372**; importation and colonisation of, in U.S.A., **295**.

Phanerotoma hendecasiella, parasite of Lepidoptera in India, 74, 75, 183.

Phanerotoma planifrons (parasite of Cydia glycinivorella), bionomics of, in Manchuria, 98.

Phanerotoma tibialis, parasite of Cydia spp. in U.S.A., 193, 406.

Phanomeris spectabilis (see Rhogas dendrolimi).

Phanurus (see Telenomus).

Phaonia personata (see Peronia rostrata).

Phaseolus (see Beans).

Phaseolus lunatus (see Lima Beans). Phaseolus mungo (Mungo Bean), Araecerus fasciculatus in stored, in Venezuela, 293.

phasiana, Anoplocnemis.

Pheasants, little danger to, from grasshopper baits, 205.

Pheidole, destroying Isaniris ater in Nyasaland, 62.

Pheidole javana var. dolenda, destroying Phaenacantha marcida in Formosa, 105.

Pheidole megacephala, destroying Promecotheca papuana in New Guinea Territory, 378; measures against, in tobacco seed-beds in Nyasaland, 588.

Pheidole oceanica, attacking antspredacious on Amblypelta in Solomon Islands, 325.

Pheidole punctulata, damaging turf in S. Africa, 615.

Pheletes ectypus, damaging potato in New York, 643, 644, 645; measures against, 643; characters of, 645.

Phellinus cryptarum, relation of Xestobium rufovillosum to, in oak timber in Britain, 178, 315.

Phellodendron, insecticidal properties of fruit of, 414.

Phenacoccus gossypii, on rose in U.S.A., 256.

Phenacoccus hirsutus, measures against, on roots of leguminous

plants in Egypt, 30; uncommon on roots of cotton, 30.

Phenacoccus icervoides, on cacao in Malaya, 513.

Phenacoccus insolitus, not experimentally transmitting virus disease to tomato in Madras, 542.

Phenol (see Carbolic Acid).

Phenomenal Berry, Metallus albipes on, in Britain, 544.

Phenothiazine (see Thiodiphenylamine).

Philaenus leucophthalmus, Phyllocoptes fockeui spread by, in Ontario, 306.

Philippines, cotton pests in, 648, 649; Coleoptera on cucurbits and Flacourtia indica in, 649; Aphids and virus diseases of Musa textilis and Canna indica in, 57, 317; new parasite of Megamelus proserpina in, 280; insects introduced into Formosa from, 92, 103.

phillyreae, Siphoninus.

Phlegethontius (see Protoparce).

Phleum pratense, Septis finitima on, in Maine, 560; resistance of, to Cephus cinctus, 393.

Phlugiola dahlemica, 489.

Phlugiolopsis henryi, gen. et sp. n., in hothouse in Britain, 489; feeding habits and suggested utilisation of, 489.

Phlyctaenia rubigalis, laboratory

rearing of, 557.

Phoenix, Pseudococcus brevipes on roots of, in Egypt, 30. (See Date

Phoracantha semipunctata, on Eucalyptus in Argentina, 512.

Phorocera (parasite of Lepidoptera), in Colombia, 53; in Oregon, 406; in Tasmania, 151.

Phorocera claripennis, parasite of Datana integerrima in Missouri, 167.

Phorocera erecta, 406.

Phorodon humuli, predacious enemies of, on hops in Britain, 282.

Photinia serrulata, Scirtothrips citri on, in California, 565.

Phragmites communis, Locusta migratoria associated with, in Russia, 570.

Phrydiuchus topiarius, type of injury to Salvia sclarea by, in Crimea, 15.

Phryxe (see Zenillia).

Phthalic Glyceryl Alkyd Resin, experiments with, against mites, 535, 536.

phthisicus, Mischocyttarus.

Phthorimaea (see Gnorimoschema).

phthorimaeae, Euplectrus. phycidis, Habrocytus.

phycitis, Eutettix.

Phygadeuon, parasite of Dendrolimus in Sakhalin, 95.

Phygadeuon subfuscus, parasite of Paradexodes epilachnae in Ohio, 115.

Phylacteophaga eucalypti, bionomics of, in New South Wales, 237.

Phylaitis pterospermi (see Pempherulus).

scutellaris (see Chiro-Phylaitis zetes).

Phyllanthus discoideus, Achaea spp. on, in Gold Coast, 107.

Phyllocoptes destructor, sp. measures against, on tomato in California, 565.

Phyllocoptes fockeui, bionomics and control of, on plum in Ontario, 304-306; food-plants and distribution of, 304.

Phyllocoptes lycopersici, on tomato in Australia, 32, 565; measures against, 32; characters distinguishing P. destructor and, 565.

Phyllocoptes oleivorus (see Phyllocoptruta).

Phyllocoptruta oleivorus (on Citrus), in Argentina, 263; in Nyasaland, 589; in Queensland, 480; in Trinidad, 648; in U.S.A., 135, 597; measures against, 135, 597, 648.

Phyllodecta spp., effect of insecticides on, hibernating on apple in Germany, 70.

Phyllodes consobrina, damaging Citrus fruit in Ceylon, 594.

Phyllognathus, parasite of, on cork oak in Morocco, 83.

Phyllonorycter (see Lithocolletis). Phyllopertha (see Anomala).

Phyllophaga (see Lachnosterna).

Phyllotoma nemorata, on birch in New Brunswick, 348.

Phyllotreta atra, on cabbage in Germany, 232.

Phylloxera vitifoliae (on vines), in S. Africa, 320; in Bermuda, 139; in France, 111; in Germany, 231; studies on, in Russian Union, 578-580; measures against, 578, 579, 580.

Phymateus leprosus, bionomics and control of, in S. Africa, 435.

Phymatioderus bizonatus, on Eucalyptus in Chile, 512.

Physalis angulata, Heliothis assulta on, in Sumatra, 319; reservoir of tobacco mosaic, 319.

Phytalus smithi (see Lachnosterna). Phytodietus celsissimus, parasite of Tortrix postvittana in Tasmania,

Phytolyma lata, parasites of, on Chlorophora excelsa in Gold Coast,

phytolymae, Psyllaephagus.

Phytometra (see Plusia).

Phytomyptera leucoptera, parasite of Tortrix cerasivorana in Quebec, 626.

Phytomyza atricornis, on flax in Formosa, 93; on chrysanthemum in Scotland, 381.

Phytomyza ilicicola, bionomics and control of, on holly in U.S.A., 606.

Phytomyza ilicis (on Ilex aquifolium), in Britain, 417, 489, 490; in Br. Columbia, 417; parasites and biological control of, 417, 489, 490.

Phytomyza rufipes, on rape in Germany, 235.

Phytomyza spicata, parasites of, on maize in Fiji and Guam, 170, 331, 496; on Sorghum and sugarcane, 170; spray against, 170.

Phytonomus (see Hypera).

Phytophaga (see Mayetiola). Phytorus lineolatus, on mango etc. in Guam, 332.

hytorus pinguis, recorded as, 332. P. Phytorus lineolatus

Phytotribus platyrhinus, sp. n., on Cocos coronata in Brazil, 468.

Picea (see Spruce).

Picea abies, pests of, in Canada, 37, 509; pests of, in U.S.A., 422, 537, 607.

Picea albertiana, Chermes cooleyi on, in Montana, 537.

Picea engelmanni, Chermes spp. on, in Montana, 537.

Picea excelsa (see P. abies).

Picea glauca, pests of, in Canada, 37, 38, 532.

Picea jezoensis, new Rhynchites on, in Japan, 40.

Picea mariana, pests of, in Canada, 37, 532.

Picea rubra, Gilpinia polytoma on, in Quebec, 37.

Picea sitchensis, decay not rendering timber of, suitable for Xestobium, **315.**

piceaperda, Dendroctonus.

piceus, Attagenus; Hylobius (Hypomolyx).

picipes, Rhabdopterus; Tenebrio.

picitarsis, Ceuthorrhynchus.

Picromerus bidens, doubtful value of, against Leptinotarsa decemlineata in Germany, 70.

picta, Dexiomorpha; Zenillia (Exorista).

Pieris, pyrethrum sprays against,

on cabbage, 585.

Pieris brassicae, in Britain, 314; new spirochaete in, in France, 224: not occurring in New Zealand, 314; importance of, as host of Apanteles glomeratus, 314: effects on Tettigoniids of feeding on larvae of, 489; experiments with bacteria against, 583, 584; test of insecticide on, 584.

Pieris rapae (on crucifers), introduction and spread of, in Australia, 290, 436, 467, 546; in Britain, 145, 313; in Hawaii, 34, 650; in Japan, 173; in New Zealand, 333; in Russia, 654; in U.S.A., 199; bionomics of, 313, 546; natural enemies and biological control of, 34, 313, 314, 333, 468, 546, 650; diseases of, 173, 314, 650; measures against, 145, 199, 290, 468; tests of insecticides on, 242, 654; laboratory rearing of, 557.

Piezosternum fallax, on cacao and Solanum in Nigeria, 516.

Pigeon Pea (see Cajanus cajan). Pigs. suggested utilisation against Polyphylla perversa, 509. Pikonema alaskensis, on spruce in Canada, 36, 530; against, 36. measures

Pikonema dimmocki, on spruce in Quebec, 36.

pilipes, Parendymia.

pilistriatus, Circopes.

pilleriana, Sparganothis.

pilosa, Anoxia.

pilosum, Sparasion. pilosus, Paratetranychus.

Pimpla, application of name, 220. Pimpla aequalis, parasite of Cydia

spp. in U.S.A., 220. Pimpla conquisitor, parasite of Tortricids in Canada, 415, 626.

Pimpla disparis, parasite of Dendrolimus in Sakhalin, 95.

Pimpla evetriae, parasite of Rhyacionia buoliana in Br. Columbia, 415.

Pimpla inflata, sp. n., parasite of Cydia spp. in N. America, 220.

Pimpla instigator, parasite of Dendrolimus in Sakhalin, 95; type of genus, 220.

Pimpla montana, hosts and utilisation of, in U.S.A., **552.**

Pimpla nuda, parasite of Neodiprion in Maine, 638; reared on Gilpinia polytoma, 638.

Pimpla obesa, parasite of Rhyacionia buoliana in Br. Columbia, 415.

Pimpla pedalis, parasite of sawflies in U.S.A. 637, 638.

Pimpla pomorum, parasite of Anthonomus pomorum in Germany, 70.

Pimpla roborator (parasite of Margaronia pyloalis), bionomics of, in Japan, 101.

Pimplini, of N. America, 220.

pinastri, Sphinx.

Pine (Pinus), Euproctis terminalis on, in S. Africa, 18; pests of, in Britain, 323, 382, 383, 593; pests of, in Canada, 37, 312, 415; Dendrolimus punctatus on, in China, 545; Sipalus formosanus on, in Formosa, 104; pests of, in Japan, 480; pests of, in Latvia, 85, 86; Sphinx pinastri on, in Poland, 86; Dendrolimus on, in Sakhalin, 95; pests of, in U.S.A., 48, 49, 87, 116, 164, 211, 357, 371, 396, 400, 422, 426, 427, 507, 551, 563, 605, 607, 637, 638, 648; fungi associated with Coleoptera in, 357; effect of felling on incidence of pests on, 382, 593; pests in timber of, 138, 234.

Pine, Black (see *Pinus nigra*). Pine, Grey or Jack (see *Pinus bank*-

siana).

Pine, Hoop (see Araucaria cunninghami).

Pine, Japanese Black (see Pinus thunbergi).

Pine, Japanese Table (see Pinus densiflora umbraculifera).

Pine, Jeffrey (see Pinus jeffreyi).

Pine, Limber (see Pinus flexilis).

Pine, Loblolly (see *Pinus taeda*). Pine, Lodgepole (see *Pinus contorta*).

Pine, Pitch (see *Pinis rigida*). Pine, Red (see *Pinus resinosa*).

Pine, Scots (see Pinus sylvestris).

Pine, Shortleaf (see *Pinus echinata*). Pine, Sugar (see *Pinus lambertiana*).

Pine, White (see *Pinus strobus*). Pine Beetle, Mountain (see *Dendroc*-

tonus monticolae).

Pine Beetle, Southern (see Dendroctomus frontalis)

droctonus frontalis).

Pine Beetle, Western (see Dendroctonus brevicomis).

Pine Butterfly (see Neophasia mena-

pia).
Pine Moth (see Dendrolimus punctatus).

Pine Scale, Scotch (see Toumeyella numismatica).

Pine Shoot Moth, European (see Rhyacionia buoliana).

Pine Weevil (see Hylobius abietis).

Pineapple, pests of, in Fiji, 389, 390; pests of, in Hawaii, 49, 140, 273, 274, 298, 346, 390; Pseudococcus spp. on, in Mauritius, 514; P. brevipes on, in Nyasaland, 589; pests of, in W. Indies, 346, 373, 472, 498; host-range, vector and distribution of virus disease of, 139-141; P. brevipes causing wilt of, 346; key to mealybugs of, 514.

Pineapple Caterpillar (see Tmolus

echion).

Pineapple Mealybug (see Pseudo-coccus brevipes).

pineoides, Chermes (Pineus).

pinetum, Neodiprion. Pineus (see Chermes).

pinguis, Phytorus.

pini, Cinara; Coelonotus (Aphidius);
 Dendrolimus; Diprion; Exenterus; Ips; Microdus (Bassus);
 Pissodes; Platygaster; Trichogramma.

piniarius, Bupalus. pinicola, Cinara.

pinicolus, Ooencyrtus.

pinifoliae, Chermes (Pineus).

piniperda, Myelophilus.

Pinnaspis buxi (on coconut), Coccinellids introduced into Seychelles and Hawaii against, 324, 331.

Pinnaspis minor, on cotton in Dominican Republic, 498.

Pinnaspis minor strachani, factors affecting attacks of, on Citrus in St. Lucia, 262.

Pinthaeus sanguinipes, doubtful value of, against Leptinotarsa decemlineata in Germany, 70.

Pinus (see Pine).

Pinus banksiana, Toumeyella numismatica on, in Michigan, 605; Rhyacionia albicapitana on, in

Quebec, 37.

Pinus contorta, Rhyacionia buoliana on, in Br. Columbia, 415; Cole-optera on, in U.S.A., 49, 371, 426, 427, 648; little attacked by Polyphylla perversa, **509.**

Pinus contorta var. latifolia (murrayana), new weevil on, in Montana.

371.

Pinus densiflora, Eulachnus agilis on, in U.S.A., 607.

densiflora umbraculifera, Cinara pini on, in U.S.A., 607. Pinus echinata, Dendroctonus front-

alis on, in U.S.A., 49, 116. Pinus flexilis, Dendroctonus pondevosae on, in Colorado, 426, 427. Pinus halepensis, Eulachnus agilis

on, in U.S.A., 607.

Pinus jeffreyi, Coleoptera on, in California, 371, 551.

Pinus lambertiana, Dendroctonus monticolae on, in California, 49. Pinus leiophylla, Euproctis termin-

alis on, in S. Africa, 18.

Pinus massoniana, Dendrolimus punctatus on, in China, 545; Sipalus formosanus on, in Formosa, 104.

Pinus mugo, Rhyacionia buoliana on, in Br. Columbia, Eulachnus agilis on, in U.S.A., 607.

Pinus nigra, pests of, in Japan, 480; Eulachnus agilis on, in U.S.A., 607.

Pinus patula, Euproctis terminalis

on, in S. Africa, 18.

Pinus pinaster, Eulachnus agilis
on, in U.S.A., 607.

Pinus ponderosa, pests of, in U.S.A., 48, 49, 357, 371, 422, 426, 427, 551; effects on, of defoliation by Neophasia menapia, 422; fungi associated with Coleoptera in, injury to, by Anabrus 357; simplex, 48.

Pinus resinosa, 422; pests of, in U.S.A., 164, 400, 607, 637, 638. Pinus rigida, Dendroctonus frontalis

on, in U.S.A., 49.

Pinus strobus, 422; Pissodes strobi on, in Quebec, 37; pests of, in U.S.A., 164, 400, 607.

Pinus sylvestris, Rhyacionia spp. on, in Britain, 382, 383; pests of, in U.S.A., 422, 607; decay not rendering timber of, suitable for Xestobium rufovillosum, 315.

Pinus taeda, pests of, in Texas, 49. Pinus taiwanensis, Sipalus formosanus on, in Formosa, 104.

Pinus thunbergi, Dendrolimus punctatus on, in China, 545; Aphid on, in U.S.A., 607.

Piper betle, Lophobaris piperis on, in Netherlands Indies, 33.

Piper methysticum, Brachylybas variegatus on, in Fiji, 390.

Piper nigrum, Lophobaris spp. on, in Netherlands Indies, 32, 33, 328; susceptibility of varieties of, to L. piperis, 33; arsenicals causing injury to, 33.

Piperidine, toxicity of, to eggs of

Lygaeus kalmii, **352.**

piperis, Lophobaris; Spathius.

Pipunculus silvaticus, parasite of Laevicephalus striatus in Russia, 618.

piri, Anthonomus.

piricola, Toxoptera.

Piripiri (see Acaena).

pisi, Macrosiphum (Illinoia) (see M. onobrychis).

pisorum, Bruchus.

Pissodes approximatus, on Pinus resinosa in Connecticut, 164.

Pissodes notatus, effect of felling on, on conifers in Britain, 382.

Pissodes pini, effect of felling on, on conifers in Britain, 382.

Pissodes strobi (on pines), Quebec, 37; character of populations of, in U.S.A., 396.

pissodis, Coeloides.

Pistol Case-bearer (see Coleophora malivorella).

Pitch Acids, preparation and use in dust insecticides of, 247, 248; pitchates obtained from, 247.

Pitch Pine (see Pinus rigida).

Pithecolobium lobatum, pests of, in

Pithecolobium saman (lac host-tree), Aspidiotus destructor on, in Siam, 95.

Pityogenes (see Ips).

Gnorimoschema plaesiosema,

(Phthorimaea).

Plaesius javanus (predacious on Cosmopolites sordidus), establishment of, in Jamaica, 445; introduced into Rarotonga from Fiji, 496.

plagiata, Anaitis.

Plagiodera metallica, bionomics and control of, on Flacourtia indica in Philippines, 649.

Plagiodera versicolora, bionomics and control of, on willow in U.S.A., 438, 607.

Plane (see Platanus).

planicollis, Lyctus.

planifrons, Phanerotoma.

Plant Pest Legislation, against Diarthronomyia chrysanthemi in N. Ireland, 64; in U.S.A., 444, 498, 568; summaries of, in other countries, 349, 499; discussion of, 543.

Plantago major (Plantain), winter host of yellows disease of lettuce and endive in New York, 646.

Plants, survey of, as sources of insecticides, 483.

platanoides, Drepanosiphum.

Platanus, Xyleborus dispar on, in Br. Columbia, 328.

Platanus acerifolia, Teragra guttifera on, in S. Africa, 18

platensis, Aphidius.

Platybolium alvearium, gen. et sp. n., bionomics and control of, infesting beehives in India, 343.

platycephalus, Egius.

Platyedra gossypiella (Pink Cotton Bollworm), distribution of, in Argentina, 498; in Colombia, 53; in Ecuador, 539; in Egypt, 338; in Nyasaland, 153, 588; danger of introduction of, into Peru, 539; in Philippines, 648; in Punjab, 266; in Sudan, 386, 387; in Uganda, 228; in U.S.A., 262, 430; quarantine against introduction of, into U.S.A. from Porto Rico, 445; factors affecting susceptibility of varieties of cotton to, 266; other food-plants of, 386, 430, 445; bionomics of, 53; aerial distribution of, 262; resting stage in larvae of, 53, 338, 386, 430, 588; parasites and biological control of, 153, 386, 430, 588; treatment of cottonseed against, 53, 338.

Platygaster, parasite of Mayetiola destructor in Russia, 582.

Platygaster hiemalis, parasite of Mayetiola destructor in Russia, 618, 619; bionomics of, 619.

Platygaster minutula (see P. hie-

Platygaster pini, parasite of Contarinia juniperina in U.S.A., 433.

Platygaster zosine, parasite of Mayetiola destructor in Russia, 618. platyhypenae, Euplectrus.

Platynota flavedana, measures against, on greenhouse roses in New Jersey, 132.

Platyplectrus, gen. n., 514.

Platyplectrus natadae, sp. n., parasite of Natada nararia in Ceylon, 514.

Platyplectrus ornatus, sp. n., in South Africa, 514; parasite of Niphadolepis alianta in Tangan-yika, 514.

Platyplectrus orthocraspedae, sp. n., hosts of, in Java and Celebes, 514.

Platyptilia pusillidactyla, on Lantana in India, 289; destroyed by Heliothis, 289.

platyrhinus, Phytotribus.

Platystolus martinezi, in Spain, **529.** Platystolus surcularius, in Spain, **529.** plebeja, Anomala.

Plecoptera reflexa, natural enemies and biological control of, on Dalbergia sissoo in India, 74, 514.

plecopterae, Microgaster. Plesiocoris rugicollis, sprays against,

on apple in Britain, 281.

pleuralis, Eubadizon.

Pleurotropis, probably a hyperparasite of Antestia in Uganda, 230. Pleurotropis detrimentosus, hyperparasite of Artona catoxantha in Java and Malaya, 79.

Pleurotropis howardi, parasite of Apanteles liparidis in Japan, 172. Pleurotropis metallica, parasite of Mayetiola destructor in Russia,

582, 619.

Pleurotropis parvulus (parasite of Promecotheca papuana), in Fiji, 378; introduction of, into New Guinea Territory, 377, 378; bionomics of, 378.

Pleurotropis ptychomyiae, sp. n., hyperparasite of Artona catoxantha in Java and Malaya, 79.

Pleurotropis simiolus, parasite of Epilachna vigintioctomaculata in Korea, 103.

plicatalis, Macalla.

Plodia interpunctella, in stored cereals in Japan, 97, 102; in dried fruit in U.S.A., 426; effect of humidity on eggs of, 102; measures against, 97, 426; caught on treated fly-papers, 255.

plorabunda, Chrysopa. plorans, Euprepocnemis. Plotheia celtis (see Selepa).

Plum, Cydia pomonella on, in S. Africa, 415; Ceratitis capitata on, in Algerian Sahara, 616;

pests of, in Australia, 291, 335; pests of, in Britain, 108, 284, 286, 304; pests of, in Canada, 304, 307, 508, 548; pests of, in Germany, 231, 339; pests of, in India, 267, 518, 523; Drosophila suzukii on, in Japan, 622; Lecanium persicae on, in New Zealand, 58; Hoplocampa flava on, in Palestine, 622; Scolytus rugulosus on, in Peru, 539; pests of, in Russian Union, 581, 653; pests of, in U.S.A., 113, 192, 223, 415, 416, 536, 630, 635; forms of Rhagoletis pomonella developing on, 226; fumigation of fruit of, against Anarsia lineatella, 113.

Plum Curculio (see Conotrachelus

nenuphar).

Plum Leafhopper (see Macropsis

trimaculata).

Plum Nursery Mite (see Phyllocoptes fockeui).

Plum Sawfly (see Hoplocampa flava).

Plumbago, Engytatus geniculatus on, in Hawaii, 34.

Plume Moth, Grape (see Oxyptilus periscelidactylus).

Plumeria, termite attacking, in Barbados, 7.

Plusia brassicae (on crucifers), measures against, in S. Carolina, 199; in Hawaii, 651; laboratory rearing of, 557.

Plusia chalcites, on potato in Hawaii, 650; on Lantana in India, 289; on cacao in Malaya, 513; on weeds in Sumatra, 319; destroyed by Heliothis armigera, 289.

Plusia intermixta, bionomics of, on flax and vegetables in Formosa, 93.

Plusia moneta, measures against, on Delphinium in Britain, 448.

Plusia orichalcea, new parasite of, on soy beans in S. Africa, 514.

Plusia peponis, on Trichosanthes anguina in India, 264; Microbracon hebetor reared on, 264.

Plusia signata, bionomics of, on tobacco in Sumatra, 319.

Plutella cruciferarum (see P. maculi-

Plutella maculipennis (on crucifers), in S. Africa, 19; in Australia, 142, 546; in greenhouses in Britain, 144; in S. Carolina, 199; in Germany, 232; in Holland, 65; bionomics of, in Hong Kong,

153; in New Caledonia, 391; in New Zealand, 64, 333; in Russian Union, 585; parasites and biological control of, 19, 64, 65, 142, 333; fungus infesting, 19; measures against, 199, 585; tests of plant insecticide on, 242; laboratory rearing of, 557.

plutellae, Apanteles.

Pnyxia scabiei, relation of, to potato scab in New York, **644**.

Poa spp., Sitodiplosis cambriensis affecting seed production in, in Britain, 315; Crambus spp. damaging lawns of, in California, 408; pests of, in New Zealand, 334.

Pochazia fasciata, on cacao and Solanum in Nigeria, 516.

Podagrica, on tea and Bidens pilosa in Nyasaland, 588.

Podagrica breweri, on cotton in Queensland, 466.

Podagrica gemella, on cacao in Malaya, 513.

Podagrica puncticollis, on cotton, etc., in Sudan, 387.

Podagrica suahelorum, flea-beetle misidentified as, on tea in Nyasaland, 588.

Podapolipus, parasitic on locust in S. Africa, 322.

Podisma pedestris, fungi infesting, in Siberia, 485.

Podisus maculiventris, unlikely to be of value against Leptinotarsa decemlineata in Germany, 70; predacious on other insects in U.S.A., 121, 443; probably not injuring tomatos, 121.

Podisus serieventris, predacious on Gilpinia polytoma in Maine, 637.

Podomyia setosa, parasite of Plecoptera reflexa in India, 74; parasite of Prodenia litura in Java, 492.

Podothrips lucasseni, in Hawaii, 272, 273; on Valota insularis, 272; predacious on sugar-cane mite, 273.

Poecilips cardamomi, not greatly reducing yield of cardamom in India, 74.

poecilus, Dysdercus.

Pogonortalis doclea, Dacus confused with, in New South Wales, 237.

Poinciana regia, Xylopsocus castanoptera on, in Fiji, 170.

Poland, Sphinz pinastri in, 86; parasite of Diprion similis in, 294.

Polia oleracea, measures against, on greenhouse crucifers in Britain, 144, 145.

Polilla (see Calotermes brevis).

Polistes macaensis, predacious on Heliothis armigera in Fiji, 148.

Polistes major, predacious on Terastia meticulosalis in Porto Rico, 469.

politus, Chilocorus: Telenomus (Phanurus).

Polybia occidentalis, predacious on Thyridia themisto in Brazil, 292.

Polyblastus, parasite of Gilpinia polytoma in Maine, 637.

Polychrosis botrana, arsenicals against, on grape vines in Spain,

Polychrosis incultana, on wattle in S. Africa, 19.

Polychrosis viteana, measures against, on vines in U.S.A., 117, 425, 558.

Polygonum, not attacked by Brotolomia meticulosa, 525.

Polygraphus, on spruce in Canada,

Polygraphus gracilis, on spruce in Sakhalin, 96.

Polygraphus proximus, on spruce in Sakhalin, 96.

Polyodaspis endogena, 71.

Polyodaspis ruficornis, parasite of Pyrausta nubilalis and Hoplocampa in Europe, 71.

Polyphylla fullo, Scoliids parasitic on, in Russia, **572.**

Polyphylla perversa, bionomics of, in Br. Columbia, 508, 509.

Polyphylla variolosa, bionomics of, in forest nurseries in New York,

polyraphoides, Amniscus.

Polystictus abietinus, attacking spruce killed by bark-beetles in Sakhalin, 96.

polytoma, Gilpinia (Diprion).

pomaria, Typhlocyba.

Pomegranates, cold storage against Trypetids in Cyprus, 544. pometaria, Alsophila.

pomi, Aphis (Doralis).

pomonella, Cydia (Carpocapsa, Laspeyresia); Rhagoletis.

pomorum, Anthonomus; Pimpla. ponderosae, Dendroctonus; Ips. Pontia rapae (see Pieris).

Popillia japonica, precautions against introduction of, into Bermuda, 138; in U.S.A., 42, 118, 119, 123, 124, 125, 126, 161,

364, 368, 369, 370, 426, 431, 498, 499, 504, 594, 595; legislation against, in U.S.A., 499; feeding habits of adults of, 118; damaging turf, 124, 161; factors affecting winter mortality of, 594; natural enemies and diseases of, and their utilisation, 124, 125, 161, 368, 369, 370, 426, 431, 504; baits and traps for, 125, 126, 364, insecticides against 504, 595; adults of, 119, 595; other measures against, 42, 123, 426, 498, 499; survival of, in artificial media, 592; determination of instars of, 224.

popilliavora, Tiphia.

Poplar (Populus), 603; Teragra guttifera on, in S. Africa, 18; Membracid on, in Brazil, 392; pests of, in Canada, 347, 626; Lithocolletis populifoliella on, in Russia, 487; pests of, in U.S.A., 438, 632.

Popular Names (of insects), list of,

in U.S.A., 280.

Population Studies, of insects, 11, 152, 312, 396, 397, 450, 490, 592. populifoliella, Lithocolletis.

Populus (see Poplar).

Populus tremuloides, fungi associated with Trypodendron retusum on, in Minnesota, 165; Agrilus anxius on, in New Brunswick, 347.

Porina (see Oxycanus).

porosa, Hoplophora (see Hoplophorion pertusa).

porosus, Myzus.

Porotermes adamsoni, rearing of, in Australia, 142.

Porricondyla, not causing club gall on Cornus in U.S.A., 606.

Porthetria (see Lymantria).

Porthmologa paraclina, on Zizyphus jujuba in India, 264; Microbracon hebetor reared on, 264.

Porto Rican Mole-cricket (see Scap-

teriscus vicinus).

Porto Rico, Tingid on Citrus in, 51; forest and timber pests in, 51, 52, 427, 473; tests of timbers resistant to termites in, 157; Lathridiid in houses in, 361; Kolla fasciata damaging lawns in, 217; pests of leguminous plants in, 51, 52, 345, 469, 470, 473; pests and disease of maize in, 469, 470; miscellaneous pests in, 469 473; Pseudococcus brevipes on pineapple, etc., in, 346, 472; Agromyzid on sweet potato in, 261; beneficial insects and biological control in, 346, 356, 469, **470, 471, 472, 473, 552**; failure to establish Amazon strain of Metagonistylum minense in, 4: pests intercepted in quarantine in, 348; introduction of beneficial insects into other countries from, 4, 354, 431, 601; quarantines against introduction of pests into U.S.A. from, 444; pests from, intercepted in U.S.A., 349.

Portugal, Cryptolaemus montrouzieri introduced from Spain into, against Pseudococcus citri, 63.

Portulaca oleracea, Hymenia recurvalis recorded on, 46.

postfasciatus, Euscepes.

postica, Hypera (see H. variabilis). postvittana, Tortrix.

Potassium Ammonium Sulphide, a constituent of Selocide, 559.

Potassium Antimoniate, dust formula containing, against termites, 610.

Potassium Antimony Tartrate (see Tartar Emetic).

Potassium Caseinate, 559.

Potassium Chromate, ineffective as a preservative against Lyctus, 147.

Potassium Cyanide, in solution for treating nests of Solenopsis, 168; and sulphuric acid, for generating hydrocyanic acid gas (q.v.), 455.

Potassium Fluoaluminate, dusting with, against Pyralids on beans, 345.

Potassium Hydroxide, increasing injury to Citrus by sprays containing sulphur, 296.

Potato, Typophorus on, in Argentina, 512; pests of, in Bermuda, new Agromyzid on, in Brazil, 36; Aphids on, in Br. Isles, 28, 329, 375, 381, 590, 591; Leptinotarsa decemlineata on, in Jersey, 63; pests of, in Canada, 90, 302, 303, 330, 508; Pseudococcus citri on, in Egypt, 30; Epilachna vigintioctopunctata on, in Fiji, 390, 495; pests of, in France, 23, 24, 25; pests of, in Germany, 70, 449; pests of, in Hawaii, 34, 650, 651; pests of, in India, 265, 266, 516, 517, 523; Acanthocoris sordidus on, in Japan, 171; Gryllotalpa on, in Jugoslavia, 72; Epilachna vigintioctomaculata on, in Korea, 103; Gnorimoschema operculella on, in Nyasaland, 154; Empoasca on, in Peru, 511; relation of wireworms to, in Russia, 572, 573; pests of, in Victoria, 31; pests of, in U.S.A., 39, 153, 162, 163, 165, 188, 200, 201, 214, 221, 222, 247, 253, 278, 301, 302, 325, 440, 501, 503, 508, 537, 596, 638, 643, 644, 645; types of injury to, by Paratrioza cockerelli and other Rhynchota, 278, 302, 596, 638, 639; relation of Arthropods to Actinomyces scabies in, 501, 644; insects and virus diseases in, 23, 24, 27, 140, 165, 179, 188, 301, 302, 329, 339, 375, 500, 638; Paris green causing injury to, 326; (tubers), fumigation of, against Gnorimoschema operculella, 222; ment of, before planting against wireworms, 643; bait for Ceratitis capitata prepared 462.

Potato Aphis, Pink and Green (see Macrosiphum solanifolii)

Potato Beetle, Colorado (see Leptinotarsa decemlineata).

Potato Flea-beetle (see Epitrix cucumeris).

Potato Leafhopper (see Empoasca fabae).

Potato Psyllid (see Paratrioza cockerelli).

Scab Potato (see Actinomyces scabies).

Potato Tuber Moth (see Gnorimoschema operculella).

Potentilla spp., Aphids on, 27, 28; Nodonota puncticollis on roots of, in Pennsylvania, 192.

potentillae, Capitophorus (Pentatri-

chopus).

wilcoxii, fumigation Pothosagainst pests in U.S.A., 555.

Potosia aeruginosa (speciosissima), as host for breeding Scoliids in Russia, **572.**

Poverty Weed (see Iva axillaris).

praepotens, Apanteles.

praetiosa, Bryobia.

pratensis, Lygus; Trigonophymus. Predacious Insects, list of, in India, 175.

quindecimpunctatus, on Prepodes Torrubia fragans in Porto Rico, 473.

pretiosum, Trichogramma.

Prickly Lettuce (see scariola).

Prickly-pear (see Opuntia).

Prince Edward Island, potato Aphids in, 302; Incurvaria rubiella on raspberry in, 307.

Prionobrachium fuscum, on Abutilon and cotton in Argentina, 512.

Pristiphora erichsoni, on larch in Canada, 36, 532; parasites of, 37, 532.

Pristogonatopus conjunctus, parasite of Delphacodes striatellus in Siberia, 14.

Pristomeridia (Pristomerus) agilis, parasite of Cydia pomonella in U.S.A., 193.

Pristomerus chinensis, parasite of Cydia spp. in Far East, 33; characters of, 33.

Pristomerus chinensis f. erythrothoracis, status of, 34.

Pristomerus hawaiiensis, parasite of Lepidoptera in Hawaii, 650.

Pristomerus vulnerator, characters of, 33.

Pristomerus vulnerator f. erythrothoracis, a form of P. chinensis, 34.

Privet (Ligustrum), Tortrix unifasciana on, in Britain, 628; Amphimallus majalis on, in Normandy, 24; insects on, in U.S.A., 628, 629.

Probits, for expressing toxicity of insecticides, 433, 434.

proboscidaria, Fiorinia.

Prociphilus oriens, food-plants of, in Japan, 173.

Prodenia eridania (see Laphygma). Prodenia litura, on Colocasia in Fiji and New Guinea Territory, 170, 377; on banana in Guam, 331; on Lantana and jute in India, 289, 447; on tobacco in Netherlands Indies, 319, 492; on cotton in Philippines, 649; outbreak of, in Spain, 463; bionomics of, 319, 463; parasites and biological control of, 332, 492; other measures against, 171, 463, 492.

profana, Mictis.

profligator, Stylocryptus. proletella, Aleurodes.

prolongata, Xiphydria. Promachus, predacious on Nodonota puncticollis in Pennsylvania, 192.

Promecotheca papuana, bionomics and biological control of, in New Guinea Territory, 377–379.

promecothecae, Eurytoma.

Prontaspis citri (on Citrus), in Fiji, 389; in W. Indies, 471, 498; Hyperaspis billoti predacious on, 471.

pronubana, Tortrix.

Propylaea quatuordecimpunctata, predacious on cotton Aphids in Russia, **582**.

Prorops nasuta (parasite of Stephanoderes hampei), failure to breed, in Ceylon, 594.

Prosapegus atrellus, parasite of Sexava in New Guinea, 380.

proserpina, Megamelus.

Prosopis, Chlorochroa ligata associated with, in U.S.A., 500.

Prospaltella, parasite of Aulacaspis pentagona in Peru, 511.

Protanagyrus aciculatus, gen. et sp. n., parasite of Coccids in Argentina, **344.**

Protein Baits, for Rhagoletis pomonella, 563.

proteus, Urbanus (Goniurus).

Protoparce, lead arsenate against, on tobacco in U.S.A., 365.

Protoparce quinquemaculata, bionomics of, on tobacco in Ontario, 303.

Protoparce sexta, on tobacco and tomato in U.S.A., 218, 219, 641; parasite of 218; measures against, 641.

Protostrophus avidus, sp. n., on Citrus in Transvaal, 341.

Protostrophus salignae, sp. n., on Eucalyptus saligna in Zululand, 341.

Proutista moesta, on sugar-cane in Guam, 331.

provancheri, Sagaritis. proximus, Polygraphus.

proximus, 1 oiygraphus. pruininus, Bruchus.

Prune, Paratetranychus pilosus on, in Br. Columbia, 22; pests of, in U.S.A., 223, 418.

pruni, Hyalopterus (see H. arundinis); Illiberis.

pruniella, auct., Argyresthia (see A. ephippella).

prunifoliae, Rhopalosiphum. prunivorella, Parornix (Ornix).

Prunus spp., Lymantria dispar on, in China, 545; relations of pests to: in Ontario, 305, 306, 307; in U.S.A., 188, 190, 222. (See Cherry, Plum, etc.)

Prunus augustifolia, jumping buds due to weevil infesting, in Florida,

475.

Prunus emarginata, Rhagoletis spp. on, in California, **222**, **565**. pryeri, Margaronia (Glyphodes).

Psallus seriatus, on cotton in U.S.A., 198, 258, 599; wild food-plants

affecting incidence of, 198; tests of insecticides against, 258, 599.

Psara ambitalis, bionomics and control of, on tobacco in Netherlands Indies, 319, 492.

Psara bipunctalis, laboratory rearing of, 557; tests of plant insecticide on, 242.

Pseudaonidia articulata (see Selenaspidus).

Pseudaphycus angelicus, parasite of Pseudococcus adonidum in Cali-

fornia, 297.

Pseudaphycus utilis (parasite of Pseudococcus nipae), imported into Porto Rico from Hawaii, 472. pseudobrassicae, Rhopalosiphum. pseudobrevipes, Pseudococcus.

Pseudocentrinus lucidulus, sp. n., on Cocos spp. in Brazil, 468.

pseudococci, Anagyrus.

pseudococcina, Hambletonia.

Pseudococcus, new parasite of, on Acacia in 'Argentina, 344: measures against, in orchards in Br. Columbia, 23, 330; in Dominican Republic, 498; on coffee in New Guinea Territory, 377; on cotton in Peru, 511; in Seychelles, 82; key to species of, on pineapple, 514; utilisation of Coccinellids against, 82, 498.

Pseudococcus adonidum, in Australia, 280, 297, 298, 355; in Brazil, 298; in greenhouses in Britain, in California, 297, 298, 355; in Hawaii, 298; in Nyasaland, 62; in Porto Rico, 356; on Aleurites, 62; on Citrus, 280, 297, 298, 356; other food-plants of, 297, 298, 356; natural enemies and biological control of, 280, 297, 298, 355, 356, 489; other measures against, 280.

Pseudococcus brevipes (bromeliae, auct.), in Egypt, 30; in Fiji, 389, 390; in Hawaii, 346; in Mauritius, 514; in Nyasaland, 589; in W. Indies, 346, 472, 498; on pineapple, 346, 389, 390, 472, 498, 514, 589; causing wilt of pineapple, 346; on other plants, 30, 346; ants associated with, 346; natural enemies and biological control of, 346, 389, 472; other measures against, 30, 346.

Pseudococcus citri, in greenhouses in Canada, 310; in Dominican Republic, 498; measures against, in Egypt, 30; in Eritrea, 155; in Fiji, 389; in Java, 155; in

Palestine, 339; in Portugal, 63; in U.S.A., 256; on cacao, 498; on Citrus, 155, 389; other foodplants of, 30, 155, 256; factors affecting distribution of, 155; natural enemies and biological control of, 63, 310, 339, 498.

Pseudococcus comstocki, fungi and ants associated with, on Citrus in Brazil, 167, 168; in Japan, 425; bionomics of, on Citrus in Palestine, 73, 339; on apple and Catalpa in U.S.A., 48, 195, 425, 500; natural enemies and biological control of, 74, 168, 196, 339, 425, 500; other measures against, 73, 168.

Pseudococcus lilacinus (crotonis), on cacao in Malaya, 513.

Pseudococcus longispinus (see P. adonidum).

Pseudococcus maritimus, in California, 113, 223, 310; in greenhouses in Canada, 310; parasites of, 310; spray against, on pear, 223; fumigation of packed pears against, 113.

Pseudococcus nipae, in Hawaii, 472; food-plants and biological control of, in Porto Rico, 52, 472.

Pseudococcus perniciosus, on tea in Nyasaland, 154, 588; Eublemma costimacula predacious on, 588. Pseudococcus pseudobrevipes, sp. n.,

on pineapple in Mauritius, 514. Pseudococcus virgatus (see Ferrisiana).

Pseudoleucopis, species of, associated with Eriococcus coriaceus in Tasmania, 545.

Pseudoleucopis benefica (predacious on Eriococcus coriaceus), introduced into New Zealand from Australia, 545; bionomics of, 545.

pseudomagnoliarum, Coccus.

pseudosolani, Myzus (see Macrosiphum solani).

Pseudotsuga taxifolia, effect felling, on outbreaks of pests in Britain, 382; Polyphylla perversa on, in Br. Columbia, 509; pests of, in U.S.A., 49, 370, 427.

pseudotsugae, Dendroctonus.

pseudotsugata, Hemerocampa.

psidii, Dacus (Strumeta).

Psylla buxi, measures against, on Buxus sempervirens in U.S.A.,

Psylla mali, in Austria, 232; Britain, 284; in Russia, 585, 653; measures against, on apple, 232, 284, 585, 653.

Psylla pyricola, on pear in U.S.A., 48, 425, 533; bionomics and control of, 533.

Psylla sterculiae, bionomics of, on Brachychiton populneus in New South Wales, 467.

Psyllaephagus phytolymae, parasite of Phytolyma lata in Gold Coast, 492.

Psylledontus insidiosus, parasite of Psylla pyricola in Connecticut, 533.

Psyllid Yellows (caused by Paratrioza cockerelli), in tomato, 21; in potato, 278, 302, 596, 638; Pentatomid causing symptoms resembling, 596.

Psylliodes chrysocephala, on rape in Germany, 235.

Psyllobora luctuosa, apparently damaging cotton in Peru, 510.

Ptaeroxylon, defoliated by Achaea lienardi in S. Africa, 19.

Pteromalus, parasite of Contarinia medicaginis in Hungary, 12.

Pteromalus puparum (parasite of Pieris rapae), in Britain, 314; in Hawaii, 35; utilisation of, in New Zealand, 333; proposed introduction of, into Tasmania 468; imported into Victoria, 546.

Pteropus giganteus, carriage of Aspidiotus perniciosus by, in India, 272.

pterospermi, Pempherulus (Phylaitis).

Ptinids, keys to, in Canada and Britain, 152, 340; review of species of, of economic importance, 340.

Ptinus fur, in Colorado, 596; in grain sweepings in Russian Union, 576.

Ptinus tectus, in stored cacao, 616; daily rhythm of activity of, 527. ptychis, Hyalospila.

Ptychomyia (parasite of Artona catoxantha), parasites of, 79.

Ptychomyia remota, imported into India from Burma against teak pests, 75.

Ptychomyia selecta, parasite of Pristiphora erichsoni in Quebec, 37.

ptychomyiae, Pleurotropis. pubescens, Calotermes (Kalotermes).

puera, Hyblaea.

Pueraria hirsuta (thunbergiana), Anticarsia gemmatalis on, in U.S.A., 252, 599; in crop rotation against *Horistonotus uhleri*, 602.

pulchellus, Gampsocoris.

pulicaria, Chaetocnema.

pulverea, Holcocera.

Pulvinaria, Exochomus flavipes predacious on, in Seychelles, 83.

Pulvinaria aurantii, measures against, on Citrus in Transcaucasia, 653.

Pulvinaria convexa, new parasite of, in Argentina, 344.

Pulvinaria jacksoni, Coccinellid predacious on, on Aleurites in Nyasaland, 62.

pulvinariae, Coccophagus.

Pumice, as a carrier for cryolite dust, 145.

pumila, Fenusa (see F. pusilla).

pumilus, Metallus. Pumpkin, Phymateus leprosus

damaging, in S. Africa, 435.
punctata, Xanthopimpla.

punctatum, Anobium. punctatus, Dendrolimus.

puncticollis, Cylas; Nodonota; Podagrica.

punctiferalis, Margaronia (Gly-phodes).

punctigera, Urostylis.
punctillum, Stethorus (Scymnus).
punctiventris, Brachymeria; Cleo-

nus.
punctulata, Pheidole.

Punjab, new weevils on apple and Zizyphus jujuba in, 341; cotton pests in, 266, 516; miscellaneous pests in, 271; pests of sugarcane and cereals in, 81, 272, 520, 521; beneficial insects in, 82.

punjabensis, Empoasca. puparum, Pteromalus.

pupivora, Trichospilus. purchasi, Icerya.

purgatus, Enicospilus.

purpurea, Clausenia. purpureus, Tetrastichus.

purpurifascia, Papaipema.

Purslane (see Portulaca oleracea). pusilla, Fenusa.

pusillidactyla, Platyptilia.

Putnam's Scale (see Aspidiotus ancylus).

Pycnoderes quadrimaculatus, on beans in Hawaii, 650.

Pycnoscelus surinamensis, on pineapple in Fiji and Hawaii, **390.** pyloalis, Margaronia (Glyphodes). pyralis, Sacadodes.

Pyrausta nubilalis (European Corn Borer), in China, 446; in Danube countries, 71, 72; in Guam, 331; in Japan, 99; in Ontario, 303; in Russian Union, 478, 487, 488 in U.S.A., 43, 48, 162, 164, 253, 370, 471, 505, 508, 529; strains of maize in relation to, 253, 303, 505, 529; on potato, 162, 508; other food-plants of, 446, 488; bionomics and physiology of, 43, 99, 304, 487; insect parasites and biological control of, 71, 331, 471; Neoaplectana glaseri experimentally parasitic in, 370; measures against, 162, 505; effect of mechanical husker on, 253; larva of, 478.

Pyrethrins, extraction of, from green parts of pyrethrum plants, 586; content of, in pyrethrum dusts and sprays, 21, 46, 47, 52, 121, 134, 136, 248, 345, 361, 434, 441, 527, 557, 633, 640, 647.

Pyrethrum (including extracts), against ants, 588; against Aphids, 96, 145, 381, 585, 586; against Jassids, 19, 134, 248, 278, 440, 441, 647; against other Rhynchota, 19, 21, 22, 31, 52, 121, 171, 230, 278, 279, 281, 290, 361, 373, 465, 513, 557, 590, 596, 603; against Coleoptera, 70, 281, 361, 390, 434, 585, 595, 600, 617; injected into apple trees against Saperda candida, 243; against Diptera, 99, 527; against Lepidoptera, 18, 19, 46, 47, 129, 132, 154, 318, 345, 408, 411, 434, 585, 633, 640, 642; protecting stored potatoes against Gnorimoschema operculella, 138; against Lepismids, 56; against sawflies, 585; against thrips, 25, 31, 121, 136, 146, 413; increasing efficacy of oil for treating maize ears against pests, 217, 469, 470, 481; tests of toxicity of, to insects, 19, 99, 121, 233, 278, 281, 434, 483, 585, 586; toxicity to house-flies of smoke from, 127; duration of activity of, 233; dusting with, 18, 19, 21, 22, 25, 31, 46, 47, 52, 121, 134, 145, 146, 154, 171, 233, 248, 278, 290, 318, 345, 390, 441, 513, 557, 590, 596, 600, 633, 640, 647; carriers for, 21, 22, 25, 46, 134, 171, 278, 290, 318, 513, 557, 596; proprietary dusts of, 18, 21, 22, 46, 121, 134, 278, 557, 596, 633; preparation of extracts of,

586: and butyl carbitol thiocyanate, 56; and dichlorethyl ether, 136; compatible with dinitro-o-cyclohexylphenol, 249; and nicotine, 31; and oil emulsions, 411, 642; atomised sprays of oils containing, 19, 56, 230, 381, 434; method of dispersing extract of, in oil, 586; and rotenone insecticides, 233, 361, 527, 640, 642; Selocide compatible with, 560; and soap, 96, 132, 279, 290, 603, 617; other spreaders and wetters for, 136, 361, 595; and sulphur, 25, 46, 47, 52, 248, 441, 647; other insecticides compared with, 19, 21, 121, 233, 318, 414, 434, 435; study of dermatitis caused by, 527, 528.

Pyrethrum Plants, extraction of pyrethrins from green parts of, 586.

Pyrgota, parasite of Lamellicorns in Kentucky, 299.

pyri, Anisandrus (see Xyleborus dispar); Anthonomus (see A. piri).

pyricola, Psylla.

Pyridine, toxicity of, to eggs of Lygaeus kalmii, 352.

Pyrilla, natural enemies of, on sugar-cane in India, 82, 183, 184. Pyrilla perpusilla, bionomics and control of, on sugar-cane and cereals in Punjab, 81, 82.

pyrillae, Ageniaspis; Cheiloneurus; Dryinus (Lestodryinus); Tetra-

stichus.

Pyrilloxenos compactus, parasite of Pyrilla perpusilla in Punjab, 82. Pyrocide, Dry, 18, 21, 22, 46, 121, 134, 557, 633.

Pyrocide 20, 134.

Pyroderces rileyi, feeding habits of, on pineapple in Hawaii, 274.

Pyrophyllite, effect of electrostatic charges on dusts of, **569**.

pyrrhaspis, Pexopsis.

Pyrus sinensis, Cydia molesta on, in Japan and Korea, 372.

Pyrus spectabilis, pests of, in China, 446.

Q.

Quadraspidiotus (see Aspidiotus) quadridens, Ceuthorrhynchus. quadridentata, Ascogaster; Colobogaster. quadrifoveatus, Strategus. quadrimaculatus, Bruchus (see B. maculatus); Pycnoderes; Scym-

quadrinotata, Indarbela. quadripunctata, Scolia.

quadripunctatus, Oecanthus nigri-

quadripustulata, Winthemia.

quadrispinosus, Scolytus.

Quamoclit vulgaris, Cylas formicarius on, in Lousiana, 602.

Quarantine, Macrodactylus spinosus intercepted in, in Bermuda, 139; pests intercepted in, in Fiji, 148; Pachyrrhynchus moniliferus intercepted in, in Formosa, 103; pests intercepted in, in Porto Rico and Hawaii, 348; pests intercepted in, in U.S.A., 45, 114, 118, 133, 223, 348; data on insect pests obtained by interceptions in, 133. Quartz, dusts of, mixed with stored

wheat against Calandra, 541. Quassia, preparation of sprays of, against plum sawfly, 623; dusts containing, ineffective against Anthonomus pomorum, 70.

quaternus, Luperodes.

quatuordecimpunctata, Propylaea.

Quebec, forest pests in, 36-39, 431, 531-532, 626; methods for surveys of forest pests in, 311; orchard pests in, 224, 625; Cydia nigricana on peas in, 310: potato Aphids in, 302; mestes lardarius boring in timber in, 392; beneficial insects and biological control in, 36, 37, 38, 220, 311, 432, 532, 625, 626, 627.

Quebracho Tannin, fixed nicotines prepared with, 162, 256.

Queensland, banana pests in, 436, 465, 466; pests of Citrus in, 437, 465, 480; miscellaneous pests in, 149, 170, 465, 467; virus disease of pineapple in, 140; sugar-cane pests in, 388, 622; starch content of Eucalyptus in relation to Lyctus in, 150; Tortrix postvittana not a pest of apple in, natural enemies and biological control of insects in, 389, 466, 545; Arthropods and biological control of noxious plants in, 143, 289, 467, 587.

Queensland Fruit-fly (see Dacus ferrugineus tryoni). quenavadi, Pandesma.

Ouercus (see Oak).

Quercus ilex var. suber (Cork Oak), Lamellicorn larvae damaging, in Morocco, 83.

Quercus palustris, Euproctis terminalis reared on, in S. Africa, 18.

Quince, pests of, in Australia, 141, 298; Tortricids on, in Baluchistan, 267, 518; Typhlocyba froggatti on, in Britain, 284; Cydia molesta on, in Japan and Korea, 372; pests of, in U.S.A., 190, 242.

quindecimpunctatus, Prepodes.

Quinoline, toxicity of, to eggs of Lygaeus kalmii, 352. quinquecincta, Myzine. quinquemaculata, Protoparce (Phle-

R.

Racial Segregation, practical importance of, in insects, 548.

Radish, Galeruca tanaceti on, in Germany, 68; Murgantia histrionica on, in Hawaii, 651; Aphids on, in India and U.S.A., 266, 596.

Ragi (see Eleusine coracana). Ragwort (see Senecio jacobaea). Ragwort-seed Fly (see Hylemyia

jacobaeae).

gethontius).

Rainfall, relation of locusts and grasshoppers to, 48, 141, 388, 484, 485, 486, 520, 523, 570, 589 relation of other insects to, 1, 19, 38, 39, 41, 42, 83, 87, 94, 98, 105, 148, 151, 161, 181, 182, 270, 313, 338, 368, 404, 431, 433, 442, 531, 533, 547, 627.

Raisin Moth (see Ephestia figuli-

lella).

Raisins, in baits for Otiorrhynchus ligustici, 252, 357; used for feeding insects, 310, 395, 619; effect on Oryzaephilus surinamensis of feeding on, 187. (See Fruit, Dried.)

ramburi, Chrysopa. rameus, Xyleborus. rapae, Pieris (Pontia). rapax, Aspidiotus.

Rape, Myzus persicae wintering on. in Eire, 329; pests of, in Germany, 231, 235; Meligethes aeneus on, in Russian Union, 585.

Rape Blossom Weevil (see Meligethes aeneus).

rapidus, Adelphocoris. rapo, Tetrastichus.

Rarotonga, Plaesius javanus imported from Fiji into, against Cosmopolites sordidus, 496.

Raspberry, pests of, in Britain, 281, 283, 287, 544; pests of, in Canada, 307, 508; pests of, in U.S.A., 287, 643; susceptibility of varieties of, to Incurvaria rubiella, 307; insects and virus diseases of 287, 288 diseases of, 287, 288.

Raspberry Beetle (see **Byturus**

tomentosus).

Rayon (Artificial Silk), silverfish damaging, 56, 401.

recta, Thosea.

rectangula, Autographa. recurvalis, Hymenia.

Redbud (see Cercis canadensis).

Reeds (see Phragmites).
reflexa, Halyomorpha; Plecoptera. Reinecke's Salt, insoluble compound of nicotine and, 207.

relictum, Anomalon (Paranomalon). religiosus, Xylothrips.

remota, Ptychomyia. remulus, Trichacis.

repandus, Calotermes (Kalotermes). Resin, uses of, in sprays, 102, 419, 420, 437, 517.

Resin Residue, emulsions of, in sprays, 419, 420, 606; efficiency of dispersing agents for, 112, 113.

Resin Soap (see Soap, Resin). resinella, Rhyacionia (Evetria).

Reticulitermes flavipes, ants predacious on, in Massachusetts, 499. tibialis, damaging Reticulitermes trees, timber, etc., in U.S.A., 400, 537.

Reticulitermes tumiceps, in Utah, 400.

retusum, Trypodendron. reuteri, Drepanothrips. reversa, Sarcophaga.

Reviews:—Chamberlin (W. J.), Entomological Nomenclature and Literature, 526; Clausen (C. P.), Entomophagous Insects, 431; Comstock (J. H.), An Introduction to Entomology—9th edn., 275; Cotton (R. T.), Insect Pests of Stored Grain and Grain Products . . ., 605; Dodd (A. P.), The Biological Campaign Prickly-pear, against Fraenkel (G. S.) & Gunn (D. L.), The Orientation of Animals . 178; Hamilton (C. C.), Ed., Entoma. A Directory of Insect Pest Control, 438; Holmes (F. O.), Handbook of Phyto-

pathogenic Viruses, 437; Knight (P.), The Problems of Insect Study, 189; Large (E. C.), The Advance of the Fungi, 111; Leech (J. G.), Insect Transmission of Plant Diseases, 239; Tothill (J. D.), Ed., Agriculture in Uganda, 175.

Rhabdocnemis obscura (Sugar-cane Beetle Borer), in Hawaii, 50, 274; in Queensland, 389; study of adult movements of, 50; natural

enemies of, 274, 389.

Rhabdophaga heterobia, population fluctuations and natural enemies of, on Salix triandra in Britain, 312, 313.

Rhabdopterus picipes, measures against, on apple in New York,

642, 643.

Rhaconotus, parasite of Scirpophaga in India, 183.

Rhaconotus caulicola, sp. n., parasite of Scirpophaga and Chilo in India, 137.

Rhaconotus cleanthes, ecology of, parasitising Pempherulus affinis in Madras, 522.

Rhaconotus menippus, hosts and ecology of, in Madras, 522.

Rhadiurgus (see Hemiteles).

Rhagoletis, tests of sprays against, on cherry in U.S.A., 196, 641; nicotine in honey repellent to, 206.

Rhagoletis cerasi, on cherry in Jugoslavia, 72.

Rhagoletis cingulata, on cherry in U.S.A., 196, 222; quarantine against, in California, 222; possible race of, confined to Prunus emarginata in California, 222, 565; parasite of, 565.

Rhagoletis fausta, in New York, 196; quarantine against, in California, 565; possible race of, on Prunus emarginata in Cali-

fornia, 565.

Rhagoletis pomonella (on apple), in U.S.A., 161, 364, 563; fumigation of harvested apples against, 364; other measures against, (blueberry race), 161, 163; measures against, in Maine, 165; cross-breeding experiments with forms of, on various fruits in Nova Scotia, 226.

Rhagoletis suavis completa, insecticides against, on walnuts in California, 561.

rhamni, Aphis.

Rhamnus spp., winter food-plants of Aphis abbreviata in Maine, 188.

Rhinaria perdix, on strawberry in Australia, 494.

rhinoceros, Oryctes.

Rhinoscapha lagopyga, measures against, on Citrus in Fiji, 389.

Rhinotermes intermedius, laboratory rearing of, in Australia, 142.

Rhizoecus amorphophalli, sp. n., on Amorphophallus variabilis

Java, 88.

Rhizopertha dominica (in stored cereals), measures against, in U.S.A. and Australia, 209, 380; factors affecting food-finding and oviposition by, 451, 452; review of data on, 480.

Rhizotrogus spp., search for parasites of, in N. Africa, 83; in

Normandy, 24.

Rhodanates (see Thiocyanates).

Rhodes Grass (see Chloris gayana). Rhodesia, Northern, locusts and

Scelio howardi in, 491.

Rhodesia, Southern, Calandra oryzae on field and stored maize in, 184; Gonipterus scutellatus in, 30; Myzus persicae and rosette disease of tobacco in, 84, 184; Tachinids of, 655; other beneficial insects in, 30, 184, 514.

Rhodnius, action of pyrethrum on,

483.

rhododendri, Giardomyia.

Rhododendron, pests of, in U.S.A., **86, 606.** (See Azalea.)

rhodophila, Diacrisia.

rhodophilodes, Diacrisia rhodophila. Rhoecocoris sulciventris, sprays against, on Citrus in Queensland,

Rhogas, in Colombia, 53; nomics of, in Ceylon, 174; in Nyasaland, 154, 588; in Peru, 510; parasite of Noctuids on cotton, **53**, **510**, **588**; parasite of Limacodids on tea, **154**, **174**; parasitised by *Mesochorus orna*tus, 588.

Rhogas dendrolimi (metanastriae), parasite of Dendrolimus in Sakhalin, 95; synonymy of, 95. rhopaloceros, Xanthopimpla.

Rhopalosiphum nymphaeae, experiments with virus diseases of Musa textilis and, in Philippines, 317.

Rhopalosiphum prunifoliae, on apple in U.S.A., 164, 194, 195, 248;

measures against, 194, 248; technique for testing ovicides against. 195.

Rhopalosiphum pseudobrassicae, on crucifers and solanaceous plants in India and Hawaii, 266, 651; bionomics and control of, in U.S.A., 596.

Rhopobota naevana, measures against, on cranberry in Washington, 640, 642.

Rhoptrocerus eccoptogastri, parasite of Dendroctonus frontalis in Virginia, 211.

Rhoptromeris (see Eucoila).

Rhubarb, Galeruca tanaceti on, in Germany, 68.

Rhus, Achaea lienardi on, in S. Africa, 19; Acrocercops on, in Japan, 99.

Rhyacionia, new parasite of, on pines in U.S.A., 87.

Rhyacionia albicapitana, on Pinus banksiana in Quebec, 37.

Rhyacionia buoliana (European Pine Shoot Moth), in Britain, 383; parasites of, in Br. Columbia, 415; in U.S.A., 396, 563; character of populations of, 396; measures against, 415, 563.

Rhyacionia comstockiana, new para-

site of, in U.S.A., 86.

Rhyacionia frustrana, in U.S.A., 87, 396; character of populations of, on pine, 396; new parasite of, 87.

Rhyacionia resinella, on Pinus sylvestris in Britain, 383.

Rhyacionia turionana, bionomics and control of, on Pinus sylvestris in Britain, 383.

Rhyacionia washiyai, sp. n., on Pinus nigra in Japan, 480.

Rhynchites bacchus, meteorological conditions affecting, on fruit trees in Russia, 580; Monilia cinerea spread by, 581.

Rhynchites haradai, sp. n., on Picea jezoensis in Japan, 40.

Rhynchophorus, on coconut in New Caledonia, 391.

Rhynchophorus ferrugineus, natural enemies of, on coconut in India, 270.

Rhynchophorus palmarum, bionomics and control of, on coconut, etc., in Colombia, 8, 9.

Rhynchophorus papuanus, on coconut in New Britain, 377.

Rhynchosia memnonia, Empoasca lybica on, in Sudan, 387.

787

Rhyparida morosa, susceptibility of varieties of banana to, in Queensland, 465.

ribesii, Nematus.

ribis, Eriophyes.

Ricania speculum, on cotton in

Philippines, 649.

Rice, pests of, in Ceylon, 594; Spodoptera mauritia on, in Fiji, 495; Chondracris rosea attacking, in Formosa, 94; pests of, in Guam, 331; pests of, in Japan, 97, 98, 100, 101, 272; pests of, in Peru, 511; Pyralid on, in Spain, 458; Blissus insularis on, in Trinidad, 373; locusts damaging, in Uganda, 229; experiments with Nephotettix and virus disease of, 272; Verania discolor feeding on pollen of, 95; Lema cyanella sapporensis not feeding on, 93; insecticides and injury to, 599.

Rice (Stored), pests in, 92, 102, 172, 453, 455, 493, 619; fumigation of, against insects, 429; in baits for *Gryllotalpa* spp., 237.

Rice, Wild, Septis on, in Maine, 560. Rice Borer (see Chilo simplex).

ricini, Anychus.

Ricinus communis (see Castor). rileyi, Eulachnus; Pyroderces. rimosalis, Evergestis.

Ripersia internodia, measures against, on maize roots in Egypt, **30.**

ritchiei, Achrysocharella; Chelonus (Chelonella).

Robinia, Anobium punctatum in timber of, 233.

Robinia pseudacacia, Chalepus dorsalis on, in U.S.A., 354.

roborator, Pimpla (Exeristes). robusta, Eurytoma; Hypsipyla.

robustus, Gordius.

Rock Phosphate, dusts of, mixed with stored wheat against Calandra, 541.

Rodolia breviuscula, predacious on Icerya aegyptiaca in India, 263.

Rodolia cardinalis (predacious on Icerya purchasi), introduced into S. Africa, 320; in S. Australia, 280; utilisation of, in California, 280; establishment of, in Cyprus, 544; introduced into Dominican Republic, 498; utilisation of, in Peru, 511; failure to establish, against I. seychellarum in Seychelles, 82.

Rodolia chermesina, bionomics and utilisation of, in Seychelles, 82.

Rodolia obscura, predacious on Icerya purchasi in Nyasaland, 62. Rondania cucullata, bionomics of, in Italy, 316.

Rooks, destroying Melolonthids, 24. Rosa (see Rose).

rosae, Typhlocyba.

rosana, Tortrix (Cacoecia).

Rose (Rosa), Macrodactylus subspinosus intercepted on, in Bermuda, 139; Thrips fuscipennis on, in greenhouses in Britain, 145; pests of, in Canada, 508, 532; pests of, in Chile, 512; Myzus circumflexus on, in Hawaii, 651; Myzus porosus on, in Mexico, 46; Niphadolepis alianta on, in Nyasaland, 59; Typhlocyba froggatti on, in Tasmania, 291; pests of, in U.S.A., 46, 132, 188, 207, 256, 532, 595; relation of Capitophorus spp. to, 27, 28; virus diseases of, 255, 532; low absorption of selenium from nutrient solutions by, 207.

Rose Leaf Beetle (see Nodonota

puncticollis).

Rose Thrips (see *Thrips fuscipennis*). Rose Weevil, Fuller's (see *Pantomorus godmani*).

rosea, Chondracris. roseus, Anuraphis.

roseveari, Aprostocetus. rosimon, Castalius.

rossica, Locusta migratoria.

rossica, Locusta migratoria rostrata, Peronia.

Rotenoids, definition and determination of, 135.

Rotenone, against Aphids, 135, 250, 360, 504, 544, 640; against Coccids, 349, 350, 426; against Coleoptera, 120, 247, 595, 630, 640; injected into apple trees Saperda, 243, 244; against against Hoplocampa, 109; against Lepidoptera, 10, 46, 47, 135, 345, 640, 642; not affecting midgut wall of Laphygma eridania, 481; against mites, 237, 249, 639, 641; against Rhagoletis spp., 161, 562; of little value against Pentatomids, 603; ineffective against Thrips tabaci, 121, 122; in dusts, 46, 47, 121, 135, 247, 249, 345, 544, 562, 630, 640; pH of carriers affecting toxicity of, 301; oil-impregnated dusts containing, 161; in sprays, 10, 109, 120, 121, 122, 345, 349, 350, 360, 426, 595, 603, 640, 642; activator and wetting agents for, 360, 595;

formulae containing, 109, 120, 237, 243, 244, 360; compatible with dinitro-o-cyclohexylphenol, 249; and nicotine, 250; and oil emulsions, 109, 121, 122, 237, 349, 350, 426, 642; atomised sprays of oils containing, 250, 504; and pyrethrum, 640, 642; solvents for, 237, 243, 244, 349, 350. (See Cubé, Derris, Tephrosia, and Timbo.)

rouxi, Habrolepis.
rubecula, Apanteles.
rubens, Ceroplastes.
rubi, Amphorophora.
rubicola Aleundes

rubicola, Aleurodes (see Asterobemisia carpini).

rubiella, Incurvaria (Lampronia). rubiellum, Macrosiphum.

rubifolium, Macrosiphum. rubigalis, Phlyctaenia.

rubripes, Scotussa.

Rubus, pests of, in Britain, 287, 544; virus diseases of, 287, 288. (See Blackberry, Raspberry, etc.)

rufescens, Dolichocolon. ruficauda, Parabrachycoma.

ruficeps, Scolia. ruficollis, Dysdercus.

ruficornis, Polyodaspis; Scolia. rufinodus, Zethus.

rufipes, Hylastes (Hylurgopinus);

Phytomyza. rufiscutellaris, Glypta. rufiventris, Dissolcus. rufovillosum, Xestobium.

rufus, Triaspis curculionis.

rugiceps, Euetheola.

rugicollis, Minthea; Plesiocoris. rugosa, Lachnosterna (Phyllophaga). rugulosus, Scolytus.

Rumania, forest pests in, 12, 474; beneficial insects, 13, 71.

rumicis, Aphis; Hypera.

Russia (U.S.S.R.), beet pests in, 455, 456, 569, 577, 583, 584, 586, 654; cereal pests in, 487, 569, 571, 572, 573, 574, 576, 582, 583, 618, 619; Laevicephalus striatus and mosaic of cereals in, 15, 617, 618; pests of Citrus in, 582, 652, 653, 654; cotton pests in, 235, 582, 584; pests of crucifers in, 110, 478, 585, 654; flea-beetles on flax in, 181; pests of leguminous crops in, 572, 573, 575, 576; miscellaneous pests in, 159, 487, 575; orchard pests in, 528, 569, 580, 581, 582, 585, 652, 653; weevil on Salvia sclarea in, 15; pests of sweet potato and

Convolvulus in, 479; insects on Taraxacum kok-saghyz in, 484; Phylloxera on vines in, 578-580 pests of stored grain in, 576; Braconids of, 480; Coccid fauna of, 480; studies on Elaterids, Melolonthids and Tenebrionids identification of in, 571–573; injurious Lepidoptera in, 478; locusts and grasshoppers in, 487, **569–571, 583**; beneficial insects and biological control in, 235, 456, 479, 569, 572, 573, 575, 581, 582, 618, 619; utilisation of fungi and bacteria against insects in, 582-584. (See Siberia and Asia, Central.)

Russian Far East, Aphthona euphor-

biae on flax in, 181. rusti, Coccophagus. rustica, Dexia.

ryderi, Eriophyes.
Rye, Tipulids on, in Germany, 68;
pests and virus diseases of, in
Russian Union, 13, 14, 574, 617,
618; pests of, in U.S.A., 254,
300, 603.

Rye (Stored), pests in, and their control, 73, 576.

Rye-grass (see Lolium).

S.

Sabal bermudiana, Coccotrypes dactyliperda in fallen fruits of, in Bermuda, 139.

sabinae, Cinara (Lachnus).

sabulifera, Anomis (Cosmophila).

sabulosum, Opatrum.

Sacadodes pyralis, bionomics and biological control of, on cotton in Colombia, 8, 53; other measures against, 53.

saccharalis, Diatraea. sacchari, Trionymus.

saccharina, Lepisma; Tomaspis. Saccharine, tests of, in sprays and

baits, 296, 612.

Safflower (see Carthamus tinctorius). Sagaritis, parasite of Gilpinia polytoma in Maine, 637.

Sagaritis provancheri, parasite of Heliothis armigera in U.S.A., 598.

Sage, Clary (see Salvia sclarea). sagittiferella, Citripestis.

Sago Palm (see Metroxylon sagu). Sahlbergella, on cacao in Gold Coast, 492, 516; spray against, 492.

Sahlbergella singularis, on cacao in W. Africa, 492, 515.

Sahlbergella theobroma, on cacao in W. Africa, 492, 515.

St. Croix, Heliothis virescens on cotton in, 11.

St. John's Wort (see Hypericum).

St. Lucia, miscellaneous pests and biological control in, 4, 262, 263; Metagonistylum minense introduced into St. Vincent from, 263.

St. Vincent, Metagonistylum minintroduced into, ense against Diatraea saccharalis, 263.

saintebeauvei, Coccophagus.

Saipan, Hispid on coconut in, 331. Saissetia coffeae (hemisphaerica), on Citrus, coffee, etc., in Dominican Republic, 498; on cotton in Philippines, 649.

Saissetia nigra, history of, in California, 114; popular names and erroneous records of, 564.

Saissetia oleae, in E. and S. Africa, 385, 386; in Argentina, 344; in California, 209, 211, 349, 350, 385, 386, 561, 564; in Dominican Republic, 498; in Peru, 511, 539; on Citrus, 209, 220, 349, 386, 498, 561, 564; on olive, 511, 539; natural enemies and biological control of, 211, 219, 220, 344, 385, 386, 511, 539; importance of winter mortality of, 209; tests of insecticides against, 209, 349, 350, 561; S. nigra confused with, 564.

saissetiae, Microterys.

Sakhalin, Agrotis ypsilon on beet in, 100; Criocerid on cereals in, 93; pests of conifers in, 95, 96, 480; parasites of Dendrolimus superans in, 95.

salaminia, Eumaenas (Othreis).

salicis, Stilpnotia.

salignae, Protostrophus.

Salix (see Willow).

salomonis, Euplectromorpha.

Salt (Sodium Chloride), in baits, 269, 406, 461, 612, 613; ineffective against Calandra oryzae in stored wheat, 541; increasing ovicidal action of nicotine, 556.

saltator, Conocephalus.

Salvia sclarea, Phrydiuchus topia-rius on, in Crimea, 15.

Samanea saman (see Pithecolobium). Samoa, Bostrychid on Artocarpus and Hibiscus in, 170; introduction of insects into, against Oryctes and Lantana, 148, 496.

San José Scale (see Aspidiotus

perniciosus).

sanborni, Macrosiphum (Macrosiphoniella).

Sand Pear (see Pyrus sinensis). Sand Wireworm (see Horistonotus

uhleri).

Sandal, studies of insects and spike disease of, in India, 75, 382.

sanguinea, Cycloneda; Tomaspis (Cercopis)

sanguineus, Helopeltis.

sanguinipes, Pinthaeus.

sanguinolenta, Aceratagallia. sanninoideae, Microbracon.

Santalum album (see Sandal).

Saperda candida, bionomics and control of, in New York, 242-244.

Saperda tridentata (see Eutetrapha). Saponin, as a spreader and wetting agent, 145, 556.

sapporensis, Lema cyanella.

Sapromyza fuscicornis, Dacus confused with, in New South Wales, 237.

Sarcocephalus cordatus, Margaronia punctiferalis on, in Burma, 78. Sarcophaga, parasite of Alabama

argillacea in Colombia, 53. Sarcophaga aculeata, parasite grasshoppers in Canada, 308. 309; introduced into Argentina, 308.

Sarcophaga atlanis, parasite of grasshoppers in Canada, 308, 309; introduced into Argentina, 308.

Sarcophaga caridei, parasite of grasshoppers in Argentina, 308; troduction and rearing of, in Canada, 308.

Sarcophaga coloradensis, parasite of grasshoppers in Canada, 308.

Sarcophaga exuberans, parasite of Polyphylla perversa in Br. Columbia, 509.

Sarcophaga fuscicauda, parasite of Rhynchophorus ferrugineus India, 270.

Sarcophaga hunteri, parasite grasshoppers in Canada, 308, 309; introduced into Argentina, 308.

Sarcophaga kellyi, parasite of grasshoppers in Canada, 308.

Sarcophaga lineata, parasite of grasshoppers in Siberia, 486.

Sarcophaga opifera, parasite of grasshoppers in Canada, 308; introduced into Argentina, 308.

Sarcophaga reversa, parasite grasshoppers in Canada,

sarcostega, Tortrix (Cacoecia).

Sardine Oil, 255.

Saskatchewan, new Curculionid on *Iva axillaris* in, **530**; grasshoppers and their natural enemies in, **280**, **308**, **309**, **393**, **394**.

satyriniformis, Melittia.

saucia, Coelophora.

saundersi, Crossotarsus.

sauteri, Dinarmus.

Savanna Grass (see Axonopus compressus).

Sawdust, in baits for locusts and grasshoppers, 20, 144, 461, 612; as bait for termites, 387.

Saw-toothed Grain Beetle (see Ory-zaephilus surinamensis).

Sawfly, Apple (see Hoplocampa testudinea).

Sawfly, European Spruce (see Gilpinia polytoma).

Sawfly, Gooseberry (see Nematus ribesii).

Sawfly, Larch (see Pristiphora erichsoni).

Sawfly, Plum (see Hoplocampa flava).

Sawfly, Western Hemlock (see Neo-diprion tsugae).

Sawfly, Western Wheat-stem (see Cephus cinctus).

sayi, Chlorochroa.

scabiei, Pnyxia.

scalaris, Megaselia; Stauroderus; Taeniotes.

Scale, Bamboo (see Asterolecanium bambusae and A. miliaris).

Scale, Black, application of name, 564.

Scale, Brown Elm (see Lecanium corni).

Scale, California (see Aspidiotus perniciosus).

Scale, Citrus or California Red (see Aonidiella aurantii).

Scale, Coconut (see Aspidiotus destructor).

Scale, Florida Red (see Chrysom-phalus ficus).

Scale, Olive (see *Parlatoria oleae*). Scale, Oyster-shell (see *Lepido*-

saphes ulmi).
Scale, Pink Wax (see Ceroplastes rubens).

Scale, Purple (see Lepidosaphes beckii).

Scale, Putnam's (see Aspidiotus

ancylus). Scale, San José (see Aspidiotus

perniciosus).
Scale, Scotch Pine (see Toumeyella numismatica).

Scale Insects (Coccids), catalogue of, of Argentina, 343; species of, described by de Charmoy from 424: of Russian Mauritius, Union, 480; natural enemies of, 17, 34, 52, 62, 63, 74, 82, 94, 102, 168, 172, 196, 210, 211, 219, 220, 263, 264, 265, 280, 288, 290, 297, 298, 310, 320, 324, 331, 339, 344, 346, 355, 356, 363, 385, 386, 389, 425, 445, 471, 472, 489, 498, 500, 511, 539, 544, 545, 546, 582, 588, 594, 610, 625; ants associated with, 82, 168, 332, 346, 475; classification and new species of, 88, 514, 622.

scammelli, Myzus.

Scapteriscus vicinus, introduction of Larra americana into Porto Rico against, 552; treatment of lawns against, in Trinidad, 374. Scelio, parasite of locust eggs in

Formosa, 94.

Scelio chortoicetes, egg-parasite of Austroicetes cruciata in W. Australia, 169.

Scelio howardi, egg-parasite of Nomadacris septemfasciata in N. Rhodesia, 491.

Scelio pembertoni, egg-parasite of Oxya chinensis in Hawaii, 49. Sceloporus, destroying Toxoptera

graminum in Oklahoma, 251. Schinus molle, Bruchus elegans on, in Chile, 512; Bostrychid boring

in, in New South Wales, 335. Schist Oils, emulsions of, unsatisfactory against *Psylla mali*, 653.

schistaceana, Eucosma.

Schistocerca, of Argentina, 528. Schistocerca gregaria, in Afghanistan, 520; in India, 269, 271, 323, 341–343, 519, 520, 523; in Sudan, 388; effects of humidity on, 341–343; food preferences of, 519; causes and course of outbreaks of, 519, 520, 523; migrations of solitary phase of, 323; measures against, 269, 519.

Schistocerca paranensis, in Argentina, 308, 322, 538; in Br. Honduras, 498; in Peru, 539; phases of, 322; periodicity of, in relation to sun spots, 538; crops attacked by, 539; parasite from Canada not developing in, 308; measures against, 498, 538, 539.

Schizoneura lanigera (see Eriosoma). Schizonotus sieboldi, parasite of Plagiodera versicolora in U.S.A., 439.

Schizonycha, parasites of, in Zanzibar, 83.

Schleicheria trijuga (lac host-tree), pests of, in India, 263, 264.

schmeltzi, Papilio.

Schotia, Achaea lienardi on, in S. Africa, 19.

schultzei, Allodontermes; Frankliniella.

Sciara, relation of, to potato scab in New York, 644.

Sciara pauciseta, associated with pit scab in potatoes in U.S.A., 501.

Scirpophaga, parasites of, on sugarcane in India, 137, 183.

Scirtothrips aurantii, on Citrus in S. Africa, 18, 320; new Anthocorid predacious on, 320; measures against 18

sures against, 18.
Scirtothrips citri, on Citrus in California, 295–297, 407, 561, 565; on ornamental plants, 565; in-

secticides against, 295-297, 407, 561.

Scirtothrips signipennis, measures against, on banana in Queensland, 436, 437, 465; susceptibility of varieties of banana to, 465.

scitaria, Striglina.

Sclerotinia fructicola, 418.

Scobicia declivis, fumigation of filled wine casks against, in California, 222.

Scolia dejeani, observations on, in

Russia, 572.

Scolia flavifrons, observations on, in Russia, 572, 573.

Scolia hirta, observations on, in Russia, 572.

Scolia oryctophaga, in Madagascar and Mauritius, 148; experimentally breeding in Oryctes rhinoceros in Java, 148; methods of transporting, to Samoa, 148.

Scolia quadripunctata, observations

on, in Russia, 572.

Scolia ruficeps, doubtful relation of, to Oryctes rhinoceros in Malaya, 148.

Scolia ruficornis, in Zanzibar, 148; possible value of, against Oryctes rhinoceros in Samoa, 149.

Scolothrips sexmaculatus, predacious on mites in California, 210.

Scolytids, new species of, 528, 592. (See Bark-beetles.)

Scolytoplatypus daimio, control of, possibly transmitting fungous disease of chestnut in Japan, 97.

Scolytus destructor (see S. scolytus). Scolytus manglissiensis, sp. n., on fruit tree in Transcaucasia, 528.

Scolytus multistriatus (on elm in U.S.A.), relation of, to Ophiostoma ulmi, 116, 212, 364, 502, 563, 606, 644, 645, 646; isolation of O. ulmi from, when stored at different temperatures, 166; other fungi associated with, 116; bionomics of, 163; probably spread by wind, 606; measures against, 163, 212, 427, 644.

Scolytus quadrispinosus (Hickory Bark-beetle), bionomics and control of, in U.S.A., 116.

Scolytus rugulosus, measures against, on fruit trees in Peru, 539.

Scolytus scolytus, absence of digestive enzymes in larvae of, 340.

Scolytus sulcatus, experimentally transmitting Ophiostoma ulmi on elm and apple in U.S.A., 119, 644. scolytus, Scolytus.

Scopula emissaria, on jute in Bengal,

447.

Scots Pine (see Pinus sylvestris). Scots Pine Scale (see Toumeyella

numismatica). Scotussa rubripes, damaging pas-

tures in Argentina, **512.**Screens, experiments with, against

tobacco pests, 318.

scripta, Lina (see Melasoma lineatopunctata).

scrophulariae, Anthrenus.

scrupulosa, Teleonemia.

scutellare, Xanthogramma (Ischiodon).

scutellaris, Chirozetes (Phylaitis).

scutellatus, Gonipterus.

Scutellista cyanea (parasite of Saissetia oleae), in S. Africa, 385; introduced into California, 385, 386; establishment of, in Peru, 511; failure to establish, in Japan against Ceroplastes rubens, 172.

Scutigerella immaculata, destroyed by dichloroethyl ether in New

Jersey, 123.

Scutiphora pedicellata, measures against, on fruit trees, etc., in New South Wales, 290.

Scyllina, introduced parasite of, in Argentina, 308.

Scyllinops variabilis, in Argentina, 497,511; damaging lucerne, 511.

Scymnillus nunenmacheri, predacious on Pseudococcus nipae in Porto Rico, 52.

Scymnus, predacious on Pseudococcus comstocki in Brazil, 168; introduced into Porto Rico from Trinidad against Asterolecanium spp., 471.

Scymnus fenestratus, possibly predacious on Pseudococcus comstocki

in Palestine, 339.

Scymnus punctillum (see Stethorus).
Scymnus quadrimaculatus, not readily attacking Pseudococcus comstocki in Palestine, 339.

Sedge (see Carex).

Seed-corn Maggot (see Hylemyia cilicrura).

Seekay Wax R.93, 147.

segetum, Agrotis (Euxoa).

Seiulus, predacious on Phyllocoptes fockeui in Ontario, 306.

sejuncta, Microphthalma.

Selatosomus aeneus (see Corymbites). selecta, Autographa; Ptychomyia (Bessa).

selenaria, Boarmia (Ascotis).

Selenaspidus articulatus, liberation of Hyperaspis billoti on wild orange infested by, in Porto Rico, 471.

Selenium, introduction of, into plants against Aphids and mites, 165, 207, 275, 276. (See Selocide.) Selepa celtis, on mango in Ceylon,

594.

sellatum, Colasposoma.

Selocide, spraying with, against mites, 559, 639; question of risks of using, 559, 639; preparation of, 559.

Semiothisa fidoniata, parasite of, on Acacia catechu in India, 263.

Semiothisa granitata, on conifers in Quebec, 36.

semipunctata, Phoracantha. semistriatus, Microphanurus.

semivillosus, Amphimallus.

senatoria, Anisota.

Senecio jacobaea, Harpobittacus preventing establishment of Tyria jacobaeae against, in Australia, 143; establishment of Hylemyia jacobaeae against, in New Zealand, 334.

senex, Bruchoctonus.
senilis, Ceromasia.
sensoriatus, Myzus.
Sepetir (see Sindora).
septemfasciata, Nomadacris.
septempunctata, Coccinella.
septendecim, Magicicada.
Septis, on wild rice in Maine,
560.

Septis finitima, bionomics of, on wheat in Maine, 560.

Septobasidium pseudopedicellatum, Pseudococcus comstocki associated with, on Citrus in Brazil, 168.

seriatus, Psallus.

Serica, in forest nurseries in New York, 421.

sericariae, Sturmia (Crossocosmia). sericata, Lucilia.

sericeus, Metamasius.

sericiventre, Tetramorium.

serieventris, Podisus.

serinopa, Nephantis. serius, Eretmocerus.

Serraca infixaria (see Boarmia).

servatipes, Lophobaris.

serricorne, Lasioderma. sertifer, Neodiprion (Diprion).

Sesamia, introduced parasites reared on, in Mauritius, 610.

Sesamia cretica, measures against, on maize and sorghum in Sudan, 387, 388.

Sesamia inferens, on sugar-cane in India, 184, 268; parasite of, 268.

Sesamia uniformis, bionomics of, on sugar-cane in India, 184.

Sesamia vuteria, on Coix lachrymajobi in Ceylon, **594.**

Sesamum indicum, preferred by Schistocerca gregaria, **519**.

Sesbania, Anticarsia gemmatalis on, in Oklahoma, 252.

Setaria, Laevicephalus striatus on, in Russia, 617.

Setaria italica (Italian Millet), pests of, in Japan, 97, 100.

Setaria viridis, Delphacodes striatellus and mosaic disease of cereals in, in Siberia, 14.

Setora nitens, new parasites of, in Malaya and Java, 514.

setosa, Podomyia.

Seudyra subflava, on vines in China,

Sexava (on coconut), bionomics and control of, in New Guinea Territory and Netherlands Indies, 91, 379, 380.

Sexava nubila, parasites of, in Netherlands Indies, 380.

sexdens, Atta.

sexguttata, Anthia.

sexmaculata, Campsomeris.

sexmaculatus, Chilomenes; Scolo-

thrips; Tetranychus. sexnotata, Macrosteles (Cicadula).

sexta, Protoparce. seychellarum, Icerya. Seychelles, Coccids and their biological control on coconut, etc., in, 82, 323, 324; Melittoma insulare in, 340.

shastae, Cryptaspidiotus.

Sheep, toxicity of lead arsenate to, 44.

Sheep Dips, unsatisfactory for treating stacking sites against *Calandra*, **540**, **541**.

sheldoni, Eriophyes.

Shisham (see Dalbergia sissoo).

Shortleaf Pine (see Pinus echinata).

Shorts (see Bran).

Shrews, destroying Gilpinia polytoma, 38.

Siam, cultivation and pests of lac and its host-trees in, 94.

Siberia, Acridids in, 484–486; Delphacodes striatellus and mosaic of cereals in, 13; soil pests in, 577; beneficial insects in, 14, 485, 486. sibiricus, Aeropus.

Sida spp., Bemisia tabaci transmitting virus disease to tobacco from, in India, 183; cotton pests on, 387, 521, 522, 539.

sidae, Dysdercus.

side, Spilochalcis.

Sideridis unipuncta (see Cirphis). Sideroxylon, Achaea lienardi on, in

S. Africa, 19. sieboldi, Schizonotus.

sigillatus, Gryllodes.

signata, Plusia.

signipennis, Scirtothrips. silhetana, Terias blanda.

Silica, and tobacco, as diluent for pyrethrum dusts, 46.

Silkworm (see Bombyx mori).

Silpha opaca, factors affecting bionomics and control of, in Kola Peninsula, 577.

silvaticus, Pipunculus.

Silverfish (see Ctenolepisma, Lepisma and Thermobia).

Sima (see Tetraponera).

similis. Diprion.

simillima, Cryptognatha.

simiolus, Pleurotropis. simoni, Trissolcus.

simonovi, Microbracon (Habrobracon).

simonyi, Lygus.

simplex, Agromyza; Anabrus; Chilo; Taeniothrips.

simpsoni, Cydia (Carpocapsa) pomonella.

Sindora, damage to wood of, by Calotermes domesticus in Malaya, 543.

sinensis, Ceroplastes.

singularis, Euplectrus; Otiorrhynchus; Sahlbergella.

sinicus, Adoretus.

Sinoxylon sudanicum, parasite of, on cotton in Madras, **522**.

Sipalus formosanus, on pines in Formosa, 104.

Sipha flava (on sugar-cane), Coelophora inaequalis established against, in Porto Rico, 471; C. inaequalis introduced into U.S.A. against, 431.

Siphoninus phillyreae, parasitised by Encarsia partenopea in Britain,

374.

Sisalkraft, penetration of, by larvae of Lasioderma serricorne, 493.

sissoo, Disophrys.

Sisymbrium irio, Lygus spp. on, in U.S.A., 441.

Sitka Spruce (see *Picea sitchensis*). Sitodiplosis cambriensis, sp. n., on *Poa* spp. in Britain, 315; periodicity in oviposition of, 315.

Sitodiplosis dactylidis, on Dactylis glomerata in Britain, 315.

Sitodiplosis mosellana, on wheat in U.S.A., 254; periodicity in oviposition of, 315.

Sitodrepa panicea, in imported coriander seeds in Fiji, 496.

Sitophilus (see Calandra).

Sitotroga crealella, measures against, in stored wheat in Australia, 380; in Colombia, 9; bionomics of, on field and stored grain in Jugoslavia, 72; effect of harvesting machines on infestation by, in U.S.A., 254; use of eggs of, for rearing Trichogramma spp., 9, 87, 92, 192, 457; factors affecting parasitism of eggs of, by T. minutum, 553; Habrocytus cerealella parasitising stocks of, 193; tests of oils on eggs of, 585.

Skunks, destroying egg-pods of Anabrus simplex, 223.

slossonae, Spilocryptus (Agrothe-reutes).

smaragdina, Oecophylla; Nezara viridula.

Smicronyx utilis, sp. n., on Iva axillaris in Saskatchewan, **530.** smithi, Diversinervus; Lachno-

sterna (Phytalus).

Smoke, against Nysius vinitor, 31; against Schistocerca paranensis, 498.

Snake Gourd (see Trichosanthes anguina).

Symphoricarpus Snowberry (see vacemosus).

snyderi, Calotermes (Kalotermes).

Soap, spraying with, 169, 373, 603, 636, 639; for emulsifying oils, 231, 373, 435, 540, 585, 603, 653, 654; increasing injury to apple by oils, 585; use of dips containing, against Aphids on strawberry, 283, 375; in bait for Trypetids, 94; formulae containing, 25, 121, 145, 170, 233, 277, 279, 284, 291, 319, 330, 360, 373, 412, 435, 437, 540, 546, 603, 607, 636, 653.

Soap, Dolphin-oil, as an emulsifier for oils, 654.

Soap, Resin, spraying with, against Empoasca devastans, 517; unsatisfactory in sprays against Scirtothrips citri, 296.

Sod Webworms (see Crambus).

Soda (see Sodium Carbonate).

Soda, Caustic (see Sodium Hydroxide).

Sodium Arsenate, in baits and sprays against locusts, 461, 529, 612.

Sodium Arsenite, in bait-spray for Antestia, 589, 590; in baits, 19, 20, 107, 142, 169, 206, 300, 354, 428, 461, 529, 612, 613, 614; other poisons compared with, against locusts, 326, 612, 613, 614; aeroplane dusting with, against locust swarms, 156; in sprays against grasshoppers, 169; and limesulphur, logs sprayed with, against Monochamus, 312; toxicity of solution of, to Calandra oryzae, 540.

Sodium Benzoate, preventing mould in baits, 357.

Sodium Bicarbonate, tests of, in locust baits, 612, 613.

Sodium Carbonate, uses of, sprays, 25, 350, 437, 517.

Sodium Chloracetate, unsatisfactory in baits for termites, 19.

Sodium Chloride (see Salt).

Sodium Cyanide, soil treatment with solutions of, against pests, 168, 640, 642; used in pot method for generating hydrocyanic-acid gas (q.v.), 32, 145, 336.

Sodium Dinitro-cresylate, spraying with, against mites, 641. (See Elgetol.)

Sodium Fluoaluminate (see Cryo-

lite).

Sodium Fluoride, in baits, 19, 20, 304, 357, 406, 615; in spray against thrips, 414; as a timber preservative against Lyctus, 147; solution of, unsatisfactory against Calandra oryzae, 540.

Sodium Fluosilicate, tests of, in bait-sprays for Trypetids, 54, 55, 337, 494; effect of sub-lethal doses of, on Dacus, 494, 495; in baits, 20, 252, 269, 357, 362, 428, 461, 538, 576, 577; dusting with, 571, 576, 577, 584, 587, 654; oilimpregnated dusts of, 584, 654; spraying with, 654; formulae containing, 20, 54, 252, 269, 337, 362, 428, 461, 538, 577.

Sodium Hydroxide, in sprays against Coccids, 437, 605; little affecting distribution of nicotine between water and oils, 246.

Sodium Lauryl Sulphate, uses of,

in sprays, 215, 361, 634. Sodium Nitrate, used in producing smoke mixtures, 127.

Sodium Oleate, as a spreading and wetting agent, 283, 352, 260, 556; toxicity of, to eggs of Lygaeus kalmii, 352; effect of dips containing, on strawberry, 283.

Sodium Oleyl Sulphate, uses of, for conditioning dusts, 353; unin spray against satisfactory mites, 215.

Sodium Pentasulphide (see Sulfocide).

Sodium Phosphate, not improving

locust baits, 613. Sodium Selenate, 207, 275. Selenium.)

Sodium Sulphosuccinate, dioctyl ester of, as wetting agent, 595.

Sogata furcifera, on rice in Japan, 98, 101; effect of rainfall and temperature on, 98.

Sogata orizicola, on rice in Peru, 511. Soil, sampling and population studies of insects in, 396, 524; effect of character of, on pests, 342, 449, 579, 643.

Soil Erosion, grasshoppers as factor in, in Michigan, 204.

sokolowi, Telenomus.

solani, Empoasca; Macrosiphum (Aulacorthum).

solanifolia, Empoasca. solanifolii, Macrosiphum.

Solanine, 332.

Solanum, Dysdercus ruficollis on, in Peru, 539; relation of Epitrix spp. to species of, in U.S.A., 201, 221. (See Egg-plant and Potato.) Solanum auriculatum, Trypetids on,

in India, 524.

Solanum nigrum, Bemisia tabaci transmitting virus disease to tobacco from, in India, 183; extract of, as an insecticide, 332.

Solanum nodiflorum, 651.

Solanum trilobatum, virus disease of egg-plant transmitted to, in Madras, 542.

Solanum verbascifolium, relation of cacao pests to, in Nigeria, 516.

Solanum xanthocarpum, Jassid transmitting virus disease of eggplant to, in Madras, 542.

Solenopsis, associated with Stictocephala festina in Louisiana, 601. Solenopsis geminata, bionomics of, in Florida, 476; damaging plants in Hawaii, 35; associated with

Pseudococcus brevipes in Porto Rico, 346.

Solenopsis geminata xyloni, damaging telephone equipment in Texas, 260.

Solenopsis moelleri, measures against, associated with Pseudococcus comstocki in Brazil, 168.

Solenopsis molesta, Eciton nigrescens predacious on, in Florida, 476.

solidaginis, Macrosiphum.

Solomon Islands, Amblypelta cocophaga causing nutfall of coconuts in, 324, 325; Papuana on Colocasia esculentum in, 171; beneficial insects in, 325, 514; thrips introduced into, against Clidemia hirta, 495; Tetrastichus brontispae introduced into New Guinea Territory from, 379.

solstitialis, Amphimallus (Amphi-

mallon).

somervilli, Tachardiaephagus tachardiae.

Macrosiphum (Amphorosonchi, phora).

Sonchus asper, Nezara on, in Fiji, 279. Sooty Moulds, associated with Pseudococcus, 23, 48, 168; associated with other insects, 3, 32,

sophorae, Brassolis. sorbillans, Tricholyga. Sorbitol, as a larval attractant in sprays against Cydia pomonella,

Sorbus americana, Saperda candida on, in New York, 242; apple race of Rhagoletis pomonella developing on, 226.

sordida, Diplotaxis.

sordidus, Acanthocoris; Cosmopolites; Encoptolophus.

sorghicola, Contarinia.

Sorghum, 251; pests of, in China, 446; pests of, in India, 266, 271, **519, 520, 521, 523;** pests of, in Queensland, 466; pests of, in Sudan 387, 388; locusts damaging, in Uganda, 229; pests of, in U.S.A., 3, 165, 258, 277, 299; varieties of, in relation to pests 258, 299, 466; in crop rotation against Horistonotus uhleri, 602; avoided by Melanoplus spp., 603.

Sorghum halepense (Johnson Grass), relation of Euetheola rugiceps to, in Arkansas, 2; Phytomyza spicata on, in Fiji, 170; Paratetranychus indicus on, in Punjab,

520, 521.

soror, Diabrotica. Sour Clover (see Melilotus indica).

Sour-sop (see Annona muricata). Soy Beans, Plusia orichalcea on, in S. Africa, 514; pests of, in Ceylon, 594; Cydia glycinivorella on, in Far East, 33, 98; Jassid on, in India, 266; pests of, in Java, 492; pests of, in U.S.A., 3, 252, 254, 354, 599; in crop rotation against pests, 245, 251, 602; Pyrausta nubilalis not experimentally surviving on, 44; insecticides causing injury to, 599; chloropicrin not affecting stored seed of, 210. Soy-bean Flour, in bait for Otior-

rhynchus ligustici, 252; as an adhesive for arsenical sprays, 134, 371, 406, 633, 634.

Soy-bean Oil (see Oils, Vegetable). Spain, Acridids and Tettigoniids in, 459-461, 463, 529; Coccids in, 155, 457, 463; fruit-flies in, 457, 459, 461, 462, 464; miscellaneous pests in, 457, 462, 463; Myzus persicae and disease of Capsicum in, 456; Pyralid on rice in, 458; vine pests in, 458, 462: attempted biological control of pests in, 457, 459; Cryptolaemus montrouzieri introduced into Portugal from 63.

Spalangia (parasite of fruit-flies), breeding and liberation of, in Ceylon, 594; imported into Mauritius 610.

Spanish Cedar (see Cedrela).

Sparasion pilosum, egg-parasite of Anabrus simplex in Montana, 537.

Sparganothis pilleriana, arsenicals against, on grape vines in Spain, 458.

Sparrow Hawks, destroying noxious insects in Utah, 217.

Spartium junceum, insects on, in Peru, **511.**

Spathius canadensis, parasite of Dendroctonus frontalis in Virginia, **211.**

Spathius critolaus, hosts and ecology of, in India, 268, 522.

Spathius labdacus, ecology of, parasitising *Pempherulus affinis* in Madras, **522.**

Spathius piperis, parasite of Lophobaris piperis in Netherlands Indies, 33.

vulnificus (parasite of Spathius Dinoderus), possibility of utilising, against Pempherulus affinis in India, 268; bionomics of, 269. speciosa, Orphulella.

speciosissima, Centrodora;

(see P. aeruginosa). speciosus, Microterys.

spectabilis, Coccophagus; Dendro-Phanomeris (see Rhogas limus; dendrolimi).

speculum, Ricania.

Sphaceloma fawcetti, 597.

Sphaerotrypes, not infesting barked

logs in Burma, 78.

Sphegigaster flavicornis (parasite of Phytomyza ilicis), in Britain, 417; establishment of, in Br. Columbia, 417.

sphenotoma, Aeolarchis.

Sphinx pinastri, on pine in Latvia and Poland, 86.

Spicaria fumosorosea, experimental utilisation of, against Sturmia sericariae in Japan, 88.

spicata, Phytomyza.

Spiders, destroying noxious insects,

105, 442, 443, 517.

Spilochalcis albifrons, parasite of Chalepus dorsalis in U.S.A., 354. Spilochâlcis delira, S. side recorded as, 190; synonym of S. flavopicta, 190.

Spilochalcis flavopicta, synonymy of, 190.

Spilochalcis igneoides, synonymy of, 190.

Spilochalcis leptis, sp. n., hosts of, in U.S.A., 190.

Spilochalcis side (torvina), parasite of Apanteles congregatus in N. Carolina, 219; synonymy of, 190; recorded as S. delira, 190. Spilochalcis vittata (see S. igneoides).

Spilocryptus japonicus, parasite of Margaronia pyloalis in Japan,

Spilocryptus slossonae, parasite of Gilpinia polytoma in Maine, 637. Spilonota holotephras (on guava), in Fiji, 495; in Guam, 332.

Spilonota ocellana (on fruit trees), bionomics of, in Baluchistan, 267; in U.S.A., 127, 128, 190, new parasite of, 190; measures against, 127, 128.

Spinach, pests of, in U.S.A., 46, 152, 158; Thrips tabaci transmitting virus disease to, 140; Lepinotus patruelis infesting imported seed of, in California, 416.

Spindle Oil, 653, 654. spinicollis, Myllocerus. spinipennis, Coleoxestia.

spinipes, Tarsonemus.

spinosum, Lecanium persicae. Spirochaeta pieridis, sp. n., Pieris brassicae in France, 224.

Spizella passerina arizonae, destroying noxious insects in Utah, 563. splendens, Chrysoplatycerus; terocerus.

Spodoptera exigua (see Laphygma). Spodoptera mauritia, in Ceylon, 620; on rice in Fiji, 495; on rice and grasses in Guam, 331; parasites of, **331**, **332**, **620**.

Spraying, types of equipment for, 152, 202, 232, 241, 271, 312, 569, 638, 654; use of aircraft for, 256, 438; dusting compared with, 42, 135, 197, 202, 296, 297, 463, 466.

Sprays, factors affecting deposits and retention of, on apples, 206, 371; study of dispersing agents for, 112; base oil affecting efficiency of atomised, 398; apparatus and technique for testing, 195, 311, 557, 591.

Spredite, 559.

Spruce (Picea), 603; effect of felling of, on pests in Britain, 382; pests of, in Canada, 36, 37-39, 294, 308, 348, 424, 431, 474, 509, 530, 531, 532, 626, 637; Gilpinia polytoma on, in Czechoslovakia and Finland, 474; new Rhynchites on, in Japan, 40; pests of, in Latvia, 85, 86; G. polytoma on, in Rumania and Sweden, 474; Chermes spp. on, in Switzerland, 67; pests of, in U.S.A., 37, 396, 422, 424, 427, 507, 531, 537, 552, 607, 637; susceptibility of species of, to Chermes cooleyi, 537.

Spruce, Black (see *Picea mariana*). Spruce, Norway (see *Picea abies*).

Spruce, Red (see Picea rubra).

Spruce, Sitka (see *Picea sitchensis*). Spruce, White (see *Picea glauca*). Spruce Budworm (see *Harmologa fumiferana*).

Spruce Sawfly, European (see Gil-

pinia polytoma).
Spruce Trunk Aphis (see Chermes pineoides).

squamosus, Hypomeces.

Squash, Engytatus geniculatus on, in Hawaii, 34, 650; pests of, in U.S.A., 163, 201, 405, 443, 501, 596, 603; fumigation against introduction of Trypetid into U.S.A. from Hawaii in fruit of, 499; insects and virus diseases in, 57, 501; calcium cyanide dust causing injury to, 201.

Squash Bug (see Anasa tristis). Stable Fly (see Stomoxys calcitrans). stabulans, Lydella; Muscina.

Stachytarpheta dichotoma, reservoir of tobacco virus disease in Sumatra, 318.

stali, Euborellia.

stanleyi, Metaphycus.

Starch, attraction of, for *Thermobia domestica*, **401**; (in timber), determination and depletion of, in relation to *Lyctus*, **150**, **466**.

Stathmopoda melanochra, liberation of, against Eriococcus coriaceus in New Zealand, 545.

Stauroderus scalaris (in Asiatic Russia), Empusa grylli infesting, 570; effect of climate on outbreaks of, 485, 486.

stebbingi, Drosicha.

Stellaria media, favourable food for Silpha opaca, 577.

stemmator, Xanthopimpla.

Stenotaphrum, insects damaging lawns of, in Texas and Porto Rico, 214, 217.

Stephanitis typica, not transmitting mosaic of Musa textilis in Philippines, 317.

Stephanoderes hampei (Coffee Berry Borer), ant destroying, in Brazil, 292; failure to breed *Prorops nasuta* against, in Ceylon, 594; review of literature on natural enemies of, 292.

Sterculia, Dysdercus spp. on, in Sudan, 387.

Sterculia apetala, Pseudococcus nipae on, in Porto Rico, 52.

Sterculia platanifolia, Aonidiella aurantii on, in Japan, 102.

sterculiae, Psylla; Tepperia; Tyora. Stethorus punctillum, predacious on hop pests in Britain, 282.

Stevensonia, Melittomma insulare associated with, in Seychelles, 341.

sticticalis, Loxostege.

sticticollis, Agalliana (Agallia). sticticraspis, Diatraea (Argyria).

Stictocephala festina, bionomics of, on leguminous plants in Louisiana, 601.

Stictocephala inermis, effect of sprays on eggs of, on fruit trees in U.S.A., 216.

stigma, Megaderus.

Stigmaeus, measures against, damaging turf in S. Africa, 615. stigmaterus, Microdus (Bassus). stigmatias, Euxesta.

Stilpnotia salicis, on poplar and

willow in Quebec, 626.

Stizolobium, pests of, in U.S.A., 252, 428, 599; in crop rotation against wireworms, 253, 602; insecticides causing injury to, 599.

Stock (see Matthiola).

stokesi, Antrocephalus (Stomato-ceras).

Stomatoceras, hyperparasite of Artona catoxantha in Malaya, **79.** Stomatoceras melitarae (see Antro-

cephalus).
Stomatoceras stokesi (see Antrocephalus).

Stomatoceras sulcatiscutellum, parasite of Nephantis serinopa in India, 620.

Stomatomyia bezziana (parasite of Nephantis serinopa), bionomics and value of, in Ceylon, 620, 621.

Stomatorrhina lineata, egg-parasite of Locusta in Uganda, 229.

Stomoxys calcitrans, straw in fields favouring, in U.S.A., 254.

strachani, Pinnaspis minor. strangulatus, Catolellerus.

Strategus quadrifoveatus, method of treating young coconut palms against, in Porto Rico, 472.

Strategus titanus, on sugar-cane in

Jamaica, 293.

Strawberry, Lagria villosa on, in S. Africa, 19; pests of, in Australia, 466, 494; pests of, in Britain, 27, 28, 146, 283, 374, 375; Lamellicorns on, in Canada, 90, 508; Amphimallus majalis on, in France, 24; Galerucella distincta on, in Korea, 152; Myzus porosus on, in Mexico, 46; pests of, in U.S.A., 2, 46, 258, 298, 362, 530, 536, 537, 549, 554, 602, 633, 643; Capitophorus fragariae and virus diseases of, 27; problem of lead arsenate spray residues on, 633; insecticides causing injury to, 259, 554, 633.

Strawberry Aphis (see Capitophorus

fragariae).

Strawberry Crown Borer (see Tyloderma fragariae).

striatalis, Diatraeophaga.

striatellus, Delphacodes (Delphax). striatum, Anobium (see A. punctatum).

striatus, Laevicephalus (Deltoceph-

alus).

Striglina scitaria, on Xylia dolabriformis in Burma, 78.

strobi, Chermes (Pineus); Cinara; Pissodes.

strobilella, Cydia (Laspeyresia). Stromatium longicorne (in timber), in Burma, 78; bionomics of, in

Loochoo Islands, 103.

Strongylurus decoratus, on Araucaria cunninghami in Queensland, 467.

Strophostyles spp., Chalcodermus aeneus on, in U.S.A., 405.

Strumeta (see Dacus). stupidus, Rhizotrogus.

Sturmia macrophallus, parasite of Semiothisa fidoniata in India, 263.

Sturmia sericariae (parasite of Bombyx mori), experimental utilisation of Spicaria fumosorosea against, in Japan, 88.

Stylocryptus profligator, parasite of Dendrolimus in Sakhalin, 95.

Stylocryptus subclavutus, parasite of Neodiprion tsugae in Oregon, **552**. Stylocryptus vulgaris, parasite of Gilpinia polytoma in Maine, **637**.

suahelorum, Podagrica (Nisotra).

suavis, Rhagoletis. subclavatus, Stylocryptus. subflava, Seudyra.

subfuscus, Phygadeuon.

submarginatus, Hemiteles (Astomaspis); Xyleborus.

subspinosus, Macrodactylus.

subterranea, Feltia.

Subterranean Clover (see Trifolium subterraneum).

Subterranean Grass Caterpillar (see Oxycanus).

subtruncatus, Elytroteinus.

succincta, Epicampocera; Patanga. Sudan, Anglo-Egyptian, miscellaneous pests in, 386–388; parasites of Lepidoptera in, 386, 514.

sudanensis, Hercothrips. sudanicum, Sinoxylon.

Sugar, in baits, 91, 99, 100, 142, 183, 252, 253, 267, 406, 414, 430, 435, 612, 613; in bait-sprays for Trypetids, 54, 55, 337, 494; addition of, to dusts and sprays against Rhagoletis completa, 561, 562; as larval attractant in sprays against Cydia pomonella, 129; in sprays against mites, 214, 215; in sprays against thrips, 121, 122, 204, 214, 277, 296, 297, 407, 413, 414, 491, 641; in dusts against Scirtothrips citri, 297; used for feeding insects, 98, 101, 102, 174, 225, 264, 310, 394; types of, 129, 413, 414, 561, 562.

Sugar Pine (see Pinus lambertiana). Sugar-cane, Diatraea saccharalis on, in Brazil, 4, 5, 470; pests of, in Ceylon, 609, 610; pests of, in Colombia, 8, 9; pests of, in Formosa, 92, 94, 104, 172, 622; pests of, in Guana, 331; Diatraea spp. on, in Br. Guiana, 538; pests of in Hawaii 35 538; pests of, in Hawaii, 35, 49, 50, 273, 274; pests of, in India, 81, 137, 183, 184, 266, 271, 272, 520; pests of, in Java, 92, 524; pests of, in Mauritius, 83. 609, 610; D. saccharalis on, in Peru, 511; pests of, in Queensland, 388, 389, 622; pests of, in U.S.A., 160, 354, 428, 431, 466, 471, 477, 599, 601; pests of, in W. Indies, 3-5, 6, 7, 52, 160, 263, 293, 470, 471, 498, 601; susceptiiblity of varieties of, to pests, 81, 272, 293, 389, 521, 601, 609; Aphis maidis transmitting mosaic of, 35, 523.

Sugar-cane Aphis (see Sipha flava).

Sugar-cane Beetle Borer (see Rhabdocnemis obscura).

Sugar-cane Bud Worm (see Decadarchis flavistriata).

Sugar-cane Froghopper (see Tomaspis saccharina).

Sugar-cane Leafhopper, Indian (see Pyrilla perpusilla).

Sugar-cane Moth Borer (see Diatraea saccharalis).

Sugar-cane Stalk Mite (see Tarsonemus spinipes).

sulcatiscutellum, Stomatoceras.

sulcatus, Dendrosoter; Otiorrhynchus (Brachyrrhinus); Scolytus. sulciventris, Rhoecocoris.

Sulfocide, for protecting timber against *Lyctus*, **448**.

Sulphite Lye, as an emulsifier, 110, 191, 644.

Sulphur, tests of toxicity to Empoasca fabae of, 325, 326, 440, 441; against other Rhynchota, 21, 258, 510, 597, 600, 638, 639; favouring infestation of vines by Pseudococcus adonidum, against Malacosoma americana, 118; against mites, 23, 32, 135, 164, 224, 249, 276, 321, 565, 584, 597, 602, 636, 637, 639, 641, 648; against thrips, 18, 25, 295, 296, 297; tests of action on honey bees of, 549; as a repellent, 21, 90, 643; equipment for fumigating empty greenhouses and mushroom houses with, 276; arsenic, for fumigating nests of Atta and termites, 169, 338; dusting with, 18, 21, 32, 90, 118, 135, 164, 224, 295, 296, 297, 321, 325, 326, 441, 510, 565, 584, 597, 598, 600, 602, 636, 638, 639, 648; in mixed dusts, 10, 25, 46, 47, 52, 135, 136, 248, 249, 258, 301, 325, 326, 358, 359, 405, 430, 441, 510, 600, 639, 640, 647; spraying with, 21, 23, 162, 295, 296, 440, 441, 534, 584, 597, 598, 637, 638, adhesive and wetting 641; agents for, 162, 249, 534; in mixed sprays, 21, 135, 249, 296, 297, 597, 602, 634, 638, 639; forms of, 23, 295, 296, 441, 584; formulae containing, 10, 21, 23, 25, 52, 135, 136, 249, 258, 296, 297, 301, 325, 358, 359, 405, 441 510, 600, 602, 634, 638, 639, 640; and injury to plants, 285, 296, 405; rendering tomatos unfit for canning, 22; toxicity of organic compounds containing, to Lucilia sericata, 407, 408.

Sulphur-lime, Dry-mix, formula for, against Empoasca fabae, 441.

Sulphuric Acid, and cyanides, for generating hydrocyanic-acid gas (q.v.), 336, 455.

Sultanas, Oryzaephilus surinamensis reared on, 591. (See Fruit, Dried.)

Sumatrol, 135.

Sun Spots, correlation of insect outbreaks with, 63, 538.

Sunflower, Isaniris ater on, in Nyasaland, 62; experiments with Pyrausta nubilalis and, 44; Trogoderma versicolor infesting stored seed of, 453.

Sunn Hemp (see Crotolaria juncea). Sunt Tree (see Acacia arabica).

superans, Dendrolimus.

superstitiosus, Dysdercus.

Sura uncariae, sp. n., on gambier in Sumatra, 105.

suranus, Microphanurus. surcularius, Platystolus.

surinamensis, Oryzaephilus; Pycnoscelus.

suspensa, Anastrepha.

Susumia exigua, parasite of, on rice in Guam, 331.

suturalis, Brumus.

suzukii, Drosophila. Swede, Myzus persicae wintering

on, in Eire, 329.

Sweden, parasite and forms of Gilpinia polytoma in, 294, 474; Coleoptera damaging building timbers in, 324.

Sweet Clover (see Melilotus).

Sweet Gum (see Liquidambar styraciflua).

Sweet Pepper (see Capsicum).

Sweet Potato, Typophorus nigritus on, in Argentina, 512; Euscepes postfasciatus on, in Bermuda, E. postfasciatus on, in Brazil, 511; Longitarsus spp. on, in Crimea, 479; Hippotion celerio on, in Fiji, 170; cephala tuberculata on, in India, 270; Acanthocoris sordidus on,

in Japan, 171; Cylas formicarius associated with fungus on, in Loochoo Is., 97; C. formicarius on, in Louisiana, 415, 602; H. celerio on, in New Guinea Territory, 377; pests of, in Nyasaland, 154, 589; pests of, in Porto Rico, 261, 346; locusts

damaging, in Uganda,

(tubers), methyl bromide fumigation of, against *C. formicarius*, **413**; (dried), *Calandra oryzae* in, in Japan, **173**.

Sweet Potato Weevil (see Cylas

formicarius).

Swietenia (see Mahogany).

Swintonia floribunda, Coleoptera attacking timber of, in Burma, 78.

Switzerland, Curculio nucum on hazel in, 376; orchard pests in, 66, 617; Chermes spp. on spruce in, 67; mass breeding and natural enemies of Tineola biselliella in, 375, 376; new Acarapis on honey bee in, 528.

sydneyensis, Anarhopus.

Sylepta clytusalis, bionomics of, on Brachychiton populneus in New

South Wales, 467.

Sylepta derogata, susceptibility of varieties of cotton to, in China, 446; on balsa in Formosa, 104; parasites of, in India, 524, 620; measures against, on cotton in Philippines, 649; on Hibiscus esculentus, 524.

sylvaticus, Pamphilius.

syma, Chrysocharis.

Sympherobius amicus, attempted utilisation of, against Pseudococcus comstocki in Palestine, 74, 339.

Symphoricarpus racemosus, experiments with forms of Rhagoletis pomonella and, in Nova Scotia, 226.

Sympiesis bimaculata, parasite of Parornix prunivorella in Kansas, 191.

Synanthedon (see Aegeria).

Synaplomyia galerucellae, parasite of Galerucella luteola in California, 417.

Synedrella nodiflora, reservoir of tobacco virus diseases in Sumatra, 318.

Synnematium jonesii, infesting Promecotheca papuana in New Guinea Territory, 379.

Synonycha grandis, liberation of, against Oregma lanigera in Formosa, 93.

Synopeas, parasite of Contarinia medicaginis in Hungary, 12.

Syntomosphyrum, parasite of Mayetiola destructor in Russia, **618, 619.** Syntomosphyrum indicum, utilisation of, against fruit-flies in Fiji,

170, 495.

Syntomosphyrum obscuriceps, sp. n., hyperparasite of Artona catoxantha in Java and Malaya, 79; parasite of Stomatomyia bezziana in Ceylon, 621.

Syntomosphyrum zygaenarum, hyperparasite of Artona catoxantha in

Java, 79.

INDEX.

Syntomosphyrum zygaenarum var. nigricoxis, n., hyperparasite of Artona catoxantha in Java, 79.

Syringa (see Lilac). syringella, Gracilaria.

Syrrhizus delusorius, parasite of Anthonomus pomorum in Germany, 70.

Syrup, in baits, 107, 430; as a larval attractant in sprays against *Cydia pomonella*, 129; in tartar emetic sprays against thrips, 413.

Systasis encyrtoides, parasite of Contarinia medicaginis in Hun-

gary, 12.

Systates hargreavesi, sp. n., on coffee and Markhamia in Uganda, 341.
Systena taeniata, on potato in New Mexico, 638; measures against, 639.

Systoechus autumnalis, parasite of Acridids in Siberia, 485.

Systoechus vulgaris, destroying grasshopper eggs in Canada, 280, 308, 309; introduced into Argentina, 308; immature stages of, 280.

Systole coriandri, factors affecting infestation of coriander by, in

Russia, 576.

T.

tabaci, Bemisia; Empoasca; Thrips. tabacum, Hypopteromalus.

tabidus, Trachelus.

tabonus, Chelonus.

tachardiae, Eupelmus; Tachardiaephagus.

Tachardiaephagus tachardiae (with var. somervilli), parasite of Laccifer lacca in India, 263.

Tachardiella argentina, new parasite of, in Argentina, 344.

Tachina civiloides, parasite of Lepidoptera in India, 74, 263; utilisation of, 74.

Tachinids, supplements to revision of British, 328, 592; records of, in Germany, 70; puparia of Indian species of, 480; list of, in S. Rhodesia, 655.

tachypus, Mesochorus.

taeniata, Systena.

taeniatulus, Pantomorus.

Taeniotes scalaris, on figs and Artocarpus in Brazil, 391, 392.

Taeniothrips inconsequens, on pear and prune in U.S.A., 211, 223, 261, 418; not attacked by Leptothrips mali, 211; against, 261, 418. measures

Taeniothrips simplex, measures against, on gladiolus and carnations in U.S.A., 136, 204, 214, 277,

413, 641.

Taeniothrips usitatus var. cinctipennis, on beans in Queensland, 466.

Tagetes minuta, oil of, in baits for locusts, 612.

talaca, Hyposidra.

Talc, as a carrier for bacterial spores, 125, 504; as carrier for dusts, 22, 46, 247, 249, 250, 278, 290, 296, 301, 325, 326, 353, 404, 405, 504, 557, 569, 596, 597, 630, 631, 647; pH and electrostatic charge in relation to toxicity of dusts containing, **301, 569.** Tamarindus indica, Bruchid

stored pods of, in India, 523.

tanaceti, Galeruca.

Tanaomastix abnormis (see Leptomastidea).

Tanganyika Territory, coffee pests and their parasites in, 175-177, 514, 589, 590; Argyroplace on cotton in, 227; mite infesting Locusta migratoria migratorioides in, 177; termites in, 590; locust migrating to N. Rhodesia from, 491; Coccinellids introduced into Seychelles from, 324.

Tangerine Orange (see Orange,

Mandarin).

Tapeigaster marginifrons, Dacus confused with, in New South Wales, 237.

tapetzella, Trichophaga.

Tapinoma melanocephalum, measures against, in houses in Austra-

lia, 143.

Tar Distillates, as dormant and delayed dormant sprays against orchard pests, 70, 108, 128, 134, 144, 155, 191, 194, 231, 232, 248, 284, 585, 653; effects of, on other Arthropods overwintering in apple bark, 70; causing injury to apple, 191, 194, 284, 585; use of, against columbine and iris

borers, 644; treatment of timber with, against termites and Lyctus, 138, 147; value of, against Scolytids in elm logs, unsatisfactory against Calandra oryzae, 540; studies of action of, on insects and insect eggs, 239, 281; (and derivatives), dusts containing, as insecticides, 247; types of, 70, 128, 134, 191, 231, 232, 239, 248, 281, 585, 653; formulae for, 108, 128, 144, 155, 191, 194, 232, 248, 284, 585, 644; emulsifiers for, 248, 644, 653; and lime-sulphur, 191; and mineral oils, 108, 128, 191, 194, 195, 248, 284, 585; and paraffin wax, 134; review of history of, as insecticides, 651.

taragamae, Apanteles.

Taraxacum kok-saghyz, insects attacking in Russia, 484.

Taro (see Colocasia).

Taro Leafhopper (see Megamelus proserpina).

Tarsonemus latus, bionomics of, in S. Africa, 320; on Capsicum and ivy in California, 223, 224; on papaya in Hawaii, 650; measures against, 224, 321.

Tarsonemus pallidus, on strawberry in Britain, 28, 283.

Tarsonemus spinipes (Sugar-cane Stalk Mite), thrips predacious on, in Hawaii, 273.

Tartar Emetic, in bait for ants, 430; in bait-sprays for Trypetids, 55, 337, 561, 562; tests of toxicity and sub-lethal doses of, on Dacus ferrugineus tryoni, 55, 495; and sugar, spraying with, against thrips, 121, 122, 204, 214, 277, 296, 297, 407, 413, 414, 491, 641; substitutes for sugar in sprays containing, 214, 215, 296, 413; preparation and tests of dusts containing, against thrips, 296, 297; in sprays against mites, 214, 215; question of risk of, to bees, 407.

Tarucus theophrastus, parasite of, in India, 264.

Tasmania, orchard pests in, 150, 151, 152, 291; Aphodius tasmaniae damaging pastures in, 547; Pieris rapae in, 467; natural enemies and biological property in 149, 151 control of insects in, 142, 151, 152, 291, 468, 545; Apion ulicis imported into, against Ulex

europaeus, 334; introduction of parasites of Tortrix postvittana into New Zealand from, 151, 152.

tasmaniae, Aphodius. tasmanica, Apanteles.

Tea, pests of, in Ceylon, 60, 173, 174, 175, 381; Zeuzera coffeae on, in Malaya, 513; pests of, in Nyasaland, 59, 60, 61, 154, 588; probable virus disease of, 175.

Tea Tortrix (see Homona coffearia). Teak, pests of, in Burma and India, 75, 76, 77; Zeuzera coffeae

on, in Java, 270.

Technomyrmex detorquens (albipes), attacking Cactoblastis cactorum in S. Africa, 17; measures against, in houses in Australia, 142; attacking Coccinellid predacious on Icerya in Seychelles, 82.

Tectocoris diophthalmus (lineola), on Citrus in Fiji, 389; on cotton in Philippines, 649.

Tectona grandis (see Teak).

tectonae, Nemeritis.

tectus, Ptinus.

tegularis, Brachymeria. telarius, Tetranychus.

Telenomus beneficiens, parasite of Diatraea venosata in Formosa and Java, 92; effect of rearing, on Chilo infuscatellus, 92.

Telenomus dendrolimusi, parasite of Dendrolimus in Sakhalin, 95.

Telenomus dilophonotae, parasite of Erinnyis ello in Colombia, 9.

Telenomus ichthyurae, parasite of Datana integerrima in Missouri, 167.

Telenomus nawai, hosts and utilisation of, in Hawaii and Guam, 49, 382.

Telenomus ochus (parasite of Oreta carnea), bionomics of, in Sumatra, 195.

Telenomus politus, parasite of Eurygaster integriceps in Russia, 582.

Telenomus sokolowi, parasite of Eurygaster integriceps in Russia, 575, 582.

Telenomus usipetes, parasite of Hapalia machaeralis in Burma, 77.

Teleonemia australis, on olive in Nyasaland, 589.

Teleonemia scrupulosa (lantanae), utilisation of, against Lantana in Queensland and Pacific Is., 289, 467, 496; suggested introduction of, into India, 289. Teleopterus violaceus, parasite of Agromyza on teain Nyasaland, 588.

Telephone Equipment, ants causing damage to, in Texas, 260. (See Cables, Electric.)

Telsimia nitida, predacious on Aspidiotus destructor in Guam, 331; establishment of, against Pinnaspis buxi in Hawaii, 331.

Temperature, relations of locusts and grasshoppers to, 48, 141, 157, 205, 223, 341, 342, 343, 394, 460, 484, 485, 487, 520, 570, 583, 589, 118, 128, 141, 145, 149, 151, 155, 156, 158, 159, 160, 161, 163, 172, 173, 174, 180, 181, 185, 186, 202, 218, 224, 234, 245, 308, 313, 315, 321, 329, 333, 374, 375, 376, 395, 402, 403, 418, 420, 426, 427, 428, 439, 442, 443, 448, 451, 452, 453, 454, 455, 456, 478, 515, 527, 531, 553, 562, 566, 567, 571, 572, 573, 574, 575, 578, 579, 580, 581, 582, 590, 591, 595, 597, 601, 602, 621, 622, 627, 638, 650; effects of constant and varying, on insect metabolism, 131; affecting development of milky diseases in Popillia japonica, 369; affecting toxicity of stomach insecticides, 356; apparatus for regulating, 88, 152.

Tenebrio molitor, reactions to humidity of, 224.

Tenebrio picipes, in stored cereals in U.S.A. and Japan, 506.

tenebrionis, Capnodis.

Tenebroides mauritanicus, in stored cereals in Kansas, 506; seasonal variation in resistance of, to heat, 506.

tenellus, Eutettix.

Tennessee, Dendroctonus frontalis on pines in, 49; Heliothis armigera and its natural enemies in, 534, 535; Melanoplus femur-rubrum in, 604.

tenthredinis, Mesoleius.

tenuibasalis, Hemiteles (Microtorus). Tenuipalpus australis, sprays against, on Passiflora edulis in Queensland, 465.

Tenuipalpus californicus, sprays against, on Passiflora edulis in Queensland, 465.

Tenuipalpus obovatus, on tea in Ceylon, 175.

INDEX. 803

tenuis, Engytatus; Heterotermes. Teosinte, Annual (see Euchlaena mexicana).

Tephrosia, rotenoids in, 135; preparation of extracts of, 586.

Tephrosia candida, value of cultivating, under Citrus against Diaprepes, 262.

Tephrosia hookeriana, preferred by Schistocerca gregaria in India,

Tephrosia macropoda, insecticidal

properties of, 483.

Tephrosia toxicaria, Lithocolletis on, in Porto Rico, 473; effect of defoliation on constituents of, 470.

Tephrosia virginiana, insecticidal properties and constituents of,

397, 398, 555.

Tephrosia vogeli, Lepidoptera on, in Porto Rico, 345, 473; insecticidal properties of, 483; extract of, ineffective against Antestia, 230.

Tepperia sterculiae, on Brachychiton populneus in New South Wales,

467.

Teragra guttifera, measures against, in forests in S. Africa, 18.

Terastia meticulosalis, on Erythrina berteroana in Porto Rico, 469.

Tergitol 7 Penetrant, 136.

Terias blanda silhetana, on Cassia in Burma, 78.

Termes badius, habits of, in Kenya, 338.

Termes bellicosus (see Macrotermes). Termes hispaniolae, imported into Barbados in wallaba wood, 6.

Termes natalensis (see Macrotermes). Termes transvaalensis, habits of, in Kenya, 338.

Termes vulgaris, attacking tung trees in Nyasaland, 588; sures against, 589.

terminalis, Euproctis. terminifera, Chortoicetes.

Termites, of E. Africa, 590; of Barbados, 5-7; of Katanga, 611; attacking living plants, 6, 7, 19, 62, 387, 588, 611; damaging grass, 338, 615; attacking timber, 6, 7, 19, 138, 142, 333, 400, 483, 537, 543, 610, 611; timbers resistant to, 7, 138, 157, 611; factors influencing population complex of, 396; comparison of mounds of, with adjacent soil, 592; ants destroying, 499; measures against, 6, 7, 19, 138, 142, 338, 387, 483, 589, 610.

Terpene Alcohols, wetting agent containing, 595.

Tessaratoma javanica, on Schleichera trijuga in India, 263, 264; parasite of, 263.

tessellatus, Eucalymnatus.

testaceipes, Lysiphlebus.

testaceus, Coptotermes; Hoplovrhinus; Paniscus; Xyleborus. testudinea, Hoplocampa.

testulalis, Maruca.

Tetrachlorethane, ineffective as a greenhouse fumigant against

Tortrix pronubana, 146.

Tetracnemus peregrinus (parasite of Pseudococcus adonidum), Brazil, 298; introduction and utilisation of, in California, 298, **355**; parasites of, **355**.

Tetragonia expansa, 140.

Tetramorium guineense, utilisation of, in Formosa, 93.

Tetramorium sericiventre continentalis, damaging turf in S. Africa, 615.

Tetranychus, on beans and papaya Hawaii, 650; measures against, on cotton in Russia and Philippines, 584, 648.

Tetranychus atlanticus, sp. n., on strawberry and clover in U.S.A.,

530.

Tetranychus bimaculatus (see T. telarius).

Tetranychus bioculatus (see Paratetranychus).

Tetranychus cucurbitae, sp. n., foodplants of, in Punjab, 271.

Tetranychus fici, on figs in Punjab, 271.

Tetranychus mori, sp. n., on mulberry in Punjab, 271.

Tetranychus opuntiae, attacking prickly-pear in America, 587; introduced into Australia, 587.

Tetranychus pacificus, outbreak of, in Br. Columbia, 330; in U.S.A., 330, 559, 561, 639, 641; on apple, 559, 639, 641; measures against, 330, 559, 561, 639, 641; characters distinguishing T. telarius and, 444.

Tetranychus sexmaculatus, tests of dusts against, in California, 561.

Tetranychus telarius, on strawberry in Britain, 28; on banana in Jamaica, 445; in Punjab, 271; on date palms in Sudan, 388 and experiments measures against, on miscellaneous plants in U.S.A., 165, 207, 214, 249, 275,

276, 535, 555, 561, 602, 636; characters distinguishing T. pacificus and, 444.

Tetraponera, predacious on Pempherulus affinis in Madras, 522. tetrarhodus, Capitophorus (Pentatrichopus).

Tetrastichodes brontispae (see Tetra-

stichus).

Tetrastichus, parasite of Hapalia machaeralis in Burma and India, 77; liberation of, against Bruchus brachialis in N. Carolina, 48; probably hyperparasitic on Phytolyma lata in Gold Coast, 492; hyperparasite of Artona catoxantha in Malaya, 79; para-site of Diaprepes in St. Lucia, 262; parasite of Tineola biselliella in Switzerland, 376.

Tetrastichus ayyari, bionomics and laboratory breeding of, in India,

Tetrastichus brevistigma, 417.

Tetrastichus brontispae (parasite of Brontispa longissima), introduced into New Guinea Territory, 379.

Tetrastichus carinatus, parasite of Mayetiola destructor in Russia, 619.

Tetrastichus coerulescens, parasite of Parornix prunivorella in Kansas, 191.

Tetrastichus giffardianus (parasite of Ceratitis capitata), establishment of, in Brazil, 344; not parasitising indigenous fruit-flies in Brazil, 344; attempted utilisation of, against fruit-flies in Fiji, 495; suggested introduction of, into Spain, **459**.

Tetrastichus injuriosus, hyperparasite of Saissetia oleae in Africa, 386. Tetrastichus purpureus, parasite of Laccifer lacca in India, 264.

Tetrastichus pyrillae, parasite of Pyrilla in India, 82, 184.

Tetrastichus vapo, parasite of Apan-

teles spp. in Britain, 314. Tettigonia viridissima, larvae of Pieris brassicae poisonous to, 489. texana, Anomis (see A. luridula).

Texas, cotton pests in, 198, 213, 257, 258, 404, 430, 499, 500; Chlovochroa ligata on flax in, 499; Lachnosterna spp. in, 213, 214; miscellaneous pests in, 47, 49; Acrobasis caryae on pecan in, 133; ants damaging telephone equipment in, 260; beneficial insects in, 190, 477, 500.

Textiles, insects damaging, 137, 164, 216, 255, 335, 363, 375, 644.

Thallium Acetate, in baits for ants, 430.

Thallium Sulphate, in baits: for ants, 430; for locusts, 614.

Thamnurgides cardamomi (see Poecilips)

theae, Fiorinia.

Thecodiplosis mosellana (see Sitodiplosis).

Thelaira nigripes, parasite of Diparopsis castanea in Nyasaland, 153.

themisto, Thyridia. theobaldi, Thomasiniana.

theobroma, Sahlbergella.

theobromae, Helopeltis; Leucothrips. theophrastus, Tarucus.

Theresia ampelophaga (see Theresimima).

Theresia claripalpis (parasite of Diatraea saccharalis), in Brazil, 4, 470; introduction of, into Barbados and Porto Rico, 4, 5, 470, 471; in Trinidad, 471.

Theresimima ampelophaga, banding against, on vines in Palestine,

623.

Theretra japonica, parasite of, in Japan, 40.

theretrae, Microplitis.

Thermobia domestica (in U.S.A.), bionomics of, 420; damaging rayon, 401; baits for, 406; toxicity of methyl bromide to,

thermophila, Zenillia (Carcelia). Theronia atalantae, parasite of Dendrolimus in Sakhalin, 95.

Theronini (see Pimplini).

Thiocyanates, in emulsions against larvae of Anomala, 111; against mites, 108, 109, 249; against Pseudococcus comstocki, against silverfish, 56; tests of toxicity of, to insects, 19, 434; formulae containing, 108, 111, 424; compatible with dinitro-ocyclohexylphenol, 249; and oil emulsions, 73, 108, 109; atomised sprays of oils containing, 56, 434; and pyrethrum, 56.

Thiodiphenylamine, spraying with, against Cydia pomonella, 22, 129, 206, 411, 550; addition of larval attractants to sprays of, 129; against other Tortricids, 425, 641; in oils for treating maize ears against Heliothis, 641; not affecting midgut wall of Laphygma, 481; in dusts and sprays INDEX. 805

against *Rhagoletis* spp., **196**, **561**; particle size affecting toxicity of to honey bees, **550**; dispersing agents for sprays of, **112**, **113**; effects of fruit growth and weather on deposits of, **206**.

Thiourea, insecticidal action of, 408. Thiuram Disulphide, Tetramethyl, spraying with, against Coleoptera,

595, 650.

Thomasiniana theobaldi, bionomics of, on raspberry in Britain, 283.

thompsoni, Perkinsiella.

thoracicus, Macrocentrus; Triaspis. Thosea, new parasite of, in Java, 514.

Thosea cervina, parasites of, in Ceylon, 174, 514, 620; on tea, 174.

Thosea recta, parasites of, on tea in Ceylon, 174.

thoseae, Metaplectrus.

Thrinax parviflora, Brontispa longissima on, in New Guinea Territory, 379.

thripoborus, Orius (Triphleps).

Thrips fuscipennis (Rose Thrips), on rose and strawberry in Britain, 145, 146; pyrethrum against, 146.

Thrips imaginis, measures against, on tomato in Victoria, 31.

Thrips tabaci, in Britain, 146; bionomics of, in France, 24, 25; in Hawaii, 35, 139, 140; in U.S.A., 121, 413, 476, 641; on greenhouse carnations, 413, 641; on vegetables, 25, 121, 146, 476; host-range and distribution of virus disease transmitted by, 24, 25, 139–141; not transmitting virus disease of papaya; 35; measures against, 25, 121, 146, 413, 641.

Thrips Banana Rust (see Scirto-thrips signipennis).

Thrips, Blueberry (see Frankliniella vaccinii).

Thrips, Citrus (see Scivtothrips aurantii and S. citri).

Thrips, Gladiolus (see Taeniothrips simplex).

Thrips, Onion (see Thrips tabaci). Thrips, Pear (see Taeniothrips inconsequens).

Thrips, Rose (see Thrips fusci-

pennis).

Thrips, Tobacco (see Frankliniella fusca).

thurberiae, Apanteles.

Thyanta custator, on beet in Arizona, 429.

Thymus serpyllum, propagation of, to attract Scoliids in Russia, 572.

Thyraeella collaris, establishment of, against Plutella maculipennis in New Zealand, 64, 333; oviposition habits and relation to other parasites of, 64, 65.

Thyridia themisto, bionomics and control of, on Brunfelsia hopeana

in Brazil, 292.

Thysanoptera, relation of, to plant diseases, 24, 25, 35, 139, 140, 141, 532, 542, 608; natural enemies of, 95, 210, 320; predacious species of, 175, 210, 211, 273; of Argentina, 655; of India, 480; classification and new species of, 272.

Thysanus, parasite of Tetracnemus peregrinus in California, **355.**

Thysanus elongatus, parasite of Pseudococcus comstocki in Virginia, 196.

Thysanus fax, parasite of Chrysom-phalus ficus in Brazil, 625.

Thysanus niger, parasite of Pseudococcus comstocki in Virginia, 196. Thysanus townsendi, parasite of Chrysomphalus ficus in Brazil, 625.

tibialis, Phanerotoma; Reticulitermes.

Tiliacora, Othreis fullonia on, in Gold Coast, 107.

Timber, Coleoptera in, 78, 86, 103, 139, 146, 147, 150, 178, 233, 234, 275, 284, 315, 316, 324, 333, 345, 385, 392, 427, 447, 448, 466, 467, 497, 593; digestive enzymes of Coleopterous larvae boring in, 340; Siricids in, 385; termites in, 6, 7, 19, 138, 142, 157, 333, 400, 473, 483, 495, 496, 537, 610, 611.

Timbo, dusting with, against insects, 252, 405, 408, 630; rotenone contents of, 252, 408, 630.

Timothy Grass (see *Phleum pratense*).

tineivorus, Typhlodromus.

Tineola biselliella, bionomics of, 375; passage of small openings by larvae of, 216.

Tiphia (parasite of Lamellicorn larvae), in Canada, 90, 509; in Kentucky, 299; proposed introduction of, into Mauritius against Lachnosterna smithi, 84.

Tiphia femorata, parasite of Lamellicorn larvae in Russia, 572.

Tiphia intermedia, parasite of Lamellicorn larvae in Kentucky, 299.

Tiphia popilliavora, utilisation of, against Popillia japonica in U.S.A., 161, 426, 430; strain of, from Korea, 430.

Tiphia vernalis, utilisation of, against Popillia japonica in U.S.A., 161, 426, 504, 505,

Tipula, damaging cereals and pastures in Germany, 68.

Tipula czizeki, in Germany, 68. Tipula oleracea, in Germany, 68. Tipula pagana, in Germany, 68.

Tipula paludosa, study of populations of, in grassland in Br. Isles, 524; in Germany, 68.

Tipula vernalis, in Germany, 68.
Tipulid Larvae (in Britain), measures against, on strawberry, 283; damaging grassland, 381.
Tirathaba, parasite of, in Java, 620.

Tirathaba rufivena, parasite of, in Malaya, 620.

tischeriae, Apanteles.

Titanio normalis, on Convolvulus in Russian Union, 478, 479; larva of, 478; confused with Loxostege sticticalis, 479.

titanus, Strategus.

Tits, destroying Rhabdophaga heterobia, 312.

Tmetocera (see Spilonota).

Tmolus echion, derris against, on pineapple in Trinidad, 373.

Toads (see Bufo marinus).

Tobacco, pests of, in S. Africa, 19, 491; pests of, in S. America, 512, 608; pests of, in Fiji, 390; Engytatus geniculatus on, Hawaii, 34: Hawaii, 34; pests of, in India, 20, 183, 266, 523; pests of, in Netherlands Indies, 318, 319, 492; pests of, in Mauritius, 610; pests of, in Nyasaland, 153, 154, 514, 588; Protoparce quinquemaculata on, in Ontario, 303; Myzus persicae on, in S. Rhodesia, 84, 184; pests of, in U.S.A., 11, 200, 201, 218, 253, 363, 365, 404, 428, 430, 476, 482; insects and virus diseases in, 25, 27, 57, 84, 140, 141, 154, 166, 179, 180, 183, 184, 318, 319, 438, 491, 542, 608.

Tobacco (Stored), Ephestia spp. in, 19, 362, 365, 429; E. elutella

killed by residues of lead arsenate on, 365; Lasioderma serricorne in, 362, 390, 493, 610.

Tobacco (as an insecticide), tests of duration of activity of sprays of, 233; as a carrier for dusts, 46, 301, 631; steam distillates of, in baits for locusts, 612; fixed extract of, 233.

Tobacco Aphis (see Myzus persicae). Tobacco Flea-beetle (see Epitrix

parvula).

Tobacco Hornworm (see Proto-parce).

Tobacco Moth (see Ephestia elutella). Tobacco Thrips (see Frankliniella fusca).

Tomaspis saccharina, pyrethrum dusting against, on sugar-cane in Trinidad, 52.

Tomaspis sanguinea, type of injury to apple and pear by, in Britain, 282.

Tomato, Frankliniella schultzei on, in S. Africa, 491; pests of, in Argentina, 141, 608; pests of, in Australia, 30-32, 143, 335, 466; Gnorimoschema operculella on, in Bermuda, 138; pests of, in greenhouses in Britain, 145, 146; Pseudococcus citri on roots of, in Egypt, 30; pests of, in Fiji, 148, 279, 390; Thysanoptera on, in France, 24, 25; pests of, in Hawaii, 34, 139, 273, 650, 651; pests of, in India, 183, 266, 271, 523, 542; Acanthocoris sordidus on, in Japan, 171; pests of, in U.S.A., 21, 121, 122, 134, 162, 200, 201, 207, 218, 276, 278, 429, 481, 534, 535, 536, 565, 598, 603, 631, 640, 641; Paratrioza cockerelli causing psyllid yellows in, 21; insects and virus diseases in, 24, 34, 57, 58, 134, 139-141, 183, 438, 491, 542, 608, 631; introduction of selenium into, against mites, 207; soil insecticides causing injury to, 122, 145.

Tomato (Fruit), fumigation of, against Trypetids, 499; problem of lead arsenate residues on, 32; rendered unfit for canning by sulphur dusts, 22.

Tomato Fruit Worm (see Heliothis armigera).

Tomato Hornworm (see Protoparce sexta).

Tomato Psyllid (see Paratrioza cockerelli).
tomentosus, Byturus.

Tonesia bondari, sp. n., on Cocos | spp. in Brazil, 468.

Tonga Islands, Nezara viridula in, 170.

topiarius, Phrydiuchus.

Torocampus eques, parasite of Neodiprion sertifer in Latvia, 86.

toroneura, Leucania.

Torrubia fragrans, Prepodes quinquedecimpunctatus on, in Porto Rico, 473. Tortrix, keys to Russian species of,

479.

Tortrix argyrospila, on apple in U.S.A., 2, 127, 128, 190, 195; new parasite of, 190; measures against, 127, 128; technique for testing ovicides against, 195.

Tortrix cerasivorana (Cherry Tent Caterpillar), parasites of,

Quebec, 625, 626.

Tortrix dinota, new parasite of, in Kenya, 514; on tea in Nyasaland, 60.

Tortrix excessana, on apple in New Zealand, 151.

Tortrix forskaleana, on maple in Britain, 628; in New York,

Tortrix occidentalis, new parasite of, in Nyasaland, 514.

Tortrix postvittana, distribution of, in Australia, 150-152; in New Zealand, 150, 151; parasites and biological control of, 151, 152.

measures pronubana, Tortrix against, in greenhouses in Britain,

146. Tortrix rosana, new parasite of, in Bulgaria, 514.

Tortrix sarcostega, bionomics of, in Baluchistan, 518.

Tortrix unifasciana, on privet in Britain and New York, 628.

Tortrix viridana, parasite of, Britain, 80; on oak in Latvia, 86. torvina, Spilochalcis (see S. side).

Torymus cyanimus, search for hosts and egg distribution of, 490.

Toumeyella numismatica, measures against, on Pinus banksiana in Michigan, 605.

townsendi, Drosicha; Thysanus. Toxicarol, determination of,

derris, 29, 30.

Toxoptera aurantii, on Citrus in Brazil, 168; unlikely to transmit phloem necrosis of tea in Ceylon, 175.

Toxoptera graminum (on cereals), in Argentina, 36; in India, 266; in U.S.A., 251, 254, 356; outbreak of, 251; natural enemies of, 36, 251; aeroplane survey of damage by, **356.**

Toxoptera piricola, bionomics and control of, on pear in Japan, 96. Trachelus tabidus, cultural practices affecting damage to wheat by, in

U.S.A., 254. traducalis, Pagyda. transiens, Creatonotus. translucens, Avrosia. transvaalensis, Termes.

trapezicollis, Ancylorrhynchus.

Traps, mechanical type of, for studying migrations of Aphids, construction and use of, for Popillia japonica, 125, 126, **504, 595**; mobile type of, for Meligethes aeneus, 231; for other Coleoptera, 9, 50, 255, 571, 593; for clothes moths, 255; Trypetids, 54, 183, 461, 462, 563; (Hopperdozer), for Pyrilla perpusilla, 82.

Treehoppers, Coconut (see Sexava). Trenches, against armyworms, 300; against beet weevils, 577, 578; against Schistocerca paranensis,

498.

Trialeurodes vaporariorum, Britain, 145, 374; in Canada, 310; on tomatos in Victoria, 32; parasitised by Encarsia formosa, 374; utilisation of E. formosa against, 32, 310; other measures against, 32, 145; tests of fumigants on, 132.

triangularis, Acerophagoides.

Triaspis, parasite of Chalcodermus aeneus in U.S.A., 405.

Triaspis curculionis var. rufus, parasite of Chalcodermus aeneus in U.S.A., 405.

Triaspis thoracicus, proposed introduction of; into W. Australia against Bruchus pisorum, 142; introduced into U.S.A. against B. brachialis, 48, 430; attempted rearing of, on B. obtectus, 48.

Triaspis vestiticida, parasite Anthonomus vestitus in Peru,

510.

castaneum, Tribolium measures against, in stored cereals in U.S.A. and Australia, 209, 210, 380; in imported ground-nuts in Fiji, 496; in stored cacao, 616. confusum, measures Tribolium

against, in stored wheat in Australia, 380; in U.S.A., 255, 477, 506; effects of high temperatures on, 187, 506; action of fumigants on, 327, 434, 452, 455, 477; caught on treated flypapers, 255; studies of physiology and populations of, 450, 452, 506, 592.

Tribulus terrestris, preferred by Schistocerca gregaria in India, 519.

Trichacis remulus, parasite of Mayetiola destructor in Russia, 618, 619; bionomics of, 619.

Trichlorethylene, for obtaining extracts of Tephrosia, 586.

Trichogramma, 77; forms and utilisation factors affecting of, against Diatraea saccharalis in Louisiana, 160, 601; species of, introduced into Lousiana, 601; importation of, into Mauritius against Diatraea, 610.

Trichogramma australicum, breeding and utilisation of, in For-

mosa, 92.

Trichogramma dendrolimi, parasite of Dendrolimus in Sakhalin, 95.

Trichogramma euproctidis, importation of, against Cydia molesta in U.S.A., 295.

Trichogramma evanescens, 221; bionomics, races and utilisation of, in Russian Union, 581, 582; effect of different hosts on, 340.

Trichogramma intermedium, hosts

of, in India, 183.
Trichogramma luteum, parasite of Argyroploce leucotreta in S. Africa, 228.

Trichogramma minutum (parasite of Lepidopterous eggs), in Barbados, **3**, **160**; in Colombia, **9**; in India, 184, 264, 265; in Japan and Korea, 372; imported into Spain, 457; in U.S.A., 167, 192, 193, 534, 600, 602; bionomics of 264, 553; field utilisation of, 3, 9, 160, 184, 192, 193, 534; reared on eggs of Sitotroga cerealella, 9, 192, 193, 457, 553.

Trichogramma pallida, in Russian Union, 581; utilisation of, against Cydia spp., 581.

Trichogramma pini, in Russian Union, 581.

Trichogramma pretiosum, utilisation and value of, against Cydia molesta in Connecticut, 87.

Trichogramma turkestanica, Russian Union, 581; effects of temperature and humidity on, 581.

Trichogrammatoidea nana, parasitising Hapalia machaeralis Burma, 77; bionomics of, parasitising lac predators, etc., in India, 264, 265.

Tricholyga sorbillans, bionómics of,

in Ceylon, 174.

Trichophaga tapetzella, baits for, in U.S.A., 255.

Trichopoda pennipes, bionomics of, parasitising Anasa tristis U.S.A., 120, 442-444; attempted introduction of, into Australia and Fiji against Nezara, 170, 496; effect on A. tristis of parasitism by, 120, 443.

Trichopria, parasite of Metagoni-stylum in Brazil, 470.

Trichopsidea clausa, parasite of grasshoppers in Canada, 308. Trichosanthes anguina, Plusia pep-

onis on, in India, 264.

Trichospilus pupivora, in Ceylon, 594, 610, 619, 620, 621; in India, Java and Malaya, 620; introduction of, into Mauritius, 610; utilisation and value of, against Nephantis serinopa, 619, 620, 621; other hosts of, 610, 620, 621; bionomics and laboratory breeding of, 620, 621.

tricolor, Exenterus.

Tricresylphosphate, for treating papers, etc., against Ctenolepisma, 56.

tridentata, Eutetrapha (Saperda). Triecphora vulnerata (see Tomaspis sanguinea).

Triethanolamine, and oleic acid, in preparation of inverted spray mixtures, 163.

trifasciatus, Coccophagus.

Trifolium alexandrinum, pests of, in India, 265, 266, 523.

Trifolium incarnatum, experiments with Jassid and virus disease in, 500.

Trifolium subterraneum, susceptibility of strains of, to Halotydeus destructor in W. Australia, 142.

trifurcata, Ceratoma.

Trigonophymus, in Argentina, 308. 511; introduced parasite of, 308; damaging lucerne, 511.

Trigonophymus arrogans, bionomics of, in Argentina, 497, 511.

Trigonophymus elongatus, in Argentina, **497, 511.**

Trigonophymus pratensis, in Argentina, 497.

trima, Orthocraspeda.

. trimaculata, Macropsis.

Trinidad, pests of Citrus in, 471, miscellaneous pests in, 373, 616; sugar-cane pests in, 52, 471; strain of Metagonistylum minense introduced into, 471; beneficial insects introduced into other islands from, 263, 471.

trinitatis, Azya.

Trionymus masrensis, measures against, on roots of barley in Egypt, 30.

Trionymus sacchari, on sugar-cane in Dominican Republic, 498.

Trioxys aceris, parasite of Myzocallis annulata in Tasmania, 142. Triphleps (see Orius).

Tripoli Earth, as a carrier for bacterial spores, 583.

Tripsacum, immune from Aplano-

bacter stewarti, 245. Tripterygium wilfordi, tests of insecticidal properties of, 241, 242.

trivadiatus, Eriophyes.

Trissolcus murgantiae, parasite of Murgantia histrionica in N. Carolina, 560.

Trissolcus simoni, parasite of Mayetiola destructor in Russia, 618.

Trissomalus fulvicornis (see Gonio-

Tristania conferta, Phylacteophaga eucalypti on, in New South Wales, 237.

tristigmus, Euschistus.

tristis, Ambeodontus; Anasa; Heteronychus; Lachnosterna (Phyllophaga).

tritici, Contarinia; Euxoa (Agrotis); Frankliniella; Harmolita. Triticum repens (see Agropyrum).

Tritneptis klugi, parasite and hyperparasite of Neodiprion tsugae in

Oregon, 552. Triumfetta rhomboidea, weevils on, in India, 270, 446, 521, 522.

trivittata, Diabrotica.

Trochalus, on tea in Nyasland, 154.

Trochorrhophalus dipterocarpi, in Swintonia timber in Burma, 78. Trogodendron fasciculatum, nomics of, in Australia, 608.

Trogoderma versicolor, bionomics and control of, in Central Asia,

Trogus lutorius f. jezoensis, parasite of Dendrolimus in Sakhalin, 95. Troilus luridus, doubtful value of, against Leptinotarsa decemlineata in Germany, 70.

Tropaeolum, Thrips tabaci on, in France, 25.

tropicalis, Myzus (see M. varians). Trout, insect food of, 224, 335.

truncatulus, Hypolixus (Lixus). tryoni, Dacus (Chaetodacus, Strumeta) ferrugineus.

trypetae, Habrocytus.

trypetes, Chilo.

Trypetids (Fruit-flies), bibliography of, 280.

Tryphetus incarnatus, bionomics and control of, on Pithecolobium lobatum in Java, 182.

Trypodendron betulae, fungi associated with, on birch in Minne-

sota, 165.

Trypodendron (Xyloterus) lineatum, bionomics of, on spruce in Sakhalin, 96.

Trypodendron retusum, fungi associated with, on aspen in Minnesota, 165.

tschirchii, Coccophagus.

Tsuga heterophylla, Polyphylla perversa on, in Br. Columbia, 509; Neodiprion tsugae on, in Oregon, 552.

tsugae, Exenterus; Microcryptus (Aptesis); Neodiprion.

tuberculata, Oncocephala.

tuberculatus, Coccosterphus. tumiceps, Reticulitermes.

tumida, Aethina.

tumulosus, Ligyrus. Tung Tree (see Aleurites).

Tunisia, Hypera variabilis 403.

Turf, pests of, in S. Africa, 614, 615; insects damaging, in Australia, 237, 466, 548; Lachnosterna damaging, in Canada, 90; Melolonthids damaging, France, 24, 83; insects damaging, in U.S.A., 124, 125, 161, 214, 408; insects damaging, in W. Indies, 217, 373, 374; treatment of, against pests, 90, 373, 408, 548, 614, 615.

turionana, Rhyacionia (Evetria). turkestanica, Trichogramma.

Turkey, miscellaneous pests 288.

Turnip, susceptibility of, to Lachnosterna in Canada, 90; Galeruca tanaceti on, in Germany, 68; pests of, in U.S.A., 569, 596; Listroderes obliquus on, in Victoria, 31.

Turnip Aphis (see Rhopalosiphum pseudobrassicae).

Turpentine, as a preservative for timber against Lyctus, 147.

Twelve-spotted Asparagus Beetle (see Crioceris duodecimpunctata). Twig Girdlers (see Oncideres).

Tyloderma fragariae, measures against, on strawberry in Kentucky, 362, 554.

tylodermatis, Eurytoma.

Tyora sterculiae, on Brachychiton populneus in New South Wales, 467.

Typhlocyba debilis, doubtful importance of, on apple in Britain,

Typhlocyba froggatti, on apple in Argentina, 496; on fruit trees and Crataegus in Britain, 284; bionomics and biological control of, in Tasmania, 291; other measures against, 291, 496, 497.

Typhlocyba pomaria, feeding habits of, and types of injury caused to rose and apple by, in N. America,

256, 548.

Typhlocyba rosae, doubtful importance of, on apple in Britain, 284. Typhlodromus spp., attacking Tineola, Dermestes and Anthrenus in Switzerland, 376.

typica, Stephanitis.

Typophorus, on potato in Argentina, 512.

Typophorus nigritus, 512.

Tyria jacobaeae, Harpobittacus preventing establishment of, against Senecio jacobaea in Australia, 143. Tyroglyphus spp., measures against, in stored grain in Russian Union, 576.

U.

Ufens, parasite of Diaprepes in St. Lucia, 262.

Uganda, coffee pests in, 80, 175, 229, 230, 341; cotton pests in, 175, 228, 310; locusts in, 175, 228, 229, 388, 491; termites of, 590; new weevil in, 341; beneficial insects in, 80, 229, 230, 385, 386, 514; book on agriculture and insect pests in, 175; introduction of parasites of Saissetia oleae into California from, 385, 386.

uhleri, Horistonotus.

Ulex europaeus, utilisation of Apion ulicis against, in New Zealand and Tasmania, 334.

ulicis, Apion.

ulkei, Moneilema. ulmi, Lepidosaphes.

ulmi, auct., Oligonychus (see Paratetranychus pilosus).

Ultroil, as a wetting agent, 595. umbrosus, Dacus (Chaetodacus). Uncaria gambir, pests of, in Suma-

tra, 105, 106.

uncariae, Sura. uncinata, Bostrychopsis.

undalis, Hellula.

undatus, Brachycerus.

undecimpunctata, Coccinella.

unica, Heydenia.

unicolor, Hoplorrhinus; Neoclytus. unicostata, Eufallia.

unifasciana, Tortrix. uniformis, Sesamia.

Cirphis (Leucania, unipuncta, Sideridis).

unipunctata, Cryptophasa (Maroga). uniseta, Voriella.

unispina, Gryllotalpa. United States of America, asparagus beetles in, 48, 430; beet pests in 161, 429; cereal pests in, 48, 165, 187, 217, 244, 250, 254, 299, 300, 366, 367, 428; cotton pests in, 257, 404, 428, 430, 500; pests of forest and shade trees in, 37, 40, 48, 49, 118, 119, 212, 213, 216, 220, 293, 294, 347, 348, 364, 370, 371, 395, 396, 416, 422, 424, 426, 427, 428, 438, 531, 532, 550, 551, 552, 606, 607; pests of leguminous crops in, 47, 48, 114, 353. 428, 429, 430, 431, 503, 567; pests of nut trees in, 116, 117, 159, 213, 405, 426, 549; orchard pests in, 48, 71, 127, 128, 129, 130, 160, 213, 294, 348, 360, 362, 373, 403, 409, 425, 549, 550, 557, 647; pests of small fruits in, 45, 287, 549; pests of ornamental plants in, 204, 214, 257, 532, 555, 606, 636; pests of solanaceous plants in, 11, 428, 429, 430, 501; vine pests in, 139, 425; storedproduct and household pests in. 137, 209, 255, 401, 420, 426, 429, 477, 506, 536, 605; Anabrus simplex in, 48, 187, 223, 326, 365, 428, 537, 562; grasshoppers in, 48, 144, 224, 254, 428, 603, 604; Lyctus in timber in, 447, 448; Myzus of, 45; Pantomorus Myzus of, 45; Pantomorus spp. in, 208, 370, 428, 494, 568, 608, 632; Popillia japonica in, 118, 123, 124, 125, 126, 364, 368, 369, 370, 426, 431, 594, 595; Rhopalosiphum pseudobrassicae in, 596; insects and plant diseases in, 40, 244, 256, 287, 364, 403, 502, 503, 532, 550, 606; beneficial insects and biological control in, 48, 114, 120, 129, 130, 137, 220, 293, 294, 300, 362, 373, 376, 405, 406, 416, 424, 425, 426, 427, 430, 431, 439, 549, 552, 569; Nematode parasites and diseases of P. japonica and their utilisation in, 124, 125, 368, 369, 370, 426; plant pest legislation in, 444, 498, 499, 568; restrictions on importation of bananas from Fiji into, 495; list of pests intercepted in quarantine in, 348; data on insect pests obtained by quarantine inspections in, 133; influence of cultural practices on pests in, 204, 254; standardisation in pest control in, 358; list of popular names of insects in, 280: directory of pest control in, patents relating to pest control in, 592; Macrodactylus subspinosus intercepted in Bermuda from, 139; Diarthronomyia chrysanthemi imported into Ireland from, 63; beneficial insects introduced into other countries from, 26, 143, 457. (See also under the various States.)

ununguis, Paratetranychus.

Urbanus proteus, on beans in Florida, 477.

Urena lobata, Pempherulus affinis on, in Madras, 521.

Urentius echinus, not transmitting virus disease of egg-plant in Madras, 542.

urichi, Liothrips.

Urostylis punctigera, food-plants and parasite of, in India, 74, 75. urozonus, Eupelmus.

urticana, Argyroploce.

Uruguay, summary of plant quarantines in, 499.

usipetes, Telenomus. usitatus, Taeniothrips.

Utah, beet pests in, 533, 563, 631;
Eutetix tenellus and curly top in
beet and tomato in, 631; Cydia
pomonella on cherry in, 48; miscellaneous pests in, 49, 538, 630,
631; termites in, 400; Spilochalcis leptis in, 190; birds
destroying noxious insects in,
217, 563; Necremnus leucarthros
imported from Europe into,
569.

utilis, Lecaniobius; Pseudaphycus; Smicronyx.

V.

vaccinii, Eriophyes; Frankliniella; Mineola.

Vaccinium (see Blueberry, Cranberry and Whortleberry).

Vacuum Fumigation, 222, 536, 568, 605.

vaga, Blattella.

INDEX.

vagans, Acarapis; Ceraphron (Calliceras).

vaginicola, Harmolita.

Valanga nigricornis, damaging tobacco in Sumatra, 318.

Valota insularis, predacious thrips on, in Hawaii, 272.

vanderwulpi, Cadurcia.

Vanessa cardui, on Xanthium spinosum in S. Africa, 19.

Vanilla, Arctiid on, in Porto Rico, 469.

Vanilla Essence, in baits for Trypetids, **91**, **465**.

Vapo-Fume Liquid, 250. vaporariorum, Trialeurodes.

variabilis, Hypera; Scyllinops. varians, Cyrtopeltis (see Engytatus

geniculatus); Myzus. varicornis, Monolepta; Paralito-

mastix. variegana, Argyroploce. variegata, Adonia.

variegatu, Adonia. variegatum, Atoposoma. variegatus, Brachylybas.

varinervis, Antholcus.

variolarius, Euschistus. variolosa, Polyphylla.

variolosum, Asterolecanium.

varivestis, Epilachna. vassilievi Microphanurus.

vassilievi, Microphanurus.

Vateria indica, new Calandra in seeds of, in India, 79.

vateriae, Calandra.

Vatsol, used with dichlorethyl ether for treating soil against thrips, 261.

Vatsol O.S., as a wetting agent for dusts, 250, 504, 600.

Vatsol O.T., improving fixed nicotine sprays against Cydia pomo-

nella, 160.

Vedalia cardinalis (see Rodolia).

Vegetable Marrow, Epilachna vigintioctopunctata on, in Fiji, **390.** vellicatus, Exenterus.

Velvet Beans (see Stizolobium). Velvet-bean Caterpillar (see Anticarsia gemmatalis). Venezuela, pests of stored coffee and cacao in, 292, 616.

venosata, Diatraea. ventralis, Exochomus. ventricosus, Pediculoides.

Verania discolor, bionomics of, in Formosa, 95.

Veratrum viride, insecticidal properties of, 352.

Verdol, 190.

Vermont, outbreak, disease and parasites of *Gilpinia polytoma* on spruce in, **37**, **422**, **424**, **427**, **531**, **637**, **638**; *Lymantria dispar* in, **428**.

vernalis, Tiphia; Tipula. Vernonia cinerea, Bemisia tabaci transmitting virus disease to tobacco from, in India, 183.

versicolor, Dindymus; Trogoderma. vesicularis, Eupelmella.

Vespa vulgaris, parasitised by Myopa in Europe, 474.

vestigialis, Agrotis. vestiticida, Triaspis. vestitus, Anthonomus.

Vetch Bruchid (see Bruchus brachialis).

Vetches (Vicia), pests of, in U.S.A., 47, 48, 404, 415, 418, 430, 598; quarantine regulations dealing with seed of, against Bruchids in U.S.A., 47; susceptibility of species and varieties of, to Macrosiphum onobrychis, 404; scorched by calcium cyanamide, 418.

vetula, Anomala. vetusta, Agrotis. Vicia (see Vetches). Vicia faba (see Beans). vicinus, Scapteriscus.

Victoria, Iridomyrmex humilis in, 332; miscellaneous pests in, 436, 494, 547; orchard pests in, 141, 149, 151, 216, 279, 335; vegetable pests in, 30-32, 142, 143, 290, 436, 546; beneficial insects and biological control in, 32, 141, 142, 545, 546.

viduatus, Aspongopus. vigintioctomaculata, Epilachna. Vigna (see Cowpeas).

villosa, Lagria.

Vine, Grape, pests of, in S. Africa, 19, 320; Pseudococcus adonidum on, in Australia, 355; Phylloxera on, in Bermuda, 139; Oides decempunctata on, in China, 446; pests of, in France, 83, 111; Phylloxera on, in Germany, 231; pests of, in Guam, 331, 332;

new Arthropods on, in India, 265, 271; Drosophila suzukii on, in Japan, 622; pests of, in New Zealand, 58, 333; Theresimma ampelophaga on, in Palestine, 623; pests of, in Russian Union, 569, 578-580; Ceratitis capitata on, in Spain, 462; pests of, in U.S.A., 2, 117, 133, 139, 210, 425, 536, 558; factors affecting resistance of varieties and hybrids of, to Phylloxera, 578, 579; problems of spray residues on fruit of, 117, 425.

Vine Bud Moth (see Theresimima ampelophaga).

Vine Moth, Australian (see Phalaenoides glycinae).

Vinegar, in baits for Ceratitis capitata, 462, 464.

vinitor, Nysius.

Viola papilionacea, Fenusa on, in Massachusetts, 418.

violaceum, Callidium. violaceus, Teleopterus.

Violet (see Viola papilionacea). virescens, Heliothis; Oecophylla smaragdina.

virgata, Epilachna; Ferrisiana (Ferrisia, Pseudococcus).

Virginia, Lepidoptera on maize in, 48, 367; orchard pests in, 48, 134, 195, 299, 300, 425, 500, 535; Macrosiphum onobrychis on peas in, 249, 504; other vegetable pests in, 46, 121, 122, 152, 503; pests of pines in, 49, 87, 116, 211; new mite on strawberry in, 530; pests of field and stored tobacco in, 200, 362, 365, 429; natural enemies and biological control of insect pests in, 87, 116, 196, 211, 400, 425, 500, 535.

Virginia, West, Scolytids and associated fungi on elm in, 116, 551; orchard pests in, 134, 363, 425, 562; beneficial insects and biological control in, 363, 425.

Virginia Creeper (see Parthenocissus quinquefolia).

virginica, Diacrisia. viridana, Tortrix.

viridiceps, Euplectromorpha. viridifasciata, Chortophaga.

viridis, Coccus.

viridiscutatus, Lygus hesperus. viridissima, Tettigonia.

viridula, Nezara.

Virola merendonis, fungous decay favouring infestation of timber of, by Xestobium, 315.

Virus Diseases (in Plants*), 166, 650; relations of Aphids to, 16, 23, 24, 26, 27, 35, 45, 57, 65, 66, 84, 112, 138, 140, 149, 154, 165, 175, 179, 180, 184, 188, 235, 256, 287, 302, 317, 329, 375, 415, 457, 501, 503, 523, 532, 542, 608, 638; relation of Bemisia to, 183, 318; relations of Delphacids to, 13, 16, 469; relations of Galerucids to, 501, 503; relations of Jassids to, 13, 15, 16, 35, 66, 75, 112, 134, 175, 256, 272, 287, 301, 302, 306, 500, 501, 502, 503, 510, 511, 532, 542, 548, 608, 617, 618, 631, 646; relations of Thysanoptera to, 24, 25, 35, 139, 140, 141, 491, 532, 542, 608; other suspected insect vectors of, 34, 75, 175, 382, 510; negative results with other insects and, 35, 256, 317, 382, 532, 542, 608; micromortar for studying, in insects, 349; mechanical transmission of, 25, 57, 140, 503, 542, 608; problems of quarantines against, 403; reviews of books on, 239, 437.

viteana, Polychrosis. vitellinae, Phyllodecta. vitiensis, Megatrioza. vitifoliae, Phylloxera. vitis, Eriophyes.

vittata, Diabrotica (see D. melano-Spilochalcis (see S. cephala); igneoides).

vittatus, Pantophthalmus.

vivida, Parasa.

Volck Oil, 155, 606.

voracis, Laelius.

vorax, Anthrenus.

Voriella uniseta, parasite of Tortrix postvittana in Tasmania, 151; suitable for introduction into New Zealand, 152.

vulgaris, Cyrtogaster; Stylocryptus; Termes; Vespa; Systoechus; Zenillia (Phryxe).

vulgatissima, Phyllodecta.

vulnerata, Triecphora (see Tomaspis

sanguinea).

vulnerator, Pristomerus vulnificus, Spathius. vuteria, Sesamia.

W.

W.S.C. Dynamite, 371. wahlbergi, Chilocorus Wallaba Wood (see Eperua falcata). Walnut (Juglans), Xyleborus dispar on, in Br. Columbia, 328; pests of, in U.S.A., 116, 211, 222, 549, 561, 564; tent fumigation of, 564; insecticides causing injury to, 561.

Walnut, Black (see Juglans nigra). Walnut Aphis (see Chromaphis juglandicola).

Walnut Caterpillar (see Datana integerrima).

Walnut Husk Fly (see Rhagoletis suavis completa).

Walnut-shell Flour, as a carrier for dusts, 249, 296, 640.

Washington, Thysanoptera on carnations in, 413, 641; pests of conifers in, 49, 371, 427; Cydia latiferreana on filberts in, 405, 406; pests of leguminous crops in, 48, 376, 639, 640; Heliothis armigera on maize in, 419, 640, 641; miscellaneous pests in, 506, 640-642; orchard pests in, 48, 222, 330, 411, 418, 425, 559, 639, 640, 641; beneficial insects in, 406, 569.

Washingtonia filifera, Dinapate wrighti on, in California, 137; Dinapate Brontispa longissima on, in New

Guinea Territory, 379. washiyai, Rhyacionia (Evetria). Wasmannia auropunctata, bionomics and control of, in Citrus groves in Florida, 475.

Water Snail (see Limnaea).

Water-gas Tar Oils, 134. Watercress, pests of, in Britain,

waterhousei, Epicauta.

Watermark Disease of Willows, 288. Watermelon, Itonida citrulli on, in Arizona, 596; Pseudococcus citri on roots of, in Egypt, 30; Engytatus geniculatus on, in Hawaii, 650; readily attacked by Schistocerca gregaria, 519.

Wattle (see Acacia).

Wax Beetle (see Platybolium alvearium).

Wax Scale (see Ceroplastes sinensis). Wax Scale, Pink (see Ceroplastes rubens).

West Indian Satinwood (see Zan-

thoxylum flavum).

West Indies, Cecidomyiids of, 468. (See also under the various Islands.)

westwoodi, Helopeltis.

^{*} These references are also indexed under the individual plants infected.

Wheat, pests of, in Canada, 112, 393; pests of, in Far East, 93, 100, 171; pests of, in India, 81, 266, 523; Diatraea saccharalis on, in Peru, 511; pests of, in Russian Union, 13, 14, 15, 16, 571, 573, 574, 576, 583, 617, 618, 619, 655; Lema melanopa on, in Spain, 463; pests of, in U.S.A., 205, 213, 215, 251, 254, 300, 356, 366, 505, 560, 603, 604; vectors of mosaic diseases of 13, 15, 16, 617, 618; varieties and hybrids of, in relation to pests, 215, 366, 393, 464, 505; harvesting machinery affecting pests of, 254; type of injury to, by Eurygaster, 574; treated with selenium against Tetranychus telarius, 276.

Wheat (Stored), pests in, and their control, 72, 209, 210, 254, 380, 429, 436, 453, 455, 457, 480, 506, 540, 576; olfactory reaction of *Rhizopertha dominica* to, 451; as bait for Tenebrionids, 571.

Wheat Jointworm (see Harmolita tritici).

Wheat Stem Sawfly (see Cephus cinctus).

Wheat Wireworm (see Agriotes mancus).

Whiteflies (Aleurodids), and plant diseases, 183, 318; ants associated with, 475; natural enemies of, 32, 293, 310, 374; of Br. Isles, 374; list of genera and subgenera of, 88; classification and new species of, 88, 374.

Whitefly, Cabbage (see Aleurodes proletella).

Whitefly, Greenhouse (see Trial-eurodes vaporariorum).

White-fringed Beetles (see Pantomorus leucoloma and P. peregrinus).

Whortleberry, Argyroploce urticana on, in Britain, 288.

Willow (Salix), Teragra guttifera on, in S. Africa, 18; Membracid on, in Brazil, 392; pests of, in Britain, 146, 288, 312; pests of, in Canada, 328, 347, 626; Acrocercops on, in Japan, 99; pests of, in U.S.A., 438, 595, 603, 607, 632; susceptibility of species of, to pests, 146, 438, 626; Xiphydria prolongata and bacterial disease of, 288; relation of

fungous decay to infestation of timber of, by Xestobium, 315, 316.

Willow Leaf Beetle, European (see Plagiodera versicolora).

Wilt Disease, in Gilpinia polytoma, 422, 637; in Lepidoptera, 61, 154, 314, 395, 588, 600, 650; effect of, on parasites of Pieris rapae, 650.

Wind, relation of grasshopper flights to, 428, 497, 604; relation of, to flight and spread of other insects, 50, 164, 180, 212, 213, 262, 271, 318, 329, 404, 606, 647; mites spread by, 281, 306, 330, 521.

Wine, Carpoglyphus lactis on, 488; in baits, 99, 465.

Wine Casks, vacuum fumigation of, against *Scobicia declivis*, **222**.

winnemana, Erixestus.

Winter Moth (see Operophtera brumata).

Winthemia alabamae, sp. n., parasite of Alabama argillacea in Brazil, 469.

Winthemia quadripustulata, parasite of Cirphis unipuncta in U.S.A., 300.

Wire Netting, against Saperda candida, 243; against Strategus quadrifoveatus, 472.

Wireworm, Sand (see Horistonotus uhleri).

Wireworm, Wheat (see Agriotes mancus).

Wireworms, measures against, 122, 145, 253, 282, 477, 483, 573, 643, 644, 645; summaries of data on bionomics and control of, 299, 381.

Wisconsin, miscellaneous pests in, 42, 48, 135, 361.

Wistaria, Dichomeris oceanis on, in Japan, 100; new subspecies of Lecanium persicae on, in New Zealand, 58.

Withania somnifera, Empoasca lybica on, in Sudan, 387.

woglumi, Aleurocanthus.

Wood Ash, as a carrier for calcium cyanide dust, 517.

Woodpeckers, destroying noxious insects, 78, 243, 244, 271.

Woollen Goods (see Textiles).

Woolly Apple Aphis (see Eriosoma lanigerum).

wrighti, Dinapate.

Wyoming, new weevil on *Pinus contorta* in, **371.**

X.

Xanthippus corallipes pantherinus, parasitised by Muscina stabulans in Texas, 477.

Xanthium pungens, Euaresta aequalis introduced into Australia against, 143.

Xanthium spinosum, Vanessa cardui on, in S. Africa, 19.

Xanthogramma aegyptium, predacious on Myzus persicae in S. Rhodesia, 184.

Xanthogramma scutellare, predacious on Aphis maidis in Guam, 331.

xanthomelaena, Galerucella (see G. luteola).

Xanthone, tests with, as an insecticide, 411, 412, 561, 640.

Xanthopimpla cera, parasite of Hapalia machaeralis in Burma, 77. Xanthopimpla punctata, parasite of

Nephantis serinopa in India, **620.** Xanthopimpla rhopaloceros, parasite of Tortrix postvittana in Tasmania, **151**; suitable for intro-

duction into New Zealand, 152. Xanthopimpla stemmator, parasite of Chilo zonellus in Ceylon, 610; importation of, against sugarcane pests in Mauritius, 610.

Xenocomys chrysomphali, gen. et sp. n., parasite of Chrysomphalus lahillei in Argentina, 344.

Xenocrepis bothynoderis (egg-parasite of Cleonus punctiventris), bionomics and factors affecting importance of, in Ukraine, 456.

Xenosternum ornigis, parasite of Parornix prunivorella in Kansas, 191.

xerophila, Neocoelostoma.

Xestobium rufovillosum, 233; measures and precautions against, in building timbers in Britain, 593; relation of fungous decay to infestation of timber by, 178, 315, 316.

Xiphydria prolongata, associated with bacterial disease of willow in Britain, 288.

Xyleborus, not infesting barked logs in Burma, 78; measures against, on avocado in Fiji, 170.

Xyleborus aplanatideclivis, on avocado and Inocarpus in Fiji, 170. Xyleborus compactus, on Clidemia hirta in Fiji, 170.

Xyleborus dispar, bionomics of, in Br. Columbia, 328.

Xyleborus fallax, in Swintonia timber in Burma, 78.

Xyleborus fornicatus, food-plant relations of subspecies of, in Ceylon, 174, 381.

Xyleborus fornicatus subsp. fornicatior, 381.

Xyleborus germanus, relation of, to Dutch elm disease in U.S.A., 365, 550; bionomics of, 551.

Xyleborus morstatti, on avocado in Fiji, 170.

Xyleborus mutilatus, on avocado in Fiji, 170.

Xyleborus perforans, on coconut in Fiji, 169.

Xyleborus rameus, on Endospermum in Fiji, 169.

Xyleborus submarginatus, in Swintonia timber in Burma, 78.

Xyleborus testaceus, on Hydnocarpus and Citrus in Fiji, 169, 389; measures against, 389.

Xyleutes ceramicus, bionomics of, on teak, etc., in Burma, 75, 76.

Xyleutes persona, bionomics and control of, on Cassia in Burma, 78.

Xylia dolabriformis, pests of, in Burma, 77, 78.

Xylol, permeability of insect cuticle by, 240.

Xylomyges eridania (see Laphygma). xyloni, Solenopsis geminata.

Xyloprista hexacantha, on Eucalyptus viminalis in Argentina, **512.** Xylopsocus castanoptera, on Poin-

ciana regia in Fiji, **170.** Xylosandrus (see Xyleborus). Xyloterus (see Trypodendron).

Xylothrips religiosus, plants attacked by, in Pacific Is., 170. Xylotrechus colonus, on elm in New York, 628.

Y.

Yam (see Dioscorea). Yam Beetle (see Galerucida bicolor). Yeast, in baits for Cydia pomonella, 414, 435.

ypsilon, Agrotis. Yucca, Aonidiella aurantii on, in Japan, 102.

Z.

Zacerata asparagi, on asparagus in S. Africa, 19.
Zadiprion grandis, parasite of, in Nebraska, 294.

Zagrammosoma, parasite of Lithocolletis in Porto Rico, 473.

Zagrammosoma multilineata, parasite of Parornix prunivorella in Kansas, 190.

Zanthoxylum spp., pests of, in Porto Rico, 51, 427.

Zanzibar, termites of, 590; parasites of Lamellicorns in, 83, 148.

Zarhopalus corvinus (parasite of Pseudococcus maritimus), introduced into Canada from California, 310.

Zatropis, parasite of Anthonomopsis mixtus in Florida, 475.

Zatropis incertus, parasite of Chalcodermus aeneus in U.S.A., 405.

zealandica, Odontria.

Zelleria haimbachi, new parasite of, in U.S.A., 190.

Zenillia, parasite of Pieris rapae in Britain, 314.

Zenillia blanda, parasite of Tortrix cerasivorana in Quebec, 626.

Zenillia nemea, parasite of Pieris rapae in Britain, 314.

Zenillia picta, parasite of Plecoptera reflexa in India, 74.

Zenillia thermophila, parasite of Plecoptera reflexa in India, 74.

Zenillia vulgaris, parasite of Lepidoptera in Britain, 314, 526.

Zethus rufinodus, damaging fence posts in Porto Rico, 473.

Zeugenia histrionica, in Swintonia timber in Burma, 78.

Zeuxia cinerea (parasite of Cleonus mendicus), bionomics of, in Italy, 316.

giardi (giardiana), Zeuxidiplosis attempted introduction of, into Australia against Hypericum perforatum, 143.

Zeuzera coffeae, on forest trees in Burma and Java, 78, 270, 513; on cacao, etc., in Malaya, 513; bionomics of, 270, 271, 513; measures against, 513.

Zinc, not affecting injury to Citrus

by sulphur, 296.

Zinc Arsenate, strawberry roots injured by mixture containing, 259; (basic), in sprays and dusts against Rhagoletis spp., 196, 561.

Zinc Arsenite, in sprays against Coleoptera on potato, 278, 639; strawberry roots injured by mixtures containing, 259.

Zinc Chloride, as a preservative for timber against Lyctus, 7, 147. termites and

Zinc Sulphate, uses and effects of, in sprays, 134, 196, 296, 633.

Zinc Sulphide, reducing efficiency sulphur against Scirtothrips of citri, 296.

zinckenella, Etiella.

Zinnia, Thrips tabaci transmitting spotted wilt to, in France, 25; new Jassid on, in India, 266.

Zinnia elegans, Bemisia tabaci transmitting leaf-curl to tobacco from, in India, 183.

Zizania latifolia, Agromyza oryzella on, in Japan, 98.

Zizyphus jujuba, pests of, in India, 263, 264, 271, 341; pests of, in Siam, 94; cultivation of lac on, 94, 263.

zonellus, Chilo.

Zootermopsis nevadensis, in Montana, 537.

Zootrephes, parasite of Gilpinia polytoma in Maine, 637.

zosine, Platygaster.

zygaenarum, Syntomosphyrum.

NOTICES.

Secretaries of Societies and Editors of Journals willing to exchange their publications with those of the Institute are requested to communicate with the Director.

The Annual Subscription, in advance, to the Review, Series A (Agricultural), is 30s. post free; Series B (Medical and Veterinary), 15s. post free. Subscriptions to the current Volume received after 30th June in each year will be charged at the published price, vix.: Series A, 36s.; Series B, 18s. Prices of Back Volumes on application.

Orders and Subscriptions should be sent direct to the Director, Imperial Institute of Entomology, 41, Queen's Gate, London, S.W.7, or through a bookseller.